

The following are the SPSS results of the Core Drug & Alcohol Survey conducted at IL

# Frequencies

## Notes

<b>Output Created</b>		16-JUN-2004 09:46:55
<b>Comments</b>		
<b>Input</b>	<b>Filter</b>	<none>
	<b>Weight</b>	<none>
	<b>Split File</b>	<none>
<b>N of Rows in Working Data File</b>		747
<b>Missing Value Handling</b>	<b>Definition of Missing</b>	User-defined missing values are treated as missing.
	<b>Cases Used</b>	Statistics are based on all cases with valid data.
<b>Syntax</b>		FREQUENCIES VARIABLES=isp1 isp2 isp3 isp4 isp5 .
<b>Resources</b>	<b>Total Values Allowed</b>	149796
	<b>Elapsed Time</b>	0:00:00.13

## Statistics

		Sp code row A	Sp code row B	Sp code row C	Sp code row D	Sp code row E
N	Valid	747	747	747	747	747
	Missing	0	0	0	0	0

# Frequency Table

## Sp code row A

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid	1	747	100.0	100.0

## Sp code row B

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid	5	747	100.0	100.0

## Sp code row C

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid	0	747	100.0	100.0

## Sp code row D

	Frequency	Percent	Valid Percent	Cumulative Percent

Valid	4	747	100.0	100.0	100.0
-------	---	-----	-------	-------	-------

Sp code row E

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid	3	747	100.0	100.0

# Crosstab analysis of categorical questions by gender

## Crosstabs

### Notes

<b>Output Created</b>		16-JUN-2004 09:46:55
<b>Comments</b>		
<b>Input</b>	<b>Filter</b>	<none>
	<b>Weight</b>	<none>
	<b>Split File</b>	<none>
	<b>N of Rows in Working Data File</b>	747
<b>Missing Value Handling</b>	<b>Definition of Missing</b>	User-defined missing values are treated as missing.
	<b>Cases Used</b>	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
<b>Syntax</b>	CROSSTABS /TABLES=q01 q03 q04 q06 q07 q08a q08b q08c q08d q08e q08f q08g q09 q10a q10b q11 q12a q12b q12c q12d q12e q13 q14 q16a q16b q16c q16d q16e q16f q16g q16h q16i q16j q16k q16l q17a q17b q17c q17d q17e q17f q17g q17h q17i q17j q17k q17l q18a q18b q18c q18d q18e q18f q18g q18h q18i q18j q18k q18l q19a q19b q19c q19d q19e q19f q19g q19h q19i q19j q19k q19l BY q05 /STATISTIC=CHISQ /CELLS= COUNT EXPECTED ROW COLUMN TOTAL .	
<b>Resources</b>	<b>Dimensions Requested</b>	2
	<b>Cells Available</b>	116508
	<b>Elapsed Time</b>	0:00:01.12

### Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
<b>Classification * Gender</b>	740	99.1%	7	.9%	747	100.0%
<b>Ethnic origin * Gender</b>	741	99.2%	6	.8%	747	100.0%
<b>Marital status * Gender</b>	739	98.9%	8	1.1%	747	100.0%
<b>Residence * Gender</b>	738	98.8%	9	1.2%	747	100.0%

<b>Working * Gender</b>	738	98.8%	9	1.2%	747	100.0%
<b>LIVING ARRAGEMENTS:WHERE * Gender</b>	739	98.9%	8	1.1%	747	100.0%
<b>LIVING WITH ROOMATE * Gender</b>	737	98.7%	10	1.3%	747	100.0%
<b>LIVING WITH ALONE * Gender</b>	737	98.7%	10	1.3%	747	100.0%
<b>LIVING WITH PARENTS * Gender</b>	737	98.7%	10	1.3%	747	100.0%
<b>LIVING WITH SPOUSE * Gender</b>	737	98.7%	10	1.3%	747	100.0%
<b>LIVING WITH CHILDREN * Gender</b>	737	98.7%	10	1.3%	747	100.0%
<b>LIVING WITH OTHER * Gender</b>	737	98.7%	10	1.3%	747	100.0%
<b>Grades * Gender</b>	740	99.1%	7	.9%	747	100.0%
<b>AVAILABILITY :DRUGS * Gender</b>	735	98.4%	12	1.6%	747	100.0%
<b>AVAILABILITY :ALCOHOL * Gender</b>	738	98.8%	9	1.2%	747	100.0%
<b>Student status * Gender</b>	736	98.5%	11	1.5%	747	100.0%
<b>Campus has a/d policies * Gender</b>	739	98.9%	8	1.1%	747	100.0%
<b>Campus a/d policies enforced * Gender</b>	736	98.5%	11	1.5%	747	100.0%
<b>Campus has a/d prev prog * Gender</b>	739	98.9%	8	1.1%	747	100.0%
<b>Campus concerned w/ a/d prev * Gender</b>	737	98.7%	10	1.3%	747	100.0%
<b>I am involved in a/d prev * Gender</b>	737	98.7%	10	1.3%	747	100.0%
<b>Permanent residence * Gender</b>	739	98.9%	8	1.1%	747	100.0%
<b>5+ drinks in last 2 wks * Gender</b>	739	98.9%	8	1.1%	747	100.0%
<b>First use: tobacco * Gender</b>	736	98.5%	11	1.5%	747	100.0%
<b>First use: alcohol * Gender</b>	736	98.5%	11	1.5%	747	100.0%
<b>First use: marijuana * Gender</b>	734	98.3%	13	1.7%	747	100.0%
<b>First use: cocaine * Gender</b>	734	98.3%	13	1.7%	747	100.0%
<b>First use: amphetamines * Gender</b>	735	98.4%	12	1.6%	747	100.0%
<b>First use: sedatives * Gender</b>	731	97.9%	16	2.1%	747	100.0%
<b>First use: hallucinogens * Gender</b>	734	98.3%	13	1.7%	747	100.0%
<b>First use: opiates * Gender</b>	731	97.9%	16	2.1%	747	100.0%
<b>First use: inhalants * Gender</b>	730	97.7%	17	2.3%	747	100.0%
<b>First use: designer * Gender</b>	735	98.4%	12	1.6%	747	100.0%
<b>First use: steroids * Gender</b>	733	98.1%	14	1.9%	747	100.0%
<b>First use: other * Gender</b>	735	98.4%	12	1.6%	747	100.0%
<b>Use last yr: tobacco * Gender</b>	737	98.7%	10	1.3%	747	100.0%
<b>Use last yr: alcohol * Gender</b>	736	98.5%	11	1.5%	747	100.0%
<b>Use last yr: marijuana * Gender</b>	736	98.5%	11	1.5%	747	100.0%
<b>Use last yr: cocaine * Gender</b>	735	98.4%	12	1.6%	747	100.0%
<b>Use last yr: amphetamines * Gender</b>	735	98.4%	12	1.6%	747	100.0%
<b>Use last yr: sedatives * Gender</b>	733	98.1%	14	1.9%	747	100.0%
<b>Use last yr: hallucinogens * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use last yr: opiates * Gender</b>	734	98.3%	13	1.7%	747	100.0%
<b>Use last yr: inhalants * Gender</b>	734	98.3%	13	1.7%	747	100.0%
<b>Use last yr: designer * Gender</b>	735	98.4%	12	1.6%	747	100.0%

Use last yr: steroids * Gender	736	98.5%	11	1.5%	747	100.0%
Use last yr: other * Gender	737	98.7%	10	1.3%	747	100.0%
PAST 30 DAYS USE:TOBACCO * Gender	730	97.7%	17	2.3%	747	100.0%
PAST 30 DAYS USE:ALCOHOL * Gender	730	97.7%	17	2.3%	747	100.0%
PAST 30 DAYS USE:MARIJUANA * Gender	730	97.7%	17	2.3%	747	100.0%
PAST 30 DAYS USE:COCAINE * Gender	727	97.3%	20	2.7%	747	100.0%
PAST 30 DAYS USE:AMPHETAMINES * Gender	730	97.7%	17	2.3%	747	100.0%
PAST 30 DAYS USE:SEDATIVES * Gender	730	97.7%	17	2.3%	747	100.0%
PAST 30 DAYS USE:HALLUCINOGENS * Gender	728	97.5%	19	2.5%	747	100.0%
PAST 30 DAYS USE:OPIATES * Gender	728	97.5%	19	2.5%	747	100.0%
PAST 30 DAYS USE:INHALANTS * Gender	729	97.6%	18	2.4%	747	100.0%
PAST 30 DAYS USE:DESIGNER * Gender	731	97.9%	16	2.1%	747	100.0%
PAST 30 DAYS USE:STEROIDS * Gender	728	97.5%	19	2.5%	747	100.0%
PAST 30 DAYS USE:OTHER * Gender	730	97.7%	17	2.3%	747	100.0%
AVERAGE USE:TOBACCO * Gender	733	98.1%	14	1.9%	747	100.0%
AVERAGE USE:ALCOHOL * Gender	734	98.3%	13	1.7%	747	100.0%
AVERAGE USE:MARIJUANA * Gender	735	98.4%	12	1.6%	747	100.0%
AVERAGE USE:COCAINE * Gender	733	98.1%	14	1.9%	747	100.0%
AVERAGE USE:AMPHETAMINES * Gender	729	97.6%	18	2.4%	747	100.0%
AVERAGE USE:SEDATIVES * Gender	728	97.5%	19	2.5%	747	100.0%
AVERAGE USE:HALLUCINOGENS * Gender	730	97.7%	17	2.3%	747	100.0%
AVERAGE USE:OPIATES * Gender	730	97.7%	17	2.3%	747	100.0%
AVERAGE USE:INHALANTS * Gender	729	97.6%	18	2.4%	747	100.0%
AVERAGE USE:DESIGNER * Gender	732	98.0%	15	2.0%	747	100.0%
AVERAGE USE:STEROIDS * Gender	732	98.0%	15	2.0%	747	100.0%
AVERAGE USE:OTHER * Gender	733	98.1%	14	1.9%	747	100.0%

## Classification \* Gender

Crosstab

		Gender			Total
		Male	Female	Unknown	
Freshman	Count	44	116	1	161
	Expected Count	51.3	109.0	.7	161.0
	% within Classification	27.3%	72.0%	.6%	100.0%
	% within Gender	18.6%	23.2%	33.3%	21.8%
	% of Total	5.9%	15.7%	.1%	21.8%
	Count	48	94	0	142
	Expected Count	45.3	96.1	.6	142.0

Classification	Sophomore	% within Classification	33.8%	66.2%	.0%	100.0%
		% within Gender	20.3%	18.8%	.0%	19.2%
		% of Total	6.5%	12.7%	.0%	19.2%
	Junior	Count	84	158	2	244
		Expected Count	77.8	165.2	1.0	244.0
		% within Classification	34.4%	64.8%	.8%	100.0%
		% within Gender	35.6%	31.5%	66.7%	33.0%
		% of Total	11.4%	21.4%	.3%	33.0%
	Senior	Count	60	130	0	190
		Expected Count	60.6	128.6	.8	190.0
		% within Classification	31.6%	68.4%	.0%	100.0%
		% within Gender	25.4%	25.9%	.0%	25.7%
		% of Total	8.1%	17.6%	.0%	25.7%
	Grad/professional	Count	0	1	0	1
		Expected Count	.3	.7	.0	1.0
		% within Classification	.0%	100.0%	.0%	100.0%
		% within Gender	.0%	.2%	.0%	.1%
		% of Total	.0%	.1%	.0%	.1%
	Other	Count	0	2	0	2
		Expected Count	.6	1.4	.0	2.0
		% within Classification	.0%	100.0%	.0%	100.0%
		% within Gender	.0%	.4%	.0%	.3%
		% of Total	.0%	.3%	.0%	.3%
	Total	Count	236	501	3	740
		Expected Count	236.0	501.0	3.0	740.0
		% within Classification	31.9%	67.7%	.4%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.9%	67.7%	.4%	100.0%

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	6.530(a)	10	.769
Likelihood Ratio	8.555	10	.575
Linear-by-Linear Association	.405	1	.525
N of Valid Cases	740		

a 10 cells (55.6%) have expected count less than 5. The minimum expected count is .00.

## Ethnic origin \* Gender

### Crosstab

		Gender			Total
		Male	Female	Unknown	
Ethnic origin	Amer Ind/AK native	Count	2	2	0 4
		Expected Count	1.3	2.7	.0 4.0
		% within Ethnic origin	50.0%	50.0%	.0% 100.0%
		% within Gender	.8%	.4%	.0% .5%
		% of Total	.3%	.3%	.0% .5%
	Hispanic	Count	4	6	0 10
		Expected Count	3.2	6.8	.0 10.0
		% within Ethnic origin	40.0%	60.0%	.0% 100.0%
		% within Gender	1.7%	1.2%	.0% 1.3%
		% of Total	.5%	.8%	.0% 1.3%
	Asian/Pac Isl	Count	3	5	1 9
		Expected Count	2.9	6.1	.0 9.0
		% within Ethnic origin	33.3%	55.6%	11.1% 100.0%
		% within Gender	1.3%	1.0%	33.3% 1.2%
		% of Total	.4%	.7%	.1% 1.2%
	White (non-Hisp)	Count	219	457	2 678
		Expected Count	215.9	459.3	2.7 678.0
		% within Ethnic origin	32.3%	67.4%	.3% 100.0%
		% within Gender	92.8%	91.0%	66.7% 91.5%
		% of Total	29.6%	61.7%	.3% 91.5%
	Black (non-Hisp)	Count	4	23	0 27
		Expected Count	8.6	18.3	.1 27.0
		% within Ethnic origin	14.8%	85.2%	.0% 100.0%
		% within Gender	1.7%	4.6%	.0% 3.6%
		% of Total	.5%	3.1%	.0% 3.6%
	Other	Count	4	9	0 13
		Expected Count	4.1	8.8	.1 13.0
		% within Ethnic origin	30.8%	69.2%	.0% 100.0%
		% within Gender	1.7%	1.8%	.0% 1.8%
		% of Total	.5%	1.2%	.0% 1.8%
Total		Count	236	502	3 741
		Expected Count	236.0	502.0	3.0 741.0
		% within Ethnic origin	31.8%	67.7%	.4% 100.0%
		% within Gender	100.0%	100.0%	100.0% 100.0%
		% of Total	31.8%	67.7%	.4% 100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)

<b>Pearson Chi-Square</b>	30.738(a)	10		.001
<b>Likelihood Ratio</b>	10.675	10		.383
<b>Linear-by-Linear Association</b>	1.647	1		.199
<b>N of Valid Cases</b>	741			
a 11 cells (61.1%) have expected count less than 5. The minimum expected count is .02.				

## Marital status \* Gender

Crosstab

			Gender			Total		
			Male	Female	Unknown			
Marital status	Single	Count	229	471	3	703		
		Expected Count	223.6	476.6	2.9	703.0		
		% within Marital status	32.6%	67.0%	.4%	100.0%		
		% within Gender	97.4%	94.0%	100.0%	95.1%		
		% of Total	31.0%	63.7%	.4%	95.1%		
	Married	Count	5	24	0	29		
		Expected Count	9.2	19.7	.1	29.0		
		% within Marital status	17.2%	82.8%	.0%	100.0%		
		% within Gender	2.1%	4.8%	.0%	3.9%		
		% of Total	.7%	3.2%	.0%	3.9%		
	Separated	Count	0	1	0	1		
		Expected Count	.3	.7	.0	1.0		
		% within Marital status	.0%	100.0%	.0%	100.0%		
		% within Gender	.0%	.2%	.0%	.1%		
		% of Total	.0%	.1%	.0%	.1%		
	Divorced	Count	1	5	0	6		
		Expected Count	1.9	4.1	.0	6.0		
		% within Marital status	16.7%	83.3%	.0%	100.0%		
		% within Gender	.4%	1.0%	.0%	.8%		
		% of Total	.1%	.7%	.0%	.8%		
Total			Count	235	501	3	739	
			Expected Count	235.0	501.0	3.0	739.0	
			% within Marital status	31.8%	67.8%	.4%	100.0%	
			% within Gender	100.0%	100.0%	100.0%	100.0%	
			% of Total	31.8%	67.8%	.4%	100.0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)

<b>Pearson Chi-Square</b>	4.360(a)	6		.628
<b>Likelihood Ratio</b>	5.206	6		.518
<b>Linear-by-Linear Association</b>	2.908	1		.088
<b>N of Valid Cases</b>	739			
a 8 cells (66.7%) have expected count less than 5. The minimum expected count is .00.				

## Residence \* Gender

Crosstab

		Count	Gender			Total
			Male	Female	Unknown	
Residence	On campus	Count	111	251	1	363
		Expected Count	116.1	245.9	1.0	363.0
		% within Residence	30.6%	69.1%	.3%	100.0%
		% within Gender	47.0%	50.2%	50.0%	49.2%
		% of Total	15.0%	34.0%	.1%	49.2%
	Off campus	Count	125	249	1	375
		Expected Count	119.9	254.1	1.0	375.0
		% within Residence	33.3%	66.4%	.3%	100.0%
		% within Gender	53.0%	49.8%	50.0%	50.8%
		% of Total	16.9%	33.7%	.1%	50.8%
Total	On campus	Count	236	500	2	738
		Expected Count	236.0	500.0	2.0	738.0
		% within Residence	32.0%	67.8%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	32.0%	67.8%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	.644(a)	2	.725
<b>Likelihood Ratio</b>	.644	2	.725
<b>Linear-by-Linear Association</b>	.634	1	.426
<b>N of Valid Cases</b>	738		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .98.

## Working \* Gender

Crosstab

		Gender			Total	
		Male	Female	Unknown		
Working	Full time	Count	11	26	0	37
		Expected Count	11.8	25.1	.1	37.0
		% within Working	29.7%	70.3%	.0%	100.0%
		% within Gender	4.7%	5.2%	.0%	5.0%
		% of Total	1.5%	3.5%	.0%	5.0%
	Part time	Count	105	271	1	377
		Expected Count	120.0	255.9	1.0	377.0
		% within Working	27.9%	71.9%	.3%	100.0%
		% within Gender	44.7%	54.1%	50.0%	51.1%
	No	% of Total	14.2%	36.7%	.1%	51.1%
		Count	119	204	1	324
		Expected Count	103.2	220.0	.9	324.0
		% within Working	36.7%	63.0%	.3%	100.0%
		% within Gender	50.6%	40.7%	50.0%	43.9%
		% of Total	16.1%	27.6%	.1%	43.9%
Total	Count	235	501	2	738	
	Expected Count	235.0	501.0	2.0	738.0	
	% within Working	31.8%	67.9%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	31.8%	67.9%	.3%	100.0%	

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	6.559(a)	4	.161
Likelihood Ratio	6.641	4	.156
Linear-by-Linear Association	4.857	1	.028
N of Valid Cases	738		

a 3 cells (33.3%) have expected count less than 5. The minimum expected count is .10.

## LIVING ARRAGEMENTS:WHERE \* Gender

## Crosstab

		Gender			Total
		Male	Female	Unknown	
HOUSE/APARTMENT	Count	128	262	0	
	Expected Count	124.5	264.4	1.1	3
	% within LIVING ARRAGEMENTS:WHERE	32.8%	67.2%	.0%	100

LIVING ARRAGEMENTS:WHERE	RESIDENCE HALL	% within Gender	54.2%	52.3%	.0%	52
		% of Total	17.3%	35.5%	.0%	52
		Count	97	234	1	
		Expected Count	106.0	225.1	.9	3
		% within LIVING ARRAGEMENTS:WHERE	29.2%	70.5%	.3%	100
	FRATERNITY/SORORITY	% within Gender	41.1%	46.7%	50.0%	42
		% of Total	13.1%	31.7%	.1%	42
		Count	10	4	1	
		Expected Count	4.8	10.2	.0	
		% within LIVING ARRAGEMENTS:WHERE	66.7%	26.7%	6.7%	100
	OTHER	% within Gender	4.2%	.8%	50.0%	2
		% of Total	1.4%	.5%	.1%	2
		Count	1	1	0	
		Expected Count	.6	1.4	.0	
		% within LIVING ARRAGEMENTS:WHERE	50.0%	50.0%	.0%	100
	Total	% within Gender	.4%	.2%	.0%	
		% of Total	.1%	.1%	.0%	
		Count	236	501	2	
		Expected Count	236.0	501.0	2.0	7
		% within LIVING ARRAGEMENTS:WHERE	31.9%	67.8%	.3%	100
	Total	% within Gender	100.0%	100.0%	100.0%	100
		% of Total	31.9%	67.8%	.3%	100

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	34.692(a)	6	.000
Likelihood Ratio	17.330	6	.008
Linear-by-Linear Association	.391	1	.532
N of Valid Cases	739		

a 7 cells (58.3%) have expected count less than 5. The minimum expected count is .01.

## LIVING WITH ROOMATE \* Gender

## Crosstab

	Count	Gender			Total
		Male	Female	Unknown	
		42	117	0	159

		Expected Count	50.5	108.1	.4	159.0
	NO	% within LIVING WITH ROOMATE	26.4%	73.6%	.0%	100.0%
		% within Gender	17.9%	23.4%	.0%	21.6%
		% of Total	5.7%	15.9%	.0%	21.6%
LIVING WITH ROOMATE	NO	Count	192	384	2	578
		Expected Count	183.5	392.9	1.6	578.0
		% within LIVING WITH ROOMATE	33.2%	66.4%	.3%	100.0%
	YES	% within Gender	82.1%	76.6%	100.0%	78.4%
		% of Total	26.1%	52.1%	.3%	78.4%
		Count	234	501	2	737
Total		Expected Count	234.0	501.0	2.0	737.0
		% within LIVING WITH ROOMATE	31.8%	68.0%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	68.0%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	3.305(a)	2	.192
Likelihood Ratio	3.791	2	.150
Linear-by-Linear Association	2.348	1	.125
N of Valid Cases	737		

## LIVING WITH ALONE \* Gender

## Crosstab

		NO	Gender			Total
			Male	Female	Unknown	
LIVING WITH ALONE	NO	Count	212	453	2	667
		Expected Count	211.8	453.4	1.8	667.0
		% within LIVING WITH ALONE	31.8%	67.9%	.3%	100.0%
		% within Gender	90.6%	90.4%	100.0%	90.5%
		% of Total	28.8%	61.5%	.3%	90.5%
	YES	Count	22	48	0	70
		Expected Count	22.2	47.6	.2	70.0
	YES	% within LIVING WITH ALONE	31.4%	68.6%	.0%	100.0%
		% within Gender	9.4%	9.6%	.0%	9.5%
		% of Total	3.0%	6.5%	.0%	9.5%
		Count	234	501	2	737

Total	Expected Count	234.0	501.0	2.0	737.0
	% within LIVING WITH ALONE	31.8%	68.0%	.3%	100.0%
	% within Gender	100.0%	100.0%	100.0%	100.0%
	% of Total	31.8%	68.0%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.216(a)	2	.897
Likelihood Ratio	.406	2	.816
Linear-by-Linear Association	.000	1	.992
N of Valid Cases	737		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .19.

**LIVING WITH PARENTS \* Gender**

## Crosstab

		Gender			Total
		Male	Female	Unknown	
LIVING WITH PARENTS	NO	Count	220	472	2 694
		Expected Count	220.3	471.8	1.9 694.0
		% within LIVING WITH PARENTS	31.7%	68.0%	.3% 100.0%
		% within Gender	94.0%	94.2%	100.0% 94.2%
		% of Total	29.9%	64.0%	.3% 94.2%
	YES	Count	14	29	0 43
		Expected Count	13.7	29.2	.1 43.0
		% within LIVING WITH PARENTS	32.6%	67.4%	.0% 100.0%
		% within Gender	6.0%	5.8%	.0% 5.8%
		% of Total	1.9%	3.9%	.0% 5.8%
Total		Count	234	501	2 737
		Expected Count	234.0	501.0	2.0 737.0
		% within LIVING WITH PARENTS	31.8%	68.0%	.3% 100.0%
		% within Gender	100.0%	100.0%	100.0% 100.0%
		% of Total	31.8%	68.0%	.3% 100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.135(a)	2	.935
Likelihood Ratio	.252	2	.882
Linear-by-Linear Association	.024	1	.877

N of Valid Cases	737	
a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .12.		

## LIVING WITH SPOUSE \* Gender

Crosstab

		Gender			Total
		Male	Female	Unknown	
LIVING WITH SPOUSE	NO	Count	228	477	2 707
		Expected Count	224.5	480.6	1.9 707.0
		% within LIVING WITH SPOUSE	32.2%	67.5%	.3% 100.0%
		% within Gender	97.4%	95.2%	100.0% 95.9%
		% of Total	30.9%	64.7%	.3% 95.9%
	YES	Count	6	24	0 30
		Expected Count	9.5	20.4	.1 30.0
		% within LIVING WITH SPOUSE	20.0%	80.0%	.0% 100.0%
		% within Gender	2.6%	4.8%	.0% 4.1%
		% of Total	.8%	3.3%	.0% 4.1%
Total		Count	234	501	2 737
		Expected Count	234.0	501.0	2.0 737.0
		% within LIVING WITH SPOUSE	31.8%	68.0%	.3% 100.0%
		% within Gender	100.0%	100.0%	100.0% 100.0%
		% of Total	31.8%	68.0%	.3% 100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	2.110(a)	2	.348
Likelihood Ratio	2.356	2	.308
Linear-by-Linear Association	1.861	1	.173
N of Valid Cases	737		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .08.

## LIVING WITH CHILDREN \* Gender

Crosstab

		Gender			Total
		Male	Female	Unknown	
	Count	231	477	2	710

LIVING WITH CHILDREN	NO	Expected Count	225.4	482.6	1.9	710.0
		% within LIVING WITH CHILDREN	32.5%	67.2%	.3%	100.0%
		% within Gender	98.7%	95.2%	100.0%	96.3%
		% of Total	31.3%	64.7%	.3%	96.3%
	YES	Count	3	24	0	27
		Expected Count	8.6	18.4	.1	27.0
		% within LIVING WITH CHILDREN	11.1%	88.9%	.0%	100.0%
		% within Gender	1.3%	4.8%	.0%	3.7%
		% of Total	.4%	3.3%	.0%	3.7%
Total	Count	234	501	2	737	
	Expected Count	234.0	501.0	2.0	737.0	
	% within LIVING WITH CHILDREN	31.8%	68.0%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	31.8%	68.0%	.3%	100.0%	

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	5.639(a)	2	.060
Likelihood Ratio	6.779	2	.034
Linear-by-Linear Association	5.251	1	.022
N of Valid Cases	737		

## LIVING WITH OTHER \* Gender

#### Crosstab

			Gender			Total	
			Male	Female	Unknown		
LIVING WITH OTHER	NO	Count	233	490	2	725	
		Expected Count	230.2	492.8	2.0	725.0	
		% within LIVING WITH OTHER	32.1%	67.6%	.3%	100.0%	
		% within Gender	99.6%	97.8%	100.0%	98.4%	
		% of Total	31.6%	66.5%	.3%	98.4%	
	YES	Count	1	11	0	12	
		Expected Count	3.8	8.2	.0	12.0	
		% within LIVING WITH OTHER	8.3%	91.7%	.0%	100.0%	
		% within Gender	.4%	2.2%	.0%	1.6%	
		% of Total	.1%	1.5%	.0%	1.6%	
			Count	234	501	2	
						737	

<b>Total</b>	<b>Expected Count</b>	234.0	501.0	2.0	737.0
	<b>% within LIVING WITH OTHER</b>	31.8%	68.0%	.3%	100.0%
	<b>% within Gender</b>	100.0%	100.0%	100.0%	100.0%
	<b>% of Total</b>	31.8%	68.0%	.3%	100.0%

**Chi-Square Tests**

	<b>Value</b>	<b>df</b>	<b>Asymp. Sig. (2-sided)</b>
<b>Pearson Chi-Square</b>	3.147(a)	2	.207
<b>Likelihood Ratio</b>	3.953	2	.139
<b>Linear-by-Linear Association</b>	2.951	1	.086
<b>N of Valid Cases</b>	737		

a 3 cells (50.0%) have expected count less than 5. The minimum expected count is .03.

**Grades \* Gender****Crosstab**

			<b>Gender</b>			<b>Total</b>
			<b>Male</b>	<b>Female</b>	<b>Unknown</b>	
<b>F</b>	<b>Count</b>	0	1	0	0	1
	<b>Expected Count</b>	.3	.7	.0	.0	1.0
	<b>% within Grades</b>	.0%	100.0%	.0%	.0%	100.0%
	<b>% within Gender</b>	.0%	.2%	.0%	.0%	.1%
	<b>% of Total</b>	.0%	.1%	.0%	.0%	.1%
<b>D-</b>	<b>Count</b>	0	1	0	0	1
	<b>Expected Count</b>	.3	.7	.0	.0	1.0
	<b>% within Grades</b>	.0%	100.0%	.0%	.0%	100.0%
	<b>% within Gender</b>	.0%	.2%	.0%	.0%	.1%
	<b>% of Total</b>	.0%	.1%	.0%	.0%	.1%
<b>D</b>	<b>Count</b>	2	3	0	0	5
	<b>Expected Count</b>	1.6	3.4	.0	.0	5.0
	<b>% within Grades</b>	40.0%	60.0%	.0%	.0%	100.0%
	<b>% within Gender</b>	.8%	.6%	.0%	.0%	.7%
	<b>% of Total</b>	.3%	.4%	.0%	.0%	.7%
<b>D+</b>	<b>Count</b>	2	0	0	0	2
	<b>Expected Count</b>	.6	1.4	.0	.0	2.0
	<b>% within Grades</b>	100.0%	.0%	.0%	.0%	100.0%
	<b>% within Gender</b>	.8%	.0%	.0%	.0%	.3%
	<b>% of Total</b>	.3%	.0%	.0%	.0%	.3%
	<b>Count</b>	7	4	0	0	11

Grades	C-	Expected Count	3.5	7.5	.0	11.0
		% within Grades	63.6%	36.4%	.0%	100.0%
		% within Gender	3.0%	.8%	.0%	1.5%
		% of Total	.9%	.5%	.0%	1.5%
	C	Count	20	27	0	47
		Expected Count	15.0	31.9	.1	47.0
		% within Grades	42.6%	57.4%	.0%	100.0%
		% within Gender	8.5%	5.4%	.0%	6.4%
		% of Total	2.7%	3.6%	.0%	6.4%
	C+	Count	34	41	1	76
		Expected Count	24.2	51.6	.2	76.0
		% within Grades	44.7%	53.9%	1.3%	100.0%
		% within Gender	14.4%	8.2%	50.0%	10.3%
		% of Total	4.6%	5.5%	.1%	10.3%
	B-	Count	22	55	0	77
		Expected Count	24.6	52.2	.2	77.0
		% within Grades	28.6%	71.4%	.0%	100.0%
		% within Gender	9.3%	11.0%	.0%	10.4%
		% of Total	3.0%	7.4%	.0%	10.4%
	B	Count	56	134	0	190
		Expected Count	60.6	128.9	.5	190.0
		% within Grades	29.5%	70.5%	.0%	100.0%
		% within Gender	23.7%	26.7%	.0%	25.7%
		% of Total	7.6%	18.1%	.0%	25.7%
	B+	Count	42	72	0	114
		Expected Count	36.4	77.3	.3	114.0
		% within Grades	36.8%	63.2%	.0%	100.0%
		% within Gender	17.8%	14.3%	.0%	15.4%
		% of Total	5.7%	9.7%	.0%	15.4%
	A-	Count	20	56	1	77
		Expected Count	24.6	52.2	.2	77.0
		% within Grades	26.0%	72.7%	1.3%	100.0%
		% within Gender	8.5%	11.2%	50.0%	10.4%
		% of Total	2.7%	7.6%	.1%	10.4%
	A	Count	22	91	0	113
		Expected Count	36.0	76.7	.3	113.0
		% within Grades	19.5%	80.5%	.0%	100.0%
		% within Gender	9.3%	18.1%	.0%	15.3%
		% of Total	3.0%	12.3%	.0%	15.3%
		Count	9	17	0	26

		<b>Expected Count</b>	8.3	17.6	.1	26.0
	<b>A+</b>	<b>% within Grades</b>	34.6%	65.4%	.0%	100.0%
		<b>% within Gender</b>	3.8%	3.4%	.0%	3.5%
		<b>% of Total</b>	1.2%	2.3%	.0%	3.5%
		<b>Count</b>	236	502	2	740
		<b>Expected Count</b>	236.0	502.0	2.0	740.0
	<b>Total</b>	<b>% within Grades</b>	31.9%	67.8%	.3%	100.0%
		<b>% within Gender</b>	100.0%	100.0%	100.0%	100.0%
		<b>% of Total</b>	31.9%	67.8%	.3%	100.0%

**Chi-Square Tests**

	<b>Value</b>	<b>df</b>	<b>Asymp. Sig. (2-sided)</b>
<b>Pearson Chi-Square</b>	38.189(a)	24	.033
<b>Likelihood Ratio</b>	37.623	24	.038
<b>Linear-by-Linear Association</b>	12.448	1	.000
<b>N of Valid Cases</b>	740		

a 22 cells (56.4%) have expected count less than 5. The minimum expected count is .00.

**AVAILABILITY :DRUGS \* Gender****Crosstab**

			<b>Gender</b>			<b>Total</b>
			<b>Male</b>	<b>Female</b>	<b>Unknown</b>	
<b>AVAILABILITY :DRUGS</b>	<b>Available</b>	<b>Count</b>	60	79	0	139
		<b>Expected Count</b>	44.6	94.0	.4	139.0
		<b>% within AVAILABILITY :DRUGS</b>	43.2%	56.8%	.0%	100.0%
		<b>% within Gender</b>	25.4%	15.9%	.0%	18.9%
		<b>% of Total</b>	8.2%	10.7%	.0%	18.9%
	<b>Not available</b>	<b>Count</b>	176	418	2	596
		<b>Expected Count</b>	191.4	403.0	1.6	596.0
		<b>% within AVAILABILITY :DRUGS</b>	29.5%	70.1%	.3%	100.0%
		<b>% within Gender</b>	74.6%	84.1%	100.0%	81.1%
		<b>% of Total</b>	23.9%	56.9%	.3%	81.1%
<b>Total</b>		<b>Count</b>	236	497	2	735
		<b>Expected Count</b>	236.0	497.0	2.0	735.0
		<b>% within AVAILABILITY :DRUGS</b>	32.1%	67.6%	.3%	100.0%
		<b>% within Gender</b>	100.0%	100.0%	100.0%	100.0%
		<b>% of Total</b>	32.1%	67.6%	.3%	100.0%

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	9.941(a)	2	.007
<b>Likelihood Ratio</b>	9.959	2	.007
<b>Linear-by-Linear Association</b>	9.876	1	.002
<b>N of Valid Cases</b>	735		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .38.

## AVAILABILITY :ALCOHOL \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
AVAILABILITY :ALCOHOL	Available	Count	202	412	2	616
		Expected Count	197.0	417.3	1.7	616.0
		% within AVAILABILITY :ALCOHOL	32.8%	66.9%	.3%	100.0%
		% within Gender	85.6%	82.4%	100.0%	83.5%
		% of Total	27.4%	55.8%	.3%	83.5%
	Not available	Count	34	88	0	122
		Expected Count	39.0	82.7	.3	122.0
		% within AVAILABILITY :ALCOHOL	27.9%	72.1%	.0%	100.0%
		% within Gender	14.4%	17.6%	.0%	16.5%
		% of Total	4.6%	11.9%	.0%	16.5%
Total		Count	236	500	2	738
		Expected Count	236.0	500.0	2.0	738.0
		% within AVAILABILITY :ALCOHOL	32.0%	67.8%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	32.0%	67.8%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	1.582(a)	2	.453
<b>Likelihood Ratio</b>	1.930	2	.381
<b>Linear-by-Linear Association</b>	.969	1	.325
<b>N of Valid Cases</b>	738		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .33.

## Student status \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Student status	FULL TIME	Count	223	480	2	705
		Expected Count	223.2	479.9	1.9	705.0
		% within Student status	31.6%	68.1%	.3%	100.0%
		% within Gender	95.7%	95.8%	100.0%	95.8%
		% of Total	30.3%	65.2%	.3%	95.8%
	PART TIME	Count	10	21	0	31
		Expected Count	9.8	21.1	.1	31.0
		% within Student status	32.3%	67.7%	.0%	100.0%
		% within Gender	4.3%	4.2%	.0%	4.2%
		% of Total	1.4%	2.9%	.0%	4.2%
Total	Count	233	501	2	736	
	Expected Count	233.0	501.0	2.0	736.0	
	% within Student status	31.7%	68.1%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	31.7%	68.1%	.3%	100.0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.092(a)	2	.955
Likelihood Ratio	.176	2	.916
Linear-by-Linear Association	.011	1	.916
N of Valid Cases	736		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .08.

## Campus has a/d policies \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Yes	Yes	Count	216	436	2	654
		Expected Count	208.0	444.3	1.8	654.0
		% within Campus has a/d policies	33.0%	66.7%	.3%	100.0%
		% within Gender	91.9%	86.9%	100.0%	88.5%
		% of Total	29.2%	59.0%	.3%	88.5%
		Count	1	3	0	4
		Expected Count	1.3	2.7	.0	4.0
		% within Campus has a/d policies	25.0%	75.0%	0.0%	100.0%
		% within Gender	25.0%	75.0%	0.0%	100.0%
		% of Total	25.0%	75.0%	0.0%	100.0%

Campus has a/d policies	No	% within Campus has a/d policies	25.0%	75.0%	.0%	100.0%
		% within Gender	.4%	.6%	.0%	.5%
		% of Total	.1%	.4%	.0%	.5%
	Don't know	Count	18	63	0	81
		Expected Count	25.8	55.0	.2	81.0
		% within Campus has a/d policies	22.2%	77.8%	.0%	100.0%
		% within Gender	7.7%	12.5%	.0%	11.0%
		% of Total	2.4%	8.5%	.0%	11.0%
	Total	Count	235	502	2	739
		Expected Count	235.0	502.0	2.0	739.0
		% within Campus has a/d policies	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	4.304(a)	4	.366
Likelihood Ratio	4.759	4	.313
Linear-by-Linear Association	3.652	1	.056
N of Valid Cases	739		

a 5 cells (55.6%) have expected count less than 5. The minimum expected count is .01.

## Campus a/d policies enforced \* Gender

#### Crosstab

Campus a/d policies enforced		Yes	Gender			Total
			Male	Female	Unknown	
			Count	294	2	
Campus a/d policies enforced	Yes	Expected Count	148.2	314.6	1.3	464.0
		% within Campus a/d policies enforced	36.2%	63.4%	.4%	100.0%
		% within Gender	71.5%	58.9%	100.0%	63.0%
		% of Total	22.8%	39.9%	.3%	63.0%
		Count	16	43	0	59
	No	Expected Count	18.8	40.0	.2	59.0
		% within Campus a/d policies enforced	27.1%	72.9%	.0%	100.0%
		% within Gender	6.8%	8.6%	.0%	8.0%
		% of Total	2.2%	5.8%	.0%	8.0%
		Count	51	162	0	213

		Expected Count	68.0	144.4	.6	213.0
	Don't know	% within Campus a/d policies enforced	23.9%	76.1%	.0%	100.0%
		% within Gender	21.7%	32.5%	.0%	28.9%
		% of Total	6.9%	22.0%	.0%	28.9%
		Count	235	499	2	736
		Expected Count	235.0	499.0	2.0	736.0
Total		% within Campus a/d policies enforced	31.9%	67.8%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.9%	67.8%	.3%	100.0%

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	12.227(a)	4	.016
Likelihood Ratio	13.162	4	.011
Linear-by-Linear Association	9.616	1	.002
N of Valid Cases	736		

a 3 cells (33.3%) have expected count less than 5. The minimum expected count is .16.

## Campus has a/d prev prog \* Gender

### Crosstab

			Gender			Total
			Male	Female	Unknown	
Campus has a/d prev prog	Yes	Count	137	248	1	386
		Expected Count	122.7	262.2	1.0	386.0
		% within Campus has a/d prev prog	35.5%	64.2%	.3%	100.0%
		% within Gender	58.3%	49.4%	50.0%	52.2%
		% of Total	18.5%	33.6%	.1%	52.2%
	No	Count	4	9	0	13
		Expected Count	4.1	8.8	.0	13.0
		% within Campus has a/d prev prog	30.8%	69.2%	.0%	100.0%
		% within Gender	1.7%	1.8%	.0%	1.8%
		% of Total	.5%	1.2%	.0%	1.8%
	Don't know	Count	94	245	1	340
		Expected Count	108.1	231.0	.9	340.0
		% within Campus has a/d prev prog	27.6%	72.1%	.3%	100.0%

		% within Gender	40.0%	48.8%	50.0%	46.0%
		% of Total	12.7%	33.2%	.1%	46.0%
Total	Count	235	502	2	739	
	Expected Count	235.0	502.0	2.0	739.0	
	% within Campus has a/d prev prog	31.8%	67.9%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	31.8%	67.9%	.3%	100.0%	

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	5.174(a)	4	.270
Likelihood Ratio	5.232	4	.264
Linear-by-Linear Association	5.068	1	.024
N of Valid Cases	739		

a 4 cells (44.4%) have expected count less than 5. The minimum expected count is .04.

## Campus concerned w/ a/d prev \* Gender

### Crosstab

			Gender			Total
			Male	Female	Unknown	
Campus concerned w/ a/d prev	Yes	Count	174	353	2	529
		Expected Count	168.7	358.9	1.4	529.0
		% within Campus concerned w/ a/d prev	32.9%	66.7%	.4%	100.0%
		% within Gender	74.0%	70.6%	100.0%	71.8%
		% of Total	23.6%	47.9%	.3%	71.8%
	No	Count	35	50	0	85
		Expected Count	27.1	57.7	.2	85.0
		% within Campus concerned w/ a/d prev	41.2%	58.8%	.0%	100.0%
		% within Gender	14.9%	10.0%	.0%	11.5%
		% of Total	4.7%	6.8%	.0%	11.5%
	Don't know	Count	26	97	0	123
		Expected Count	39.2	83.4	.3	123.0
		% within Campus concerned w/ a/d prev	21.1%	78.9%	.0%	100.0%
		% within Gender	11.1%	19.4%	.0%	16.7%
		% of Total	3.5%	13.2%	.0%	16.7%
		Count	235	500	2	737

Total	Expected Count	235.0	500.0	2.0	737.0
	% within Campus concerned w/ a/d prev	31.9%	67.8%	.3%	100.0%
	% within Gender	100.0%	100.0%	100.0%	100.0%
	% of Total	31.9%	67.8%	.3%	100.0%

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	11.028(a)	4	.026
Likelihood Ratio	11.917	4	.018
Linear-by-Linear Association	3.275	1	.070
N of Valid Cases	737		

a 3 cells (33.3%) have expected count less than 5. The minimum expected count is .23.

## I am involved in a/d prev \* Gender

### Crosstab

		Gender			Total
		Male	Female	Unknown	
I am involved in a/d prev	Yes	Count	14	31	0 45
		Expected Count	14.3	30.6	.1 45.0
		% within I am involved in a/d prev	31.1%	68.9%	.0% 100.0%
		% within Gender	6.0%	6.2%	.0% 6.1%
		% of Total	1.9%	4.2%	.0% 6.1%
	No	Count	220	470	2 692
		Expected Count	219.7	470.4	1.9 692.0
		% within I am involved in a/d prev	31.8%	67.9%	.3% 100.0%
		% within Gender	94.0%	93.8%	100.0% 93.9%
		% of Total	29.9%	63.8%	.3% 93.9%
Total		Count	234	501	2 737
		Expected Count	234.0	501.0	2.0 737.0
		% within I am involved in a/d prev	31.8%	68.0%	.3% 100.0%
		% within Gender	100.0%	100.0%	100.0% 100.0%
		% of Total	31.8%	68.0%	.3% 100.0%

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.142(a)	2	.931
Likelihood Ratio	.264	2	.876
Linear-by-Linear Association	.003	1	.957

N of Valid Cases	737	
a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .12.		

## Permanent residence \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Permanent residence	In state	Count	232	483	2	717
		Expected Count	229.0	486.1	1.9	717.0
		% within Permanent residence	32.4%	67.4%	.3%	100.0%
		% within Gender	98.3%	96.4%	100.0%	97.0%
		% of Total	31.4%	65.4%	.3%	97.0%
	USA, other state	Count	4	15	0	19
		Expected Count	6.1	12.9	.1	19.0
		% within Permanent residence	21.1%	78.9%	.0%	100.0%
		% within Gender	1.7%	3.0%	.0%	2.6%
		% of Total	.5%	2.0%	.0%	2.6%
	Another country	Count	0	3	0	3
		Expected Count	1.0	2.0	.0	3.0
		% within Permanent residence	.0%	100.0%	.0%	100.0%
		% within Gender	.0%	.6%	.0%	.4%
		% of Total	.0%	.4%	.0%	.4%
Total		Count	236	501	2	739
		Expected Count	236.0	501.0	2.0	739.0
		% within Permanent residence	31.9%	67.8%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.9%	67.8%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	2.591(a)	4	.628
Likelihood Ratio	3.629	4	.459
Linear-by-Linear Association	2.290	1	.130
N of Valid Cases	739		

a 5 cells (55.6%) have expected count less than 5. The minimum expected count is .01.

## 5+ drinks in last 2 wks \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
5+ drinks in last 2 wks	None	Count	62	203	1	266
		Expected Count	84.6	180.7	.7	266.0
		% within 5+ drinks in last 2 wks	23.3%	76.3%	.4%	100.0%
		% within Gender	26.4%	40.4%	50.0%	36.0%
		% of Total	8.4%	27.5%	.1%	36.0%
	Once	Count	32	86	0	118
		Expected Count	37.5	80.2	.3	118.0
		% within 5+ drinks in last 2 wks	27.1%	72.9%	.0%	100.0%
		% within Gender	13.6%	17.1%	.0%	16.0%
		% of Total	4.3%	11.6%	.0%	16.0%
	Twice	Count	41	74	0	115
		Expected Count	36.6	78.1	.3	115.0
		% within 5+ drinks in last 2 wks	35.7%	64.3%	.0%	100.0%
		% within Gender	17.4%	14.7%	.0%	15.6%
		% of Total	5.5%	10.0%	.0%	15.6%
	3-5 times	Count	57	104	1	162
		Expected Count	51.5	110.0	.4	162.0
		% within 5+ drinks in last 2 wks	35.2%	64.2%	.6%	100.0%
		% within Gender	24.3%	20.7%	50.0%	21.9%
		% of Total	7.7%	14.1%	.1%	21.9%
	6-9 times	Count	37	29	0	66
		Expected Count	21.0	44.8	.2	66.0
		% within 5+ drinks in last 2 wks	56.1%	43.9%	.0%	100.0%
		% within Gender	15.7%	5.8%	.0%	8.9%
		% of Total	5.0%	3.9%	.0%	8.9%
	10+ times	Count	6	6	0	12
		Expected Count	3.8	8.2	.0	12.0
		% within 5+ drinks in last 2 wks	50.0%	50.0%	.0%	100.0%
		% within Gender	2.6%	1.2%	.0%	1.6%
		% of Total	.8%	.8%	.0%	1.6%
Total		Count	235	502	2	739
		Expected Count	235.0	502.0	2.0	739.0
		% within 5+ drinks in last 2 wks	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	32.990(a)	10	.000
<b>Likelihood Ratio</b>	32.539	10	.000
<b>Linear-by-Linear Association</b>	25.761	1	.000
<b>N of Valid Cases</b>	739		

a 7 cells (38.9%) have expected count less than 5. The minimum expected count is .03.

## First use: tobacco \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
First use: tobacco	Did not use	Count	71	209	0	280
		Expected Count	88.3	191.0	.8	280.0
		% within First use: tobacco	25.4%	74.6%	.0%	100.0%
		% within Gender	30.6%	41.6%	.0%	38.0%
		% of Total	9.6%	28.4%	.0%	38.0%
	Under 10	Count	2	5	0	7
		Expected Count	2.2	4.8	.0	7.0
		% within First use: tobacco	28.6%	71.4%	.0%	100.0%
		% within Gender	.9%	1.0%	.0%	1.0%
		% of Total	.3%	.7%	.0%	1.0%
	10-11	Count	7	10	0	17
		Expected Count	5.4	11.6	.0	17.0
		% within First use: tobacco	41.2%	58.8%	.0%	100.0%
		% within Gender	3.0%	2.0%	.0%	2.3%
		% of Total	1.0%	1.4%	.0%	2.3%
	12-13	Count	24	64	0	88
		Expected Count	27.7	60.0	.2	88.0
		% within First use: tobacco	27.3%	72.7%	.0%	100.0%
		% within Gender	10.3%	12.7%	.0%	12.0%
		% of Total	3.3%	8.7%	.0%	12.0%
	14-15	Count	32	76	1	109
		Expected Count	34.4	74.3	.3	109.0
		% within First use: tobacco	29.4%	69.7%	.9%	100.0%
		% within Gender	13.8%	15.1%	50.0%	14.8%
		% of Total	4.3%	10.3%	.1%	14.8%
		Count	49	72	1	122
		Expected Count	38.5	83.2	.3	122.0

		% within First use: tobacco	40.2%	59.0%	.8%	100.0%
		% within Gender	21.1%	14.3%	50.0%	16.6%
		% of Total	6.7%	9.8%	.1%	16.6%
16-17	18-20	Count	44	63	0	107
		Expected Count	33.7	73.0	.3	107.0
		% within First use: tobacco	41.1%	58.9%	.0%	100.0%
		% within Gender	19.0%	12.5%	.0%	14.5%
		% of Total	6.0%	8.6%	.0%	14.5%
21-25	Total	Count	3	3	0	6
		Expected Count	1.9	4.1	.0	6.0
		% within First use: tobacco	50.0%	50.0%	.0%	100.0%
		% within Gender	1.3%	.6%	.0%	.8%
		% of Total	.4%	.4%	.0%	.8%
Total	Total	Count	232	502	2	736
		Expected Count	232.0	502.0	2.0	736.0
		% within First use: tobacco	31.5%	68.2%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.5%	68.2%	.3%	100.0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	21.023(a)	14	.101
Likelihood Ratio	21.016	14	.101
Linear-by-Linear Association	11.077	1	.001
N of Valid Cases	736		

a 12 cells (50.0%) have expected count less than 5. The minimum expected count is .02.

## First use: alcohol \* Gender

#### Crosstab

			Gender			Total
			Male	Female	Unknown	
Did not use	Did not use	Count	11	32	0	43
		Expected Count	13.6	29.3	.1	43.0
		% within First use: alcohol	25.6%	74.4%	.0%	100.0%
		% within Gender	4.7%	6.4%	.0%	5.8%
		% of Total	1.5%	4.3%	.0%	5.8%
		Count	3	5	0	8
		Expected Count	2.5	5.5	.0	8.0

First use: alcohol	Under 10	% within First use: alcohol	37.5%	62.5%	.0%	100.0%	
		% within Gender	1.3%	1.0%	.0%	1.1%	
		% of Total	.4%	.7%	.0%	1.1%	
	10-11	Count	4	6	0	10	
		Expected Count	3.2	6.8	.0	10.0	
		% within First use: alcohol	40.0%	60.0%	.0%	100.0%	
		% within Gender	1.7%	1.2%	.0%	1.4%	
		% of Total	.5%	.8%	.0%	1.4%	
	12-13	Count	20	35	0	55	
		Expected Count	17.3	37.5	.1	55.0	
		% within First use: alcohol	36.4%	63.6%	.0%	100.0%	
		% within Gender	8.6%	7.0%	.0%	7.5%	
		% of Total	2.7%	4.8%	.0%	7.5%	
	14-15	Count	45	105	1	151	
		Expected Count	47.6	103.0	.4	151.0	
		% within First use: alcohol	29.8%	69.5%	.7%	100.0%	
		% within Gender	19.4%	20.9%	50.0%	20.5%	
		% of Total	6.1%	14.3%	.1%	20.5%	
	16-17	Count	78	163	0	241	
		Expected Count	76.0	164.4	.7	241.0	
		% within First use: alcohol	32.4%	67.6%	.0%	100.0%	
		% within Gender	33.6%	32.5%	.0%	32.7%	
		% of Total	10.6%	22.1%	.0%	32.7%	
	18-20	Count	61	132	1	194	
		Expected Count	61.2	132.3	.5	194.0	
		% within First use: alcohol	31.4%	68.0%	.5%	100.0%	
		% within Gender	26.3%	26.3%	50.0%	26.4%	
		% of Total	8.3%	17.9%	.1%	26.4%	
	21-25	Count	10	22	0	32	
		Expected Count	10.1	21.8	.1	32.0	
		% within First use: alcohol	31.3%	68.8%	.0%	100.0%	
		% within Gender	4.3%	4.4%	.0%	4.3%	
		% of Total	1.4%	3.0%	.0%	4.3%	
	26+	Count	0	2	0	2	
		Expected Count	.6	1.4	.0	2.0	
		% within First use: alcohol	.0%	100.0%	.0%	100.0%	
		% within Gender	.0%	.4%	.0%	.3%	
		% of Total	.0%	.3%	.0%	.3%	
		Count	232	502	2	736	
		Expected Count	232.0	502.0	2.0	736.0	

Total	% within First use: alcohol	31.5%	68.2%	.3%	100.0%
	% within Gender	100.0%	100.0%	100.0%	100.0%
	% of Total	31.5%	68.2%	.3%	100.0%

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	5.269(a)	16	.994
<b>Likelihood Ratio</b>	6.584	16	.980
<b>Linear-by-Linear Association</b>	.016	1	.898
<b>N of Valid Cases</b>	736		

a 13 cells (48.1%) have expected count less than 5. The minimum expected count is .01.

# First use: marijuana \* Gender

**Crosstab**

			Gender			Total
			Male	Female	Unknown	
First use: marijuana	Did not use	Count	97	261	0	358
		Expected Count	112.7	244.4	1.0	358.0
		% within First use: marijuana	27.1%	72.9%	.0%	100.0%
		% within Gender	42.0%	52.1%	.0%	48.8%
		% of Total	13.2%	35.6%	.0%	48.8%
	Under 10	Count	0	1	0	1
		Expected Count	.3	.7	.0	1.0
		% within First use: marijuana	.0%	100.0%	.0%	100.0%
		% within Gender	.0%	.2%	.0%	.1%
		% of Total	.0%	.1%	.0%	.1%
	10-11	Count	0	2	0	2
		Expected Count	.6	1.4	.0	2.0
		% within First use: marijuana	.0%	100.0%	.0%	100.0%
		% within Gender	.0%	.4%	.0%	.3%
		% of Total	.0%	.3%	.0%	.3%
	12-13	Count	10	18	0	28
		Expected Count	8.8	19.1	.1	28.0
		% within First use: marijuana	35.7%	64.3%	.0%	100.0%
		% within Gender	4.3%	3.6%	.0%	3.8%
		% of Total	1.4%	2.5%	.0%	3.8%
		Count	26	53	0	79
		Expected Count	24.9	53.9	.2	79.0

		% within First use: marijuana	32.9%	67.1%	.0%	100.0%	
		% within Gender	11.3%	10.6%	.0%	10.8%	
		% of Total	3.5%	7.2%	.0%	10.8%	
14-15	16-17	Count	49	88	0	137	
		Expected Count	43.1	93.5	.4	137.0	
		% within First use: marijuana	35.8%	64.2%	.0%	100.0%	
		% within Gender	21.2%	17.6%	.0%	18.7%	
		% of Total	6.7%	12.0%	.0%	18.7%	
18-20	21-25	Count	46	73	2	121	
		Expected Count	38.1	82.6	.3	121.0	
		% within First use: marijuana	38.0%	60.3%	1.7%	100.0%	
		% within Gender	19.9%	14.6%	100.0%	16.5%	
		% of Total	6.3%	9.9%	.3%	16.5%	
Total		Count	3	5	0	8	
		Expected Count	2.5	5.5	.0	8.0	
		% within First use: marijuana	37.5%	62.5%	.0%	100.0%	
		% within Gender	1.3%	1.0%	.0%	1.1%	
		% of Total	.4%	.7%	.0%	1.1%	
		Count	231	501	2	734	
		Expected Count	231.0	501.0	2.0	734.0	
		% within First use: marijuana	31.5%	68.3%	.3%	100.0%	
		% within Gender	100.0%	100.0%	100.0%	100.0%	
		% of Total	31.5%	68.3%	.3%	100.0%	

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	19.144(a)	14	.160
Likelihood Ratio	17.098	14	.251
Linear-by-Linear Association	5.891	1	.015
N of Valid Cases	734		

a 13 cells (54.2%) have expected count less than 5. The minimum expected count is .00.

## First use: cocaine \* Gender

#### Crosstab

		Gender			Total
		Male	Female	Unknown	
		Count	204	459	2
		Expected Count	210.2	453.0	1.8

First use: cocaine	Did not use	% within First use: cocaine	30.7%	69.0%	.3%	100.0%
		% within Gender	87.9%	91.8%	100.0%	90.6%
		% of Total	27.8%	62.5%	.3%	90.6%
	12-13	Count	1	1	0	2
		Expected Count	.6	1.4	.0	2.0
		% within First use: cocaine	50.0%	50.0%	.0%	100.0%
		% within Gender	.4%	.2%	.0%	.3%
		% of Total	.1%	.1%	.0%	.3%
	14-15	Count	1	5	0	6
		Expected Count	1.9	4.1	.0	6.0
		% within First use: cocaine	16.7%	83.3%	.0%	100.0%
		% within Gender	.4%	1.0%	.0%	.8%
		% of Total	.1%	.7%	.0%	.8%
	16-17	Count	5	11	0	16
		Expected Count	5.1	10.9	.0	16.0
		% within First use: cocaine	31.3%	68.8%	.0%	100.0%
		% within Gender	2.2%	2.2%	.0%	2.2%
		% of Total	.7%	1.5%	.0%	2.2%
	18-20	Count	14	20	0	34
		Expected Count	10.7	23.2	.1	34.0
		% within First use: cocaine	41.2%	58.8%	.0%	100.0%
		% within Gender	6.0%	4.0%	.0%	4.6%
		% of Total	1.9%	2.7%	.0%	4.6%
	21-25	Count	7	4	0	11
		Expected Count	3.5	7.5	.0	11.0
		% within First use: cocaine	63.6%	36.4%	.0%	100.0%
		% within Gender	3.0%	.8%	.0%	1.5%
		% of Total	1.0%	.5%	.0%	1.5%
Total		Count	232	500	2	734
		Expected Count	232.0	500.0	2.0	734.0
		% within First use: cocaine	31.6%	68.1%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.6%	68.1%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	8.024(a)	10	.627
Likelihood Ratio	7.717	10	.656
Linear-by-Linear Association	3.949	1	.047
N of Valid Cases	734		

a 11 cells (61.1%) have expected count less than 5. The minimum expected count is .01.

# First use: amphetamines \* Gender

Crosstab

		Gender			Total	
		Male	Female	Unknown		
First use: amphetamines	Did not use	Count	202	398	2	602
		Expected Count	190.0	410.3	1.6	602.0
		% within First use: amphetamines	33.6%	66.1%	.3%	100.0%
		% within Gender	87.1%	79.4%	100.0%	81.9%
		% of Total	27.5%	54.1%	.3%	81.9%
	10-11	Count	1	1	0	2
		Expected Count	.6	1.4	.0	2.0
		% within First use: amphetamines	50.0%	50.0%	.0%	100.0%
		% within Gender	.4%	.2%	.0%	.3%
		% of Total	.1%	.1%	.0%	.3%
	12-13	Count	2	2	0	4
		Expected Count	1.3	2.7	.0	4.0
		% within First use: amphetamines	50.0%	50.0%	.0%	100.0%
		% within Gender	.9%	.4%	.0%	.5%
		% of Total	.3%	.3%	.0%	.5%
	14-15	Count	4	18	0	22
		Expected Count	6.9	15.0	.1	22.0
		% within First use: amphetamines	18.2%	81.8%	.0%	100.0%
		% within Gender	1.7%	3.6%	.0%	3.0%
		% of Total	.5%	2.4%	.0%	3.0%
	16-17	Count	9	42	0	51
		Expected Count	16.1	34.8	.1	51.0
		% within First use: amphetamines	17.6%	82.4%	.0%	100.0%
		% within Gender	3.9%	8.4%	.0%	6.9%
		% of Total	1.2%	5.7%	.0%	6.9%
	18-20	Count	12	37	0	49
		Expected Count	15.5	33.4	.1	49.0
		% within First use: amphetamines	24.5%	75.5%	.0%	100.0%

		% within Gender	5.2%	7.4%	.0%	6.7%
		% of Total	1.6%	5.0%	.0%	6.7%
21-25	Count	2	3	0	5	
	Expected Count	1.6	3.4	.0	5.0	
	% within First use: amphetamines	40.0%	60.0%	.0%	100.0%	
	% within Gender	.9%	.6%	.0%	.7%	
	% of Total	.3%	.4%	.0%	.7%	
Total	Count	232	501	2	735	
	Expected Count	232.0	501.0	2.0	735.0	
	% within First use: amphetamines	31.6%	68.2%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	31.6%	68.2%	.3%	100.0%	

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	10.318(a)	12	.588
Likelihood Ratio	11.336	12	.500
Linear-by-Linear Association	5.424	1	.020
N of Valid Cases	735		

a 13 cells (61.9%) have expected count less than 5. The minimum expected count is .01.

## First use: sedatives \* Gender

### Crosstab

			Gender			Total
			Male	Female	Unknown	
Did not use	Count	215	476	2	693	
	Expected Count	218.0	473.1	1.9	693.0	
	% within First use: sedatives	31.0%	68.7%	.3%	100.0%	
	% within Gender	93.5%	95.4%	100.0%	94.8%	
	% of Total	29.4%	65.1%	.3%	94.8%	
12-13	Count	2	0	0	2	
	Expected Count	.6	1.4	.0	2.0	
	% within First use: sedatives	100.0%	.0%	.0%	100.0%	
	% within Gender	.9%	.0%	.0%	.3%	
	% of Total	.3%	.0%	.0%	.3%	
	Count	2	5	0	7	
	Expected Count	2.2	4.8	.0	7.0	

		% within First use: sedatives	28.6%	71.4%	.0%	100.0%	
		% within Gender	.9%	1.0%	.0%	1.0%	
		% of Total	.3%	.7%	.0%	1.0%	
First use: sedatives	14-15	Count	4	6	0	10	
		Expected Count	3.1	6.8	.0	10.0	
		% within First use: sedatives	40.0%	60.0%	.0%	100.0%	
		% within Gender	1.7%	1.2%	.0%	1.4%	
		% of Total	.5%	.8%	.0%	1.4%	
	16-17	Count	6	12	0	18	
		Expected Count	5.7	12.3	.0	18.0	
		% within First use: sedatives	33.3%	66.7%	.0%	100.0%	
		% within Gender	2.6%	2.4%	.0%	2.5%	
		% of Total	.8%	1.6%	.0%	2.5%	
	18-20	Count	1	0	0	1	
		Expected Count	.3	.7	.0	1.0	
		% within First use: sedatives	100.0%	.0%	.0%	100.0%	
		% within Gender	.4%	.0%	.0%	.1%	
		% of Total	.1%	.0%	.0%	.1%	
	26+	Count	230	499	2	731	
		Expected Count	230.0	499.0	2.0	731.0	
		% within First use: sedatives	31.5%	68.3%	.3%	100.0%	
		% within Gender	100.0%	100.0%	100.0%	100.0%	
		% of Total	31.5%	68.3%	.3%	100.0%	
Total		Count	230	499	2	731	
		Expected Count	230.0	499.0	2.0	731.0	
		% within First use: sedatives	31.5%	68.3%	.3%	100.0%	
		% within Gender	100.0%	100.0%	100.0%	100.0%	
		% of Total	31.5%	68.3%	.3%	100.0%	

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	7.084(a)	10	.717
Likelihood Ratio	7.569	10	.671
Linear-by-Linear Association	1.067	1	.302
N of Valid Cases	731		

a 13 cells (72.2%) have expected count less than 5. The minimum expected count is .00.

## First use: hallucinogens \* Gender

#### Crosstab

		Gender			Total
		Male	Female	Unknown	
		Count	203	447	2
		Expected Count	204.3	445.9	1.8

First use: hallucinogens	Did not use	% within First use: hallucinogens	31.1%	68.6%	.3%	100.0%
		% within Gender	88.3%	89.0%	100.0%	88.8%
		% of Total	27.7%	60.9%	.3%	88.8%
	Under 10	Count	0	3	0	3
		Expected Count	.9	2.1	.0	3.0
		% within First use: hallucinogens	.0%	100.0%	.0%	100.0%
		% within Gender	.0%	.6%	.0%	.4%
		% of Total	.0%	.4%	.0%	.4%
	12-13	Count	0	1	0	1
		Expected Count	.3	.7	.0	1.0
		% within First use: hallucinogens	.0%	100.0%	.0%	100.0%
		% within Gender	.0%	.2%	.0%	.1%
		% of Total	.0%	.1%	.0%	.1%
	14-15	Count	3	10	0	13
		Expected Count	4.1	8.9	.0	13.0
		% within First use: hallucinogens	23.1%	76.9%	.0%	100.0%
		% within Gender	1.3%	2.0%	.0%	1.8%
		% of Total	.4%	1.4%	.0%	1.8%
	16-17	Count	9	15	0	24
		Expected Count	7.5	16.4	.1	24.0
		% within First use: hallucinogens	37.5%	62.5%	.0%	100.0%
		% within Gender	3.9%	3.0%	.0%	3.3%
		% of Total	1.2%	2.0%	.0%	3.3%
	18-20	Count	13	20	0	33
		Expected Count	10.3	22.6	.1	33.0
		% within First use: hallucinogens	39.4%	60.6%	.0%	100.0%
		% within Gender	5.7%	4.0%	.0%	4.5%
		% of Total	1.8%	2.7%	.0%	4.5%
	21-25	Count	2	6	0	8
		Expected Count	2.5	5.5	.0	8.0
		% within First use: hallucinogens	25.0%	75.0%	.0%	100.0%
		% within Gender	.9%	1.2%	.0%	1.1%
		% of Total	.3%	.8%	.0%	1.1%
Total		Count	230	502	2	734
		Expected Count	230.0	502.0	2.0	734.0
		% within First use: hallucinogens	31.3%	68.4%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.3%	68.4%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	4.064(a)	12	.982
<b>Likelihood Ratio</b>	5.442	12	.942
<b>Linear-by-Linear Association</b>	.465	1	.495
<b>N of Valid Cases</b>	734		

a 13 cells (61.9%) have expected count less than 5. The minimum expected count is .00.

## First use: opiates \* Gender

Crosstab

		Gender			Total	
		Male	Female	Unknown		
First use: opiates	Did not use	<b>Count</b>	221	484	2	707
		<b>Expected Count</b>	223.4	481.6	1.9	707.0
		<b>% within First use: opiates</b>	31.3%	68.5%	.3%	100.0%
		<b>% within Gender</b>	95.7%	97.2%	100.0%	96.7%
		<b>% of Total</b>	30.2%	66.2%	.3%	96.7%
	14-15	<b>Count</b>	2	3	0	5
		<b>Expected Count</b>	1.6	3.4	.0	5.0
		<b>% within First use: opiates</b>	40.0%	60.0%	.0%	100.0%
		<b>% within Gender</b>	.9%	.6%	.0%	.7%
		<b>% of Total</b>	.3%	.4%	.0%	.7%
	16-17	<b>Count</b>	3	4	0	7
		<b>Expected Count</b>	2.2	4.8	.0	7.0
		<b>% within First use: opiates</b>	42.9%	57.1%	.0%	100.0%
		<b>% within Gender</b>	1.3%	.8%	.0%	1.0%
		<b>% of Total</b>	.4%	.5%	.0%	1.0%
	18-20	<b>Count</b>	5	7	0	12
		<b>Expected Count</b>	3.8	8.2	.0	12.0
		<b>% within First use: opiates</b>	41.7%	58.3%	.0%	100.0%
		<b>% within Gender</b>	2.2%	1.4%	.0%	1.6%
		<b>% of Total</b>	.7%	1.0%	.0%	1.6%
	Total	<b>Count</b>	231	498	2	731
		<b>Expected Count</b>	231.0	498.0	2.0	731.0
		<b>% within First use: opiates</b>	31.6%	68.1%	.3%	100.0%
		<b>% within Gender</b>	100.0%	100.0%	100.0%	100.0%
		<b>% of Total</b>	31.6%	68.1%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	1.224(a)	6	.976
<b>Likelihood Ratio</b>	1.235	6	.975
<b>Linear-by-Linear Association</b>	1.189	1	.276
<b>N of Valid Cases</b>	731		

a 9 cells (75.0%) have expected count less than 5. The minimum expected count is .01.

## First use: inhalants \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
First use: inhalants	Did not use	<b>Count</b>	214	474	2	690
		<b>Expected Count</b>	217.4	470.7	1.9	690.0
		<b>% within First use: inhalants</b>	31.0%	68.7%	.3%	100.0%
		<b>% within Gender</b>	93.0%	95.2%	100.0%	94.5%
		<b>% of Total</b>	29.3%	64.9%	.3%	94.5%
	10-11	<b>Count</b>	1	1	0	2
		<b>Expected Count</b>	.6	1.4	.0	2.0
		<b>% within First use: inhalants</b>	50.0%	50.0%	.0%	100.0%
		<b>% within Gender</b>	.4%	.2%	.0%	.3%
		<b>% of Total</b>	.1%	.1%	.0%	.3%
	12-13	<b>Count</b>	2	2	0	4
		<b>Expected Count</b>	1.3	2.7	.0	4.0
		<b>% within First use: inhalants</b>	50.0%	50.0%	.0%	100.0%
		<b>% within Gender</b>	.9%	.4%	.0%	.5%
		<b>% of Total</b>	.3%	.3%	.0%	.5%
	14-15	<b>Count</b>	5	7	0	12
		<b>Expected Count</b>	3.8	8.2	.0	12.0
		<b>% within First use: inhalants</b>	41.7%	58.3%	.0%	100.0%
		<b>% within Gender</b>	2.2%	1.4%	.0%	1.6%
		<b>% of Total</b>	.7%	1.0%	.0%	1.6%
	16-17	<b>Count</b>	1	6	0	7
		<b>Expected Count</b>	2.2	4.8	.0	7.0
		<b>% within First use: inhalants</b>	14.3%	85.7%	.0%	100.0%
		<b>% within Gender</b>	.4%	1.2%	.0%	1.0%
		<b>% of Total</b>	.1%	.8%	.0%	1.0%
		<b>Count</b>	7	8	0	15
		<b>Expected Count</b>	4.7	10.2	.0	15.0

			% within First use: inhalants	46.7%	53.3%	.0%	100.0%
		18-20	% within Gender	3.0%	1.6%	.0%	2.1%
			% of Total	1.0%	1.1%	.0%	2.1%
	Total		Count	230	498	2	730
			Expected Count	230.0	498.0	2.0	730.0
			% within First use: inhalants	31.5%	68.2%	.3%	100.0%
			% within Gender	100.0%	100.0%	100.0%	100.0%
			% of Total	31.5%	68.2%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	4.255(a)	10	.935
Likelihood Ratio	4.311	10	.932
Linear-by-Linear Association	1.282	1	.258
N of Valid Cases	730		

a 14 cells (77.8%) have expected count less than 5. The minimum expected count is .01.

## First use: designer \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
First use: designer	Did not use	Count	217	453	2	672
		Expected Count	212.1	458.1	1.8	672.0
		% within First use: designer	32.3%	67.4%	.3%	100.0%
		% within Gender	93.5%	90.4%	100.0%	91.4%
		% of Total	29.5%	61.6%	.3%	91.4%
	16-17	Count	3	22	0	25
		Expected Count	7.9	17.0	.1	25.0
		% within First use: designer	12.0%	88.0%	.0%	100.0%
		% within Gender	1.3%	4.4%	.0%	3.4%
		% of Total	.4%	3.0%	.0%	3.4%
	18-20	Count	10	26	0	36
		Expected Count	11.4	24.5	.1	36.0
		% within First use: designer	27.8%	72.2%	.0%	100.0%
		% within Gender	4.3%	5.2%	.0%	4.9%
		% of Total	1.4%	3.5%	.0%	4.9%
		Count	2	0	0	2
		Expected Count	.6	1.4	.0	2.0

		% within First use: designer	100.0%	.0%	.0%	100.0%
	21-25	% within Gender	.9%	.0%	.0%	.3%
		% of Total	.3%	.0%	.0%	.3%
Total		Count	232	501	2	735
		Expected Count	232.0	501.0	2.0	735.0
		% within First use: designer	31.6%	68.2%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.6%	68.2%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	9.412(a)	6	.152
Likelihood Ratio	10.684	6	.099
Linear-by-Linear Association	1.196	1	.274
N of Valid Cases	735		

a 6 cells (50.0%) have expected count less than 5. The minimum expected count is .01.

## First use: steroids \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
First use: steroids	Did not use	Count	229	499	2	730
		Expected Count	229.1	498.9	2.0	730.0
		% within First use: steroids	31.4%	68.4%	.3%	100.0%
		% within Gender	99.6%	99.6%	100.0%	99.6%
		% of Total	31.2%	68.1%	.3%	99.6%
	12-13	Count	1	0	0	1
		Expected Count	.3	.7	.0	1.0
		% within First use: steroids	100.0%	.0%	.0%	100.0%
		% within Gender	.4%	.0%	.0%	.1%
		% of Total	.1%	.0%	.0%	.1%
	14-15	Count	0	1	0	1
		Expected Count	.3	.7	.0	1.0
		% within First use: steroids	.0%	100.0%	.0%	100.0%
		% within Gender	.0%	.2%	.0%	.1%
		% of Total	.0%	.1%	.0%	.1%
		Count	0	1	0	1
		Expected Count	.3	.7	.0	1.0

			<b>% within First use: steroids</b>	.0%	100.0%	.0%	100.0%
		<b>16-17</b>	<b>% within Gender</b>	.0%	.2%	.0%	.1%
			<b>% of Total</b>	.0%	.1%	.0%	.1%
			<b>Count</b>	230	501	2	733
			<b>Expected Count</b>	230.0	501.0	2.0	733.0
			<b>% within First use: steroids</b>	31.4%	68.3%	.3%	100.0%
			<b>% within Gender</b>	100.0%	100.0%	100.0%	100.0%
			<b>% of Total</b>	31.4%	68.3%	.3%	100.0%

### Chi-Square Tests

	<b>Value</b>	<b>df</b>	<b>Asymp. Sig. (2-sided)</b>
<b>Pearson Chi-Square</b>	3.113(a)	6	.795
<b>Likelihood Ratio</b>	3.840	6	.698
<b>Linear-by-Linear Association</b>	.049	1	.825
<b>N of Valid Cases</b>	733		

a 10 cells (83.3%) have expected count less than 5. The minimum expected count is .00.

## First use: other \* Gender

### Crosstab

			<b>Gender</b>			<b>Total</b>
			<b>Male</b>	<b>Female</b>	<b>Unknown</b>	
<b>First use: other</b>	<b>Did not use</b>	<b>Count</b>	218	476	2	696
		<b>Expected Count</b>	219.7	474.4	1.9	696.0
		<b>% within First use: other</b>	31.3%	68.4%	.3%	100.0%
		<b>% within Gender</b>	94.0%	95.0%	100.0%	94.7%
		<b>% of Total</b>	29.7%	64.8%	.3%	94.7%
	<b>12-13</b>	<b>Count</b>	0	1	0	1
		<b>Expected Count</b>	.3	.7	.0	1.0
		<b>% within First use: other</b>	.0%	100.0%	.0%	100.0%
		<b>% within Gender</b>	.0%	.2%	.0%	.1%
		<b>% of Total</b>	.0%	.1%	.0%	.1%
	<b>14-15</b>	<b>Count</b>	2	3	0	5
		<b>Expected Count</b>	1.6	3.4	.0	5.0
		<b>% within First use: other</b>	40.0%	60.0%	.0%	100.0%
		<b>% within Gender</b>	.9%	.6%	.0%	.7%
		<b>% of Total</b>	.3%	.4%	.0%	.7%
		<b>Count</b>	2	7	0	9
		<b>Expected Count</b>	2.8	6.1	.0	9.0

		% within First use: other	22.2%	77.8%	.0%	100.0%	
		% within Gender	.9%	1.4%	.0%	1.2%	
		% of Total	.3%	1.0%	.0%	1.2%	
16-17	18-20	Count	9	14	0	23	
		Expected Count	7.3	15.7	.1	23.0	
		% within First use: other	39.1%	60.9%	.0%	100.0%	
		% within Gender	3.9%	2.8%	.0%	3.1%	
		% of Total	1.2%	1.9%	.0%	3.1%	
21-25	Total	Count	1	0	0	1	
		Expected Count	.3	.7	.0	1.0	
		% within First use: other	100.0%	.0%	.0%	100.0%	
		% within Gender	.4%	.0%	.0%	.1%	
		% of Total	.1%	.0%	.0%	.1%	
		Count	232	501	2	735	
		Expected Count	232.0	501.0	2.0	735.0	
		% within First use: other	31.6%	68.2%	.3%	100.0%	
		% within Gender	100.0%	100.0%	100.0%	100.0%	
		% of Total	31.6%	68.2%	.3%	100.0%	

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	3.889(a)	10	.952
Likelihood Ratio	4.421	10	.926
Linear-by-Linear Association	.612	1	.434
N of Valid Cases	735		

a 13 cells (72.2%) have expected count less than 5. The minimum expected count is .00.

## Use last yr: tobacco \* Gender

#### Crosstab

			Gender			Total
			Male	Female	Unknown	
Never used		Count	98	272	0	370
		Expected Count	118.0	251.0	1.0	370.0
		% within Use last yr: tobacco	26.5%	73.5%	.0%	100.0%
		% within Gender	41.7%	54.4%	.0%	50.2%
		% of Total	13.3%	36.9%	.0%	50.2%
		Count	18	29	1	48
		Expected Count	15.3	32.6	.1	48.0

Use last yr: tobacco	Once/year	% within Use last yr: tobacco	37.5%	60.4%	2.1%	100.0%	
		% within Gender	7.7%	5.8%	50.0%	6.5%	
		% of Total	2.4%	3.9%	.1%	6.5%	
	6 times/year	Count	20	49	0	69	
		Expected Count	22.0	46.8	.2	69.0	
		% within Use last yr: tobacco	29.0%	71.0%	.0%	100.0%	
		% within Gender	8.5%	9.8%	.0%	9.4%	
		% of Total	2.7%	6.6%	.0%	9.4%	
	Once/month	Count	12	13	0	25	
		Expected Count	8.0	17.0	.1	25.0	
		% within Use last yr: tobacco	48.0%	52.0%	.0%	100.0%	
		% within Gender	5.1%	2.6%	.0%	3.4%	
		% of Total	1.6%	1.8%	.0%	3.4%	
	Twice/month	Count	11	18	0	29	
		Expected Count	9.2	19.7	.1	29.0	
		% within Use last yr: tobacco	37.9%	62.1%	.0%	100.0%	
		% within Gender	4.7%	3.6%	.0%	3.9%	
		% of Total	1.5%	2.4%	.0%	3.9%	
	Once/week	Count	13	17	0	30	
		Expected Count	9.6	20.4	.1	30.0	
		% within Use last yr: tobacco	43.3%	56.7%	.0%	100.0%	
		% within Gender	5.5%	3.4%	.0%	4.1%	
		% of Total	1.8%	2.3%	.0%	4.1%	
	3 times/week	Count	11	14	0	25	
		Expected Count	8.0	17.0	.1	25.0	
		% within Use last yr: tobacco	44.0%	56.0%	.0%	100.0%	
		% within Gender	4.7%	2.8%	.0%	3.4%	
		% of Total	1.5%	1.9%	.0%	3.4%	
	5 times/week	Count	9	19	0	28	
		Expected Count	8.9	19.0	.1	28.0	
		% within Use last yr: tobacco	32.1%	67.9%	.0%	100.0%	
		% within Gender	3.8%	3.8%	.0%	3.8%	
		% of Total	1.2%	2.6%	.0%	3.8%	
	Every day	Count	43	69	1	113	
		Expected Count	36.0	76.7	.3	113.0	
		% within Use last yr: tobacco	38.1%	61.1%	.9%	100.0%	
		% within Gender	18.3%	13.8%	50.0%	15.3%	
		% of Total	5.8%	9.4%	.1%	15.3%	
		Count	235	500	2	737	
		Expected Count	235.0	500.0	2.0	737.0	

Total	% within Use last yr: tobacco	31.9%	67.8%	.3%	100.0%
	% within Gender	100.0%	100.0%	100.0%	100.0%
	% of Total	31.9%	67.8%	.3%	100.0%

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	24.227(a)	16	.085
<b>Likelihood Ratio</b>	21.440	16	.162
<b>Linear-by-Linear Association</b>	7.143	1	.008
<b>N of Valid Cases</b>	737		

a 9 cells (33.3%) have expected count less than 5. The minimum expected count is .07.

**Use last yr: alcohol \* Gender****Crosstab**

			Gender			Total
			Male	Female	Unknown	
Never used	Never used	<b>Count</b>	17	40	0	57
		<b>Expected Count</b>	18.2	38.6	.2	57.0
		<b>% within Use last yr: alcohol</b>	29.8%	70.2%	.0%	100.0%
		<b>% within Gender</b>	7.2%	8.0%	.0%	7.7%
		<b>% of Total</b>	2.3%	5.4%	.0%	7.7%
Once/year	Once/year	<b>Count</b>	9	18	0	27
		<b>Expected Count</b>	8.6	18.3	.1	27.0
		<b>% within Use last yr: alcohol</b>	33.3%	66.7%	.0%	100.0%
		<b>% within Gender</b>	3.8%	3.6%	.0%	3.7%
		<b>% of Total</b>	1.2%	2.4%	.0%	3.7%
6 times/year	6 times/year	<b>Count</b>	9	50	0	59
		<b>Expected Count</b>	18.8	40.0	.2	59.0
		<b>% within Use last yr: alcohol</b>	15.3%	84.7%	.0%	100.0%
		<b>% within Gender</b>	3.8%	10.0%	.0%	8.0%
		<b>% of Total</b>	1.2%	6.8%	.0%	8.0%
Once/month	Once/month	<b>Count</b>	15	39	1	55
		<b>Expected Count</b>	17.6	37.3	.1	55.0
		<b>% within Use last yr: alcohol</b>	27.3%	70.9%	1.8%	100.0%
		<b>% within Gender</b>	6.4%	7.8%	50.0%	7.5%
		<b>% of Total</b>	2.0%	5.3%	.1%	7.5%
		<b>Count</b>	20	80	0	100
		<b>Expected Count</b>	31.9	67.8	.3	100.0

Use last yr: alcohol	Twice/month	% within Use last yr: alcohol	20.0%	80.0%	.0%	100.0%
		% within Gender	8.5%	16.0%	.0%	13.6%
		% of Total	2.7%	10.9%	.0%	13.6%
	Once/week	Count	62	147	0	209
		Expected Count	66.7	141.7	.6	209.0
		% within Use last yr: alcohol	29.7%	70.3%	.0%	100.0%
		% within Gender	26.4%	29.5%	.0%	28.4%
		% of Total	8.4%	20.0%	.0%	28.4%
	3 times/week	Count	87	116	0	203
		Expected Count	64.8	137.6	.6	203.0
		% within Use last yr: alcohol	42.9%	57.1%	.0%	100.0%
		% within Gender	37.0%	23.2%	.0%	27.6%
		% of Total	11.8%	15.8%	.0%	27.6%
	5 times/week	Count	14	8	1	23
		Expected Count	7.3	15.6	.1	23.0
		% within Use last yr: alcohol	60.9%	34.8%	4.3%	100.0%
		% within Gender	6.0%	1.6%	50.0%	3.1%
		% of Total	1.9%	1.1%	.1%	3.1%
	Every day	Count	2	1	0	3
		Expected Count	1.0	2.0	.0	3.0
		% within Use last yr: alcohol	66.7%	33.3%	.0%	100.0%
		% within Gender	.9%	.2%	.0%	.4%
		% of Total	.3%	.1%	.0%	.4%
Total		Count	235	499	2	736
		Expected Count	235.0	499.0	2.0	736.0
		% within Use last yr: alcohol	31.9%	67.8%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.9%	67.8%	.3%	100.0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	58.498(a)	16	.000
Likelihood Ratio	47.635	16	.000
Linear-by-Linear Association	13.185	1	.000
N of Valid Cases	736		

a 11 cells (40.7%) have expected count less than 5. The minimum expected count is .01.

## Use last yr: marijuana \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Use last yr: marijuana	Never used	Count	141	333	2	476
		Expected Count	152.0	322.7	1.3	476.0
		% within Use last yr: marijuana	29.6%	70.0%	.4%	100.0%
		% within Gender	60.0%	66.7%	100.0%	64.7%
		% of Total	19.2%	45.2%	.3%	64.7%
	Once/year	Count	31	53	0	84
		Expected Count	26.8	57.0	.2	84.0
		% within Use last yr: marijuana	36.9%	63.1%	.0%	100.0%
		% within Gender	13.2%	10.6%	.0%	11.4%
		% of Total	4.2%	7.2%	.0%	11.4%
	6 times/year	Count	23	39	0	62
		Expected Count	19.8	42.0	.2	62.0
		% within Use last yr: marijuana	37.1%	62.9%	.0%	100.0%
		% within Gender	9.8%	7.8%	.0%	8.4%
		% of Total	3.1%	5.3%	.0%	8.4%
	Once/month	Count	8	15	0	23
		Expected Count	7.3	15.6	.1	23.0
		% within Use last yr: marijuana	34.8%	65.2%	.0%	100.0%
		% within Gender	3.4%	3.0%	.0%	3.1%
		% of Total	1.1%	2.0%	.0%	3.1%
	Twice/month	Count	5	16	0	21
		Expected Count	6.7	14.2	.1	21.0
		% within Use last yr: marijuana	23.8%	76.2%	.0%	100.0%
		% within Gender	2.1%	3.2%	.0%	2.9%
		% of Total	.7%	2.2%	.0%	2.9%
	Once/week	Count	7	11	0	18
		Expected Count	5.7	12.2	.0	18.0
		% within Use last yr: marijuana	38.9%	61.1%	.0%	100.0%
		% within Gender	3.0%	2.2%	.0%	2.4%
		% of Total	1.0%	1.5%	.0%	2.4%
	3 times/week	Count	7	12	0	19
		Expected Count	6.1	12.9	.1	19.0
		% within Use last yr: marijuana	36.8%	63.2%	.0%	100.0%
		% within Gender	3.0%	2.4%	.0%	2.6%
		% of Total	1.0%	1.6%	.0%	2.6%
		Count	5	10	0	15
		Expected Count	4.8	10.2	.0	15.0

		% within Use last yr: marijuana	33.3%	66.7%	.0%	100.0%
		% within Gender	2.1%	2.0%	.0%	2.0%
		% of Total	.7%	1.4%	.0%	2.0%
Total	Every day	Count	8	10	0	18
		Expected Count	5.7	12.2	.0	18.0
		% within Use last yr: marijuana	44.4%	55.6%	.0%	100.0%
		% within Gender	3.4%	2.0%	.0%	2.4%
		% of Total	1.1%	1.4%	.0%	2.4%
Total	Total	Count	235	499	2	736
		Expected Count	235.0	499.0	2.0	736.0
		% within Use last yr: marijuana	31.9%	67.8%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.9%	67.8%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	6.498(a)	16	.982
Likelihood Ratio	7.063	16	.972
Linear-by-Linear Association	2.322	1	.128
N of Valid Cases	736		

a 10 cells (37.0%) have expected count less than 5. The minimum expected count is .04.

## Use last yr: cocaine \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
Never used	Never used	Count	217	472	2	691
		Expected Count	220.0	469.1	1.9	691.0
		% within Use last yr: cocaine	31.4%	68.3%	.3%	100.0%
		% within Gender	92.7%	94.6%	100.0%	94.0%
		% of Total	29.5%	64.2%	.3%	94.0%
Once/year	Once/year	Count	10	12	0	22
		Expected Count	7.0	14.9	.1	22.0
		% within Use last yr: cocaine	45.5%	54.5%	.0%	100.0%
		% within Gender	4.3%	2.4%	.0%	3.0%
		% of Total	1.4%	1.6%	.0%	3.0%
		Count	2	9	0	11
		Expected Count	3.5	7.5	.0	11.0

Use last yr: cocaine	6 times/year	% within Use last yr: cocaine	18.2%	81.8%	.0%	100.0%
		% within Gender	.9%	1.8%	.0%	1.5%
		% of Total	.3%	1.2%	.0%	1.5%
	Once/month	Count	3	1	0	4
		Expected Count	1.3	2.7	.0	4.0
		% within Use last yr: cocaine	75.0%	25.0%	.0%	100.0%
		% within Gender	1.3%	.2%	.0%	.5%
		% of Total	.4%	.1%	.0%	.5%
	Twice/month	Count	1	3	0	4
		Expected Count	1.3	2.7	.0	4.0
		% within Use last yr: cocaine	25.0%	75.0%	.0%	100.0%
		% within Gender	.4%	.6%	.0%	.5%
		% of Total	.1%	.4%	.0%	.5%
	Once/week	Count	1	0	0	1
		Expected Count	.3	.7	.0	1.0
		% within Use last yr: cocaine	100.0%	.0%	.0%	100.0%
		% within Gender	.4%	.0%	.0%	.1%
		% of Total	.1%	.0%	.0%	.1%
	3 times/week	Count	0	1	0	1
		Expected Count	.3	.7	.0	1.0
		% within Use last yr: cocaine	.0%	100.0%	.0%	100.0%
		% within Gender	.0%	.2%	.0%	.1%
		% of Total	.0%	.1%	.0%	.1%
	Every day	Count	0	1	0	1
		Expected Count	.3	.7	.0	1.0
		% within Use last yr: cocaine	.0%	100.0%	.0%	100.0%
		% within Gender	.0%	.2%	.0%	.1%
		% of Total	.0%	.1%	.0%	.1%
Total	Count	234	499	2	735	
	Expected Count	234.0	499.0	2.0	735.0	
	% within Use last yr: cocaine	31.8%	67.9%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	31.8%	67.9%	.3%	100.0%	

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	9.595(a)	14	.791
Likelihood Ratio	10.148	14	.751
Linear-by-Linear Association	.173	1	.677
N of Valid Cases	735		

a 19 cells (79.2%) have expected count less than 5. The minimum expected count is .00.

## Use last yr: amphetamines \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Use last yr: amphetamines	Never used	Count	220	455	2	677
		Expected Count	216.5	458.7	1.8	677.0
		% within Use last yr: amphetamines	32.5%	67.2%	.3%	100.0%
		% within Gender	93.6%	91.4%	100.0%	92.1%
		% of Total	29.9%	61.9%	.3%	92.1%
	Once/year	Count	4	15	0	19
		Expected Count	6.1	12.9	.1	19.0
		% within Use last yr: amphetamines	21.1%	78.9%	.0%	100.0%
		% within Gender	1.7%	3.0%	.0%	2.6%
		% of Total	.5%	2.0%	.0%	2.6%
	6 times/year	Count	5	8	0	13
		Expected Count	4.2	8.8	.0	13.0
		% within Use last yr: amphetamines	38.5%	61.5%	.0%	100.0%
		% within Gender	2.1%	1.6%	.0%	1.8%
		% of Total	.7%	1.1%	.0%	1.8%
	Once/month	Count	1	3	0	4
		Expected Count	1.3	2.7	.0	4.0
		% within Use last yr: amphetamines	25.0%	75.0%	.0%	100.0%
		% within Gender	.4%	.6%	.0%	.5%
		% of Total	.1%	.4%	.0%	.5%
	Twice/month	Count	1	3	0	4
		Expected Count	1.3	2.7	.0	4.0
		% within Use last yr: amphetamines	25.0%	75.0%	.0%	100.0%
		% within Gender	.4%	.6%	.0%	.5%
		% of Total	.1%	.4%	.0%	.5%
	Once/week	Count	1	1	0	2
		Expected Count	.6	1.4	.0	2.0
		% within Use last yr: amphetamines	50.0%	50.0%	.0%	100.0%

		% within Gender	.4%	.2%	.0%	.3%
		% of Total	.1%	.1%	.0%	.3%
3 times/week	Count	0	4	0	4	
	Expected Count	1.3	2.7	.0	4.0	
	% within Use last yr: amphetamines	.0%	100.0%	.0%	100.0%	
	% within Gender	.0%	.8%	.0%	.5%	
	% of Total	.0%	.5%	.0%	.5%	
5 times/week	Count	2	3	0	5	
	Expected Count	1.6	3.4	.0	5.0	
	% within Use last yr: amphetamines	40.0%	60.0%	.0%	100.0%	
	% within Gender	.9%	.6%	.0%	.7%	
	% of Total	.3%	.4%	.0%	.7%	
Every day	Count	1	6	0	7	
	Expected Count	2.2	4.7	.0	7.0	
	% within Use last yr: amphetamines	14.3%	85.7%	.0%	100.0%	
	% within Gender	.4%	1.2%	.0%	1.0%	
	% of Total	.1%	.8%	.0%	1.0%	
Total	Count	235	498	2	735	
	Expected Count	235.0	498.0	2.0	735.0	
	% within Use last yr: amphetamines	32.0%	67.8%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	32.0%	67.8%	.3%	100.0%	

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	5.100(a)	16	.995
Likelihood Ratio	6.665	16	.979
Linear-by-Linear Association	.977	1	.323
N of Valid Cases	735		

a 22 cells (81.5%) have expected count less than 5. The minimum expected count is .01.

## Use last yr: sedatives \* Gender

## Crosstab

	Gender			Total
	Male	Female	Unknown	
Count	229	485	2	716

Use last yr: sedatives	Never used	Expected Count	228.6	485.5	2.0	716.0
		% within Use last yr: sedatives	32.0%	67.7%	.3%	100.0%
		% within Gender	97.9%	97.6%	100.0%	97.7%
		% of Total	31.2%	66.2%	.3%	97.7%
	Once/year	Count	0	6	0	6
		Expected Count	1.9	4.1	.0	6.0
		% within Use last yr: sedatives	.0%	100.0%	.0%	100.0%
		% within Gender	.0%	1.2%	.0%	.8%
		% of Total	.0%	.8%	.0%	.8%
	6 times/year	Count	5	2	0	7
		Expected Count	2.2	4.7	.0	7.0
		% within Use last yr: sedatives	71.4%	28.6%	.0%	100.0%
		% within Gender	2.1%	.4%	.0%	1.0%
		% of Total	.7%	.3%	.0%	1.0%
	Once/month	Count	0	2	0	2
		Expected Count	.6	1.4	.0	2.0
		% within Use last yr: sedatives	.0%	100.0%	.0%	100.0%
		% within Gender	.0%	.4%	.0%	.3%
		% of Total	.0%	.3%	.0%	.3%
	Once/week	Count	0	1	0	1
		Expected Count	.3	.7	.0	1.0
		% within Use last yr: sedatives	.0%	100.0%	.0%	100.0%
		% within Gender	.0%	.2%	.0%	.1%
		% of Total	.0%	.1%	.0%	.1%
	3 times/week	Count	0	1	0	1
		Expected Count	.3	.7	.0	1.0
		% within Use last yr: sedatives	.0%	100.0%	.0%	100.0%
		% within Gender	.0%	.2%	.0%	.1%
		% of Total	.0%	.1%	.0%	.1%
	Total	Count	234	497	2	733
		Expected Count	234.0	497.0	2.0	733.0
		% within Use last yr: sedatives	31.9%	67.8%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.9%	67.8%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	9.781(a)	10	.460
Likelihood Ratio	12.370	10	.261
Linear-by-Linear Association	.119	1	.731

N of Valid Cases	733	
a 16 cells (88.9%) have expected count less than 5. The minimum expected count is .00.		

## Use last yr: hallucinogens \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Use last yr: hallucinogens	Never used	Count	219	474	1	694
		Expected Count	222.8	470.3	.9	694.0
		% within Use last yr: hallucinogens	31.6%	68.3%	.1%	100.0%
		% within Gender	93.2%	95.6%	100.0%	94.8%
		% of Total	29.9%	64.8%	.1%	94.8%
	Once/year	Count	11	15	0	26
		Expected Count	8.3	17.6	.0	26.0
		% within Use last yr: hallucinogens	42.3%	57.7%	.0%	100.0%
		% within Gender	4.7%	3.0%	.0%	3.6%
		% of Total	1.5%	2.0%	.0%	3.6%
	6 times/year	Count	5	7	0	12
		Expected Count	3.9	8.1	.0	12.0
		% within Use last yr: hallucinogens	41.7%	58.3%	.0%	100.0%
		% within Gender	2.1%	1.4%	.0%	1.6%
		% of Total	.7%	1.0%	.0%	1.6%
Total	Count	235	496	1	732	
	Expected Count	235.0	496.0	1.0	732.0	
	% within Use last yr: hallucinogens	32.1%	67.8%	.1%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	32.1%	67.8%	.1%	100.0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1.881(a)	4	.758
Likelihood Ratio	1.855	4	.762
Linear-by-Linear Association	1.617	1	.204
N of Valid Cases	732		

a 4 cells (44.4%) have expected count less than 5. The minimum expected count is .02.

# Use last yr: opiates \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Use last yr: opiates	Never used	Count	229	491	2	722
		Expected Count	230.2	489.9	2.0	722.0
		% within Use last yr: opiates	31.7%	68.0%	.3%	100.0%
		% within Gender	97.9%	98.6%	100.0%	98.4%
		% of Total	31.2%	66.9%	.3%	98.4%
	Once/year	Count	1	3	0	4
		Expected Count	1.3	2.7	.0	4.0
		% within Use last yr: opiates	25.0%	75.0%	.0%	100.0%
		% within Gender	.4%	.6%	.0%	.5%
		% of Total	.1%	.4%	.0%	.5%
	6 times/year	Count	4	4	0	8
		Expected Count	2.6	5.4	.0	8.0
		% within Use last yr: opiates	50.0%	50.0%	.0%	100.0%
		% within Gender	1.7%	.8%	.0%	1.1%
		% of Total	.5%	.5%	.0%	1.1%
Total		Count	234	498	2	734
		Expected Count	234.0	498.0	2.0	734.0
		% within Use last yr: opiates	31.9%	67.8%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.9%	67.8%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1.331(a)	4	.856
Likelihood Ratio	1.283	4	.864
Linear-by-Linear Association	.912	1	.340
N of Valid Cases	734		

a 6 cells (66.7%) have expected count less than 5. The minimum expected count is .01.

# Use last yr: inhalants \* Gender

Crosstab

	Gender		Total
	Male	Female	

			Male	Female	Unknown	
Use last yr: inhalants	Never used	Count	228	489	2	719
		Expected Count	230.2	486.8	2.0	719.0
		% within Use last yr: inhalants	31.7%	68.0%	.3%	100.0%
		% within Gender	97.0%	98.4%	100.0%	98.0%
		% of Total	31.1%	66.6%	.3%	98.0%
	Once/year	Count	4	3	0	7
		Expected Count	2.2	4.7	.0	7.0
		% within Use last yr: inhalants	57.1%	42.9%	.0%	100.0%
		% within Gender	1.7%	.6%	.0%	1.0%
		% of Total	.5%	.4%	.0%	1.0%
	6 times/year	Count	3	3	0	6
		Expected Count	1.9	4.1	.0	6.0
		% within Use last yr: inhalants	50.0%	50.0%	.0%	100.0%
		% within Gender	1.3%	.6%	.0%	.8%
		% of Total	.4%	.4%	.0%	.8%
	Once/month	Count	0	2	0	2
		Expected Count	.6	1.4	.0	2.0
		% within Use last yr: inhalants	.0%	100.0%	.0%	100.0%
		% within Gender	.0%	.4%	.0%	.3%
		% of Total	.0%	.3%	.0%	.3%
Total	Count	235	497	2	734	
	Expected Count	235.0	497.0	2.0	734.0	
	% within Use last yr: inhalants	32.0%	67.7%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	32.0%	67.7%	.3%	100.0%	

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	3.924(a)	6	.687
Likelihood Ratio	4.337	6	.631
Linear-by-Linear Association	.398	1	.528
N of Valid Cases	734		

a 10 cells (83.3%) have expected count less than 5. The minimum expected count is .01.

## Use last yr: designer \* Gender

### Crosstab

		Gender	Total

			Male	Female	Unknown	
Use last yr: designer	Never used	Count	228	482	2	712
		Expected Count	227.6	482.4	1.9	712.0
		% within Use last yr: designer	32.0%	67.7%	.3%	100.0%
		% within Gender	97.0%	96.8%	100.0%	96.9%
		% of Total	31.0%	65.6%	.3%	96.9%
	Once/year	Count	7	9	0	16
		Expected Count	5.1	10.8	.0	16.0
		% within Use last yr: designer	43.8%	56.3%	.0%	100.0%
		% within Gender	3.0%	1.8%	.0%	2.2%
		% of Total	1.0%	1.2%	.0%	2.2%
	6 times/year	Count	0	5	0	5
		Expected Count	1.6	3.4	.0	5.0
		% within Use last yr: designer	.0%	100.0%	.0%	100.0%
		% within Gender	.0%	1.0%	.0%	.7%
		% of Total	.0%	.7%	.0%	.7%
	Once/month	Count	0	1	0	1
		Expected Count	.3	.7	.0	1.0
		% within Use last yr: designer	.0%	100.0%	.0%	100.0%
		% within Gender	.0%	.2%	.0%	.1%
		% of Total	.0%	.1%	.0%	.1%
	Twice/month	Count	0	1	0	1
		Expected Count	.3	.7	.0	1.0
		% within Use last yr: designer	.0%	100.0%	.0%	100.0%
		% within Gender	.0%	.2%	.0%	.1%
		% of Total	.0%	.1%	.0%	.1%
Total	Count	235	498	2	735	
	Expected Count	235.0	498.0	2.0	735.0	
	% within Use last yr: designer	32.0%	67.8%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	32.0%	67.8%	.3%	100.0%	

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	4.384(a)	8	.821
Likelihood Ratio	6.493	8	.592
Linear-by-Linear Association	.906	1	.341
N of Valid Cases	735		

a 11 cells (73.3%) have expected count less than 5. The minimum expected count is .00.

## Use last yr: steroids \* Gender

Crosstab

		Count	Gender			Total
			Male	Female	Unknown	
Use last yr: steroids	Never used	Count	234	500	2	736
		Expected Count	234.0	500.0	2.0	736.0
		% within Use last yr: steroids	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%
Total		Count	234	500	2	736
		Expected Count	234.0	500.0	2.0	736.0
		% within Use last yr: steroids	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value
Pearson Chi-Square	.(a)
N of Valid Cases	736

a No statistics are computed because Use last yr: steroids is a constant.

## Use last yr: other \* Gender

Crosstab

		Count	Gender			Total
			Male	Female	Unknown	
Use last yr: other	Never used	Count	226	486	2	714
		Expected Count	227.7	484.4	1.9	714.0
		% within Use last yr: other	31.7%	68.1%	.3%	100.0%
		% within Gender	96.2%	97.2%	100.0%	96.9%
		% of Total	30.7%	65.9%	.3%	96.9%
	Once/year	Count	5	8	0	13
		Expected Count	4.1	8.8	.0	13.0
		% within Use last yr: other	38.5%	61.5%	.0%	100.0%
		% within Gender	2.1%	1.6%	.0%	1.8%
		% of Total	.7%	1.1%	.0%	1.8%
		Count	3	5	0	8

		Expected Count	2.6	5.4	.0	8.0
6 times/year	Once/month	% within Use last yr: other	37.5%	62.5%	.0%	100.0%
		% within Gender	1.3%	1.0%	.0%	1.1%
		% of Total	.4%	.7%	.0%	1.1%
		Count	1	1	0	2
Total		Expected Count	.6	1.4	.0	2.0
		% within Use last yr: other	50.0%	50.0%	.0%	100.0%
		% within Gender	.4%	.2%	.0%	.3%
		% of Total	.1%	.1%	.0%	.3%
		Count	235	500	2	737
		Expected Count	235.0	500.0	2.0	737.0
		% within Use last yr: other	31.9%	67.8%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.9%	67.8%	.3%	100.0%

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.747(a)	6	.993
Likelihood Ratio	.776	6	.993
Linear-by-Linear Association	.633	1	.426
N of Valid Cases	737		

a 8 cells (66.7%) have expected count less than 5. The minimum expected count is .01.

## PAST 30 DAYS USE:TOBACCO \* Gender

**Crosstab**

			Gender			Total
			Male	Female	Unknown	
	0 DAYS	Count	129	324	1	454
		Expected Count	144.3	308.5	1.2	454.0
		% within PAST 30 DAYS USE:TOBACCO	28.4%	71.4%	.2%	100.0%
		% within Gender	55.6%	65.3%	50.0%	62.2%
		% of Total	17.7%	44.4%	.1%	62.2%
	1-2 DAYS	Count	27	46	0	73
		Expected Count	23.2	49.6	.2	73.0
		% within PAST 30 DAYS USE:TOBACCO	37.0%	63.0%	.0%	100.0%
		% within Gender	11.6%	9.3%	.0%	10.0%
		% of Total	3.7%	6.3%	.0%	10.0%

PAST 30 DAYS USE:TOBACCO	3-5 DAYS	Count	10	15	0	25
		Expected Count	7.9	17.0	.1	25.0
		% within PAST 30 DAYS USE:TOBACCO	40.0%	60.0%	.0%	100.0%
		% within Gender	4.3%	3.0%	.0%	3.4%
		% of Total	1.4%	2.1%	.0%	3.4%
	6-9 DAYS	Count	7	14	0	21
		Expected Count	6.7	14.3	.1	21.0
		% within PAST 30 DAYS USE:TOBACCO	33.3%	66.7%	.0%	100.0%
		% within Gender	3.0%	2.8%	.0%	2.9%
		% of Total	1.0%	1.9%	.0%	2.9%
	10-19 DAYS	Count	13	25	0	38
		Expected Count	12.1	25.8	.1	38.0
		% within PAST 30 DAYS USE:TOBACCO	34.2%	65.8%	.0%	100.0%
		% within Gender	5.6%	5.0%	.0%	5.2%
		% of Total	1.8%	3.4%	.0%	5.2%
	20-29 DAYS	Count	15	26	0	41
		Expected Count	13.0	27.9	.1	41.0
		% within PAST 30 DAYS USE:TOBACCO	36.6%	63.4%	.0%	100.0%
		% within Gender	6.5%	5.2%	.0%	5.6%
		% of Total	2.1%	3.6%	.0%	5.6%
	ALL 30 DAYS	Count	31	46	1	78
		Expected Count	24.8	53.0	.2	78.0
		% within PAST 30 DAYS USE:TOBACCO	39.7%	59.0%	1.3%	100.0%
		% within Gender	13.4%	9.3%	50.0%	10.7%
		% of Total	4.2%	6.3%	.1%	10.7%
Total		Count	232	496	2	730
		Expected Count	232.0	496.0	2.0	730.0
		% within PAST 30 DAYS USE:TOBACCO	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	10.551(a)	12	.568
Likelihood Ratio	9.624	12	.649
Linear-by-Linear Association	4.229	1	.040
N of Valid Cases	730		

a 7 cells (33.3%) have expected count less than 5. The minimum expected count is .06.

## PAST 30 DAYS USE:ALCOHOL \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
PAST 30 DAYS USE:ALCOHOL	0 DAYS	Count	34	76	1	111
		Expected Count	35.4	75.3	.3	111.0
		% within PAST 30 DAYS USE:ALCOHOL	30.6%	68.5%	.9%	100.0%
		% within Gender	14.6%	15.4%	50.0%	15.2%
		% of Total	4.7%	10.4%	.1%	15.2%
	1-2 DAYS	Count	30	118	0	148
		Expected Count	47.2	100.4	.4	148.0
		% within PAST 30 DAYS USE:ALCOHOL	20.3%	79.7%	.0%	100.0%
		% within Gender	12.9%	23.8%	.0%	20.3%
		% of Total	4.1%	16.2%	.0%	20.3%
	3-5 DAYS	Count	42	103	0	145
		Expected Count	46.3	98.3	.4	145.0
		% within PAST 30 DAYS USE:ALCOHOL	29.0%	71.0%	.0%	100.0%
		% within Gender	18.0%	20.8%	.0%	19.9%
		% of Total	5.8%	14.1%	.0%	19.9%
	6-9 DAYS	Count	61	107	0	168
		Expected Count	53.6	113.9	.5	168.0
		% within PAST 30 DAYS USE:ALCOHOL	36.3%	63.7%	.0%	100.0%
		% within Gender	26.2%	21.6%	.0%	23.0%
		% of Total	8.4%	14.7%	.0%	23.0%
	10-19 DAYS	Count	55	86	0	141
		Expected Count	45.0	95.6	.4	141.0
		% within PAST 30 DAYS USE:ALCOHOL	39.0%	61.0%	.0%	100.0%
		% within Gender	23.6%	17.4%	.0%	19.3%
		% of Total	7.5%	11.8%	.0%	19.3%
	20-29 DAYS	Count	10	5	1	16
		Expected Count	5.1	10.8	.0	16.0
		% within PAST 30 DAYS USE:ALCOHOL	62.5%	31.3%	6.3%	100.0%

		% within Gender	4.3%	1.0%	50.0%	2.2%
		% of Total	1.4%	.7%	.1%	2.2%
ALL 30 DAYS	Count	1	0	0	1	
	Expected Count	.3	.7	.0	1.0	
	% within PAST 30 DAYS USE:ALCOHOL	100.0%	.0%	.0%	100.0%	
	% within Gender	.4%	.0%	.0%	.1%	
	% of Total	.1%	.0%	.0%	.1%	
Total	Count	233	495	2	730	
	Expected Count	233.0	495.0	2.0	730.0	
	% within PAST 30 DAYS USE:ALCOHOL	31.9%	67.8%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	31.9%	67.8%	.3%	100.0%	

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	48.770(a)	12	.000
Likelihood Ratio	33.856	12	.001
Linear-by-Linear Association	12.722	1	.000
N of Valid Cases	730		

a 9 cells (42.9%) have expected count less than 5. The minimum expected count is .00.

## PAST 30 DAYS USE:MARIJUANA \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
0 DAYS	Count	185	396	2	583	
	Expected Count	185.3	396.1	1.6	583.0	
	% within PAST 30 DAYS USE:MARIJUANA	31.7%	67.9%	.3%	100.0%	
	% within Gender	79.7%	79.8%	100.0%	79.9%	
	% of Total	25.3%	54.2%	.3%	79.9%	
1-2 DAYS	Count	21	43	0	64	
	Expected Count	20.3	43.5	.2	64.0	
	% within PAST 30 DAYS USE:MARIJUANA	32.8%	67.2%	.0%	100.0%	
	% within Gender	9.1%	8.7%	.0%	8.8%	
	% of Total	2.9%	5.9%	.0%	8.8%	
	Count	6	15	0	21	

PAST 30 DAYS USE:MARIJUANA	3-5 DAYS	Expected Count	6.7	14.3	.1	21.0
		% within PAST 30 DAYS USE:MARIJUANA	28.6%	71.4%	.0%	100.0%
		% within Gender	2.6%	3.0%	.0%	2.9%
		% of Total	.8%	2.1%	.0%	2.9%
	6-9 DAYS	Count	3	12	0	15
		Expected Count	4.8	10.2	.0	15.0
		% within PAST 30 DAYS USE:MARIJUANA	20.0%	80.0%	.0%	100.0%
		% within Gender	1.3%	2.4%	.0%	2.1%
		% of Total	.4%	1.6%	.0%	2.1%
	10-19 DAYS	Count	4	15	0	19
		Expected Count	6.0	12.9	.1	19.0
		% within PAST 30 DAYS USE:MARIJUANA	21.1%	78.9%	.0%	100.0%
		% within Gender	1.7%	3.0%	.0%	2.6%
		% of Total	.5%	2.1%	.0%	2.6%
	20-29 DAYS	Count	7	9	0	16
		Expected Count	5.1	10.9	.0	16.0
		% within PAST 30 DAYS USE:MARIJUANA	43.8%	56.3%	.0%	100.0%
		% within Gender	3.0%	1.8%	.0%	2.2%
		% of Total	1.0%	1.2%	.0%	2.2%
	ALL 30 DAYS	Count	6	6	0	12
		Expected Count	3.8	8.2	.0	12.0
		% within PAST 30 DAYS USE:MARIJUANA	50.0%	50.0%	.0%	100.0%
		% within Gender	2.6%	1.2%	.0%	1.6%
		% of Total	.8%	.8%	.0%	1.6%
Total	Count	232	496	2	730	
	Expected Count	232.0	496.0	2.0	730.0	
	% within PAST 30 DAYS USE:MARIJUANA	31.8%	67.9%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	31.8%	67.9%	.3%	100.0%	

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	5.505(a)	12	.939
Likelihood Ratio	5.875	12	.922
Linear-by-Linear Association	.340	1	.560
N of Valid Cases	730		

a 9 cells (42.9%) have expected count less than 5. The minimum expected count is .03.

# PAST 30 DAYS USE:COCAINE \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
PAST 30 DAYS USE:COCAINE	0 DAYS	Count	226	486	2	714
		Expected Count	226.9	485.2	2.0	714.0
		% within PAST 30 DAYS USE:COCAINE	31.7%	68.1%	.3%	100.0%
		% within Gender	97.8%	98.4%	100.0%	98.2%
		% of Total	31.1%	66.9%	.3%	98.2%
	1-2 DAYS	Count	5	6	0	11
		Expected Count	3.5	7.5	.0	11.0
		% within PAST 30 DAYS USE:COCAINE	45.5%	54.5%	.0%	100.0%
		% within Gender	2.2%	1.2%	.0%	1.5%
		% of Total	.7%	.8%	.0%	1.5%
	3-5 DAYS	Count	0	2	0	2
		Expected Count	.6	1.4	.0	2.0
		% within PAST 30 DAYS USE:COCAINE	.0%	100.0%	.0%	100.0%
		% within Gender	.0%	.4%	.0%	.3%
		% of Total	.0%	.3%	.0%	.3%
Total	Count	231	494	2	727	
	Expected Count	231.0	494.0	2.0	727.0	
	% within PAST 30 DAYS USE:COCAINE	31.8%	68.0%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	31.8%	68.0%	.3%	100.0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1.918(a)	4	.751
Likelihood Ratio	2.495	4	.646
Linear-by-Linear Association	.018	1	.892
N of Valid Cases	727		

a 6 cells (66.7%) have expected count less than 5. The minimum expected count is .01.

# PAST 30 DAYS USE:AMPHETAMINES \*

# Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
PAST 30 DAYS USE:AMPHETAMINES	0 DAYS	Count	229	478	2	709
		Expected Count	226.3	480.8	1.9	709.0
		% within PAST 30 DAYS USE:AMPHETAMINES	32.3%	67.4%	.3%	100.0%
		% within Gender	98.3%	96.6%	100.0%	97.1%
		% of Total	31.4%	65.5%	.3%	97.1%
	1-2 DAYS	Count	2	4	0	6
		Expected Count	1.9	4.1	.0	6.0
		% within PAST 30 DAYS USE:AMPHETAMINES	33.3%	66.7%	.0%	100.0%
		% within Gender	.9%	.8%	.0%	.8%
		% of Total	.3%	.5%	.0%	.8%
	3-5 DAYS	Count	1	1	0	2
		Expected Count	.6	1.4	.0	2.0
		% within PAST 30 DAYS USE:AMPHETAMINES	50.0%	50.0%	.0%	100.0%
		% within Gender	.4%	.2%	.0%	.3%
		% of Total	.1%	.1%	.0%	.3%
	6-9 DAYS	Count	0	7	0	7
		Expected Count	2.2	4.7	.0	7.0
		% within PAST 30 DAYS USE:AMPHETAMINES	.0%	100.0%	.0%	100.0%
		% within Gender	.0%	1.4%	.0%	1.0%
		% of Total	.0%	1.0%	.0%	1.0%
	10-19 DAYS	Count	0	2	0	2
		Expected Count	.6	1.4	.0	2.0
		% within PAST 30 DAYS USE:AMPHETAMINES	.0%	100.0%	.0%	100.0%
		% within Gender	.0%	.4%	.0%	.3%
		% of Total	.0%	.3%	.0%	.3%
	20-29 DAYS	Count	0	1	0	1
		Expected Count	.3	.7	.0	1.0
		% within PAST 30 DAYS USE:AMPHETAMINES	.0%	100.0%	.0%	100.0%
		% within Gender	.0%	.2%	.0%	.1%
		% of Total	.0%	.1%	.0%	.1%

		Count	1	2	0	3
		Expected Count	1.0	2.0	.0	3.0
	ALL 30 DAYS	% within PAST 30 DAYS USE:AMPHETAMINES	33.3%	66.7%	.0%	100.0%
		% within Gender	.4%	.4%	.0%	.4%
		% of Total	.1%	.3%	.0%	.4%
Total		Count	233	495	2	730
		Expected Count	233.0	495.0	2.0	730.0
		% within PAST 30 DAYS USE:AMPHETAMINES	31.9%	67.8%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.9%	67.8%	.3%	100.0%

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	5.133(a)	12	.953
Likelihood Ratio	8.164	12	.772
Linear-by-Linear Association	1.760	1	.185
N of Valid Cases	730		

a 19 cells (90.5%) have expected count less than 5. The minimum expected count is .00.

## PAST 30 DAYS USE:SEDATIVES \* Gender

### Crosstab

			Gender			Total
			Male	Female	Unknown	
PAST 30 DAYS USE:SEDATIVES	0 DAYS	Count	229	492	2	723
		Expected Count	229.8	491.2	2.0	723.0
		% within PAST 30 DAYS USE:SEDATIVES	31.7%	68.0%	.3%	100.0%
		% within Gender	98.7%	99.2%	100.0%	99.0%
		% of Total	31.4%	67.4%	.3%	99.0%
	1-2 DAYS	Count	3	2	0	5
		Expected Count	1.6	3.4	.0	5.0
		% within PAST 30 DAYS USE:SEDATIVES	60.0%	40.0%	.0%	100.0%
		% within Gender	1.3%	.4%	.0%	.7%
		% of Total	.4%	.3%	.0%	.7%
		Count	0	1	0	1
		Expected Count	.3	.7	.0	1.0
		% within PAST 30 DAYS				

	3-5 DAYS	USE:SEDATIVES	.0%	100.0%	.0%	100.0%	
		% within Gender	.0%	.2%	.0%	.1%	
		% of Total	.0%	.1%	.0%	.1%	
	10-19 DAYS	Count	0	1	0	1	
		Expected Count	.3	.7	.0	1.0	
		% within PAST 30 DAYS USE:SEDATIVES	.0%	100.0%	.0%	100.0%	
		% within Gender	.0%	.2%	.0%	.1%	
		% of Total	.0%	.1%	.0%	.1%	
	Total		Count	232	496	2	730
			Expected Count	232.0	496.0	2.0	730.0
			% within PAST 30 DAYS USE:SEDATIVES	31.8%	67.9%	.3%	100.0%
			% within Gender	100.0%	100.0%	100.0%	100.0%
			% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	2.789(a)	6	.835
Likelihood Ratio	3.243	6	.778
Linear-by-Linear Association	.039	1	.843
N of Valid Cases	730		

a 10 cells (83.3%) have expected count less than 5. The minimum expected count is .00.

## PAST 30 DAYS USE:HALLUCCINOGENS \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
PAST 30 DAYS USE:HALLUCCINOGENS	0 DAYS	Count	224	491	2	717
		Expected Count	226.5	488.5	2.0	717.0
		% within PAST 30 DAYS USE:HALLUCCINOGENS	31.2%	68.5%	.3%	100.0%
		% within Gender	97.4%	99.0%	100.0%	98.5%
		% of Total	30.8%	67.4%	.3%	98.5%
	1-2 DAYS	Count	6	5	0	11
		Expected Count	3.5	7.5	.0	11.0
		% within PAST 30 DAYS USE:HALLUCCINOGENS	54.5%	45.5%	.0%	100.0%

	% within Gender	2.6%	1.0%	.0%	1.5%
	% of Total	.8%	.7%	.0%	1.5%
Total	Count	230	496	2	728
	Expected Count	230.0	496.0	2.0	728.0
	% within PAST 30 DAYS USE:HALLUCINOGENS	31.6%	68.1%	.3%	100.0%
	% within Gender	100.0%	100.0%	100.0%	100.0%
	% of Total	31.6%	68.1%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	2.736(a)	2	.255
Likelihood Ratio	2.547	2	.280
Linear-by-Linear Association	2.728	1	.099
N of Valid Cases	728		

a 3 cells (50.0%) have expected count less than 5. The minimum expected count is .03.

## PAST 30 DAYS USE:OPIATES \* Gender

## Crosstab

		Gender			Total	
		Male	Female	Unknown		
PAST 30 DAYS USE:OPIATES	0 DAYS	Count	231	493	2	726
		Expected Count	232.4	491.6	2.0	726.0
		% within PAST 30 DAYS USE:OPIATES	31.8%	67.9%	.3%	100.0%
		% within Gender	99.1%	100.0%	100.0%	99.7%
		% of Total	31.7%	67.7%	.3%	99.7%
	1-2 DAYS	Count	2	0	0	2
		Expected Count	.6	1.4	.0	2.0
		% within PAST 30 DAYS USE:OPIATES	100.0%	.0%	.0%	100.0%
		% within Gender	.9%	.0%	.0%	.3%
		% of Total	.3%	.0%	.0%	.3%
Total	Count	233	493	2	728	
	Expected Count	233.0	493.0	2.0	728.0	
	% within PAST 30 DAYS USE:OPIATES	32.0%	67.7%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	32.0%	67.7%	.3%	100.0%	

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	4.261(a)	2	.119
<b>Likelihood Ratio</b>	4.569	2	.102
<b>Linear-by-Linear Association</b>	4.202	1	.040
<b>N of Valid Cases</b>	728		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .01.

**PAST 30 DAYS USE:INHALANTS \* Gender****Crosstab**

			Gender			Total
			Male	Female	Unknown	
<b>PAST 30 DAYS USE:INHALANTS</b>	<b>0 DAYS</b>	Count	231	494	2	727
		Expected Count	231.4	493.6	2.0	727.0
		% within PAST 30 DAYS USE:INHALANTS	31.8%	68.0%	.3%	100.0%
		% within Gender	99.6%	99.8%	100.0%	99.7%
		% of Total	31.7%	67.8%	.3%	99.7%
	<b>1-2 DAYS</b>	Count	1	1	0	2
		Expected Count	.6	1.4	.0	2.0
		% within PAST 30 DAYS USE:INHALANTS	50.0%	50.0%	.0%	100.0%
		% within Gender	.4%	.2%	.0%	.3%
		% of Total	.1%	.1%	.0%	.3%
<b>Total</b>		Count	232	495	2	729
		Expected Count	232.0	495.0	2.0	729.0
		% within PAST 30 DAYS USE:INHALANTS	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	.308(a)	2	.857
<b>Likelihood Ratio</b>	.292	2	.864
<b>Linear-by-Linear Association</b>	.308	1	.579
<b>N of Valid Cases</b>	729		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .01.

# PAST 30 DAYS USE:DESIGNER \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
PAST 30 DAYS USE:DESIGNER	0 DAYS	Count	232	494	2	728
		Expected Count	231.0	495.0	2.0	728.0
		% within PAST 30 DAYS USE:DESIGNER	31.9%	67.9%	.3%	100.0%
		% within Gender	100.0%	99.4%	100.0%	99.6%
		% of Total	31.7%	67.6%	.3%	99.6%
	1-2 DAYS	Count	0	3	0	3
		Expected Count	1.0	2.0	.0	3.0
		% within PAST 30 DAYS USE:DESIGNER	.0%	100.0%	.0%	100.0%
		% within Gender	.0%	.6%	.0%	.4%
		% of Total	.0%	.4%	.0%	.4%
Total	Count	232	497	2	731	
	Expected Count	232.0	497.0	2.0	731.0	
	% within PAST 30 DAYS USE:DESIGNER	31.7%	68.0%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	31.7%	68.0%	.3%	100.0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1.418(a)	2	.492
Likelihood Ratio	2.321	2	.313
Linear-by-Linear Association	1.347	1	.246
N of Valid Cases	731		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .01.

# PAST 30 DAYS USE:STEROIDS \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
		Count	231	495	2	728
		Expected Count	231.0	495.0	2.0	728.0

PAST 30 DAYS USE:STEROIDS	0 DAYS	% within PAST 30 DAYS USE:STEROIDS	31.7%	68.0%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.7%	68.0%	.3%	100.0%
Total	Count	231	495	2	728	
	Expected Count	231.0	495.0	2.0	728.0	
	% within PAST 30 DAYS USE:STEROIDS	31.7%	68.0%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	31.7%	68.0%	.3%	100.0%	

## Chi-Square Tests

	Value
Pearson Chi-Square	.(a)
N of Valid Cases	728

a No statistics are computed because PAST 30 DAYS USE:STEROIDS is a constant.

## PAST 30 DAYS USE:OTHER \* Gender

## Crosstab

			Gender			Total	
			Male	Female	Unknown		
PAST 30 DAYS USE:OTHER	0 DAYS	Count	232	492	2	726	
		Expected Count	231.7	492.3	2.0	726.0	
		% within PAST 30 DAYS USE:OTHER	32.0%	67.8%	.3%	100.0%	
		% within Gender	99.6%	99.4%	100.0%	99.5%	
		% of Total	31.8%	67.4%	.3%	99.5%	
	1-2 DAYS	Count	1	2	0	3	
		Expected Count	1.0	2.0	.0	3.0	
		% within PAST 30 DAYS USE:OTHER	33.3%	66.7%	.0%	100.0%	
		% within Gender	.4%	.4%	.0%	.4%	
		% of Total	.1%	.3%	.0%	.4%	
	3-5 DAYS	Count	0	1	0	1	
		Expected Count	.3	.7	.0	1.0	
		% within PAST 30 DAYS USE:OTHER	.0%	100.0%	.0%	100.0%	
		% within Gender	.0%	.2%	.0%	.1%	
		% of Total	.0%	.1%	.0%	.1%	
			Count	233	495	2	
						730	

Total	Expected Count	233.0	495.0	2.0	730.0
	% within PAST 30 DAYS USE:OTHER	31.9%	67.8%	.3%	100.0%
	% within Gender	100.0%	100.0%	100.0%	100.0%
	% of Total	31.9%	67.8%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.486(a)	4	.975
Likelihood Ratio	.796	4	.939
Linear-by-Linear Association	.219	1	.640
N of Valid Cases	730		

a 7 cells (77.8%) have expected count less than 5. The minimum expected count is .00.

## AVERAGE USE:TOBACCO \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
	Never used	Count	5	7	0	12
		Expected Count	3.8	8.1	.0	12.0
		% within AVERAGE USE:TOBACCO	41.7%	58.3%	.0%	100.0%
		% within Gender	2.1%	1.4%	.0%	1.6%
		% of Total	.7%	1.0%	.0%	1.6%
	Once/year	Count	4	4	0	8
		Expected Count	2.6	5.4	.0	8.0
		% within AVERAGE USE:TOBACCO	50.0%	50.0%	.0%	100.0%
		% within Gender	1.7%	.8%	.0%	1.1%
		% of Total	.5%	.5%	.0%	1.1%
	6 times/year	Count	3	8	1	12
		Expected Count	3.8	8.1	.0	12.0
		% within AVERAGE USE:TOBACCO	25.0%	66.7%	8.3%	100.0%
		% within Gender	1.3%	1.6%	50.0%	1.6%
		% of Total	.4%	1.1%	.1%	1.6%
	Once/month	Count	8	11	0	19
		Expected Count	6.1	12.9	.1	19.0
		% within AVERAGE USE:TOBACCO	42.1%	57.9%	.0%	100.0%

AVERAGE USE:TOBACCO	Twice/month	% within Gender	3.4%	2.2%	.0%	2.6%
		% of Total	1.1%	1.5%	.0%	2.6%
		Count	8	18	0	26
		Expected Count	8.3	17.6	.1	26.0
		% within AVERAGE USE:TOBACCO	30.8%	69.2%	.0%	100.0%
	Once/week	% within Gender	3.4%	3.6%	.0%	3.5%
		% of Total	1.1%	2.5%	.0%	3.5%
		Count	39	68	0	107
		Expected Count	34.2	72.5	.3	107.0
		% within AVERAGE USE:TOBACCO	36.4%	63.6%	.0%	100.0%
	3 times/week	% within Gender	16.7%	13.7%	.0%	14.6%
		% of Total	5.3%	9.3%	.0%	14.6%
		Count	66	131	1	198
		Expected Count	63.2	134.3	.5	198.0
		% within AVERAGE USE:TOBACCO	33.3%	66.2%	.5%	100.0%
	5 times/week	% within Gender	28.2%	26.4%	50.0%	27.0%
		% of Total	9.0%	17.9%	.1%	27.0%
		Count	54	86	0	140
		Expected Count	44.7	94.9	.4	140.0
		% within AVERAGE USE:TOBACCO	38.6%	61.4%	.0%	100.0%
	Every day	% within Gender	23.1%	17.3%	.0%	19.1%
		% of Total	7.4%	11.7%	.0%	19.1%
		Count	47	164	0	211
		Expected Count	67.4	143.1	.6	211.0
		% within AVERAGE USE:TOBACCO	22.3%	77.7%	.0%	100.0%
	Total	% within Gender	20.1%	33.0%	.0%	28.8%
		% of Total	6.4%	22.4%	.0%	28.8%
		Count	234	497	2	733
		Expected Count	234.0	497.0	2.0	733.0
		% within AVERAGE USE:TOBACCO	31.9%	67.8%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.9%	67.8%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	46.362(a)	16	.000
Likelihood Ratio	24.380	16	.082

Linear-by-Linear Association	4.815	1	.028
N of Valid Cases	733		
a 12 cells (44.4%) have expected count less than 5. The minimum expected count is .02.			

## AVERAGE USE:ALCOHOL \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
AVERAGE USE:ALCOHOL	Never used	Count	1	3	0	4
		Expected Count	1.3	2.7	.0	4.0
		% within AVERAGE USE:ALCOHOL	25.0%	75.0%	.0%	100.0%
		% within Gender	.4%	.6%	.0%	.5%
		% of Total	.1%	.4%	.0%	.5%
	Once/year	Count	2	0	0	2
		Expected Count	.6	1.4	.0	2.0
		% within AVERAGE USE:ALCOHOL	100.0%	.0%	.0%	100.0%
		% within Gender	.9%	.0%	.0%	.3%
		% of Total	.3%	.0%	.0%	.3%
	6 times/year	Count	0	3	0	3
		Expected Count	1.0	2.0	.0	3.0
		% within AVERAGE USE:ALCOHOL	.0%	100.0%	.0%	100.0%
		% within Gender	.0%	.6%	.0%	.4%
		% of Total	.0%	.4%	.0%	.4%
	Once/month	Count	1	4	0	5
		Expected Count	1.6	3.4	.0	5.0
		% within AVERAGE USE:ALCOHOL	20.0%	80.0%	.0%	100.0%
		% within Gender	.4%	.8%	.0%	.7%
		% of Total	.1%	.5%	.0%	.7%
	Twice/month	Count	9	10	0	19
		Expected Count	6.1	12.9	.1	19.0
		% within AVERAGE USE:ALCOHOL	47.4%	52.6%	.0%	100.0%
		% within Gender	3.8%	2.0%	.0%	2.6%
		% of Total	1.2%	1.4%	.0%	2.6%
		Count	75	126	1	202
		Expected Count	64.7	136.8	.6	202.0

	Once/week	% within AVERAGE USE:ALCOHOL	37.1%	62.4%	.5%	100.0%
		% within Gender	31.9%	25.4%	50.0%	27.5%
		% of Total	10.2%	17.2%	.1%	27.5%
	3 times/week	Count	118	274	1	393
		Expected Count	125.8	266.1	1.1	393.0
		% within AVERAGE USE:ALCOHOL	30.0%	69.7%	.3%	100.0%
		% within Gender	50.2%	55.1%	50.0%	53.5%
		% of Total	16.1%	37.3%	.1%	53.5%
	5 times/week	Count	21	58	0	79
		Expected Count	25.3	53.5	.2	79.0
		% within AVERAGE USE:ALCOHOL	26.6%	73.4%	.0%	100.0%
		% within Gender	8.9%	11.7%	.0%	10.8%
		% of Total	2.9%	7.9%	.0%	10.8%
	Every day	Count	8	19	0	27
		Expected Count	8.6	18.3	.1	27.0
		% within AVERAGE USE:ALCOHOL	29.6%	70.4%	.0%	100.0%
		% within Gender	3.4%	3.8%	.0%	3.7%
		% of Total	1.1%	2.6%	.0%	3.7%
Total	Count	235	497	2	734	
	Expected Count	235.0	497.0	2.0	734.0	
	% within AVERAGE USE:ALCOHOL	32.0%	67.7%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	32.0%	67.7%	.3%	100.0%	

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	13.285(a)	16	.652
Likelihood Ratio	14.686	16	.548
Linear-by-Linear Association	2.692	1	.101
N of Valid Cases	734		

a 17 cells (63.0%) have expected count less than 5. The minimum expected count is .01.

## AVERAGE USE:MARIJUANA \* Gender

### Crosstab

	Gender	Total

			Male	Female	Unknown	
AVERAGE USE:MARIJUANA	Never used	Count	6	9	2	17
		Expected Count	5.4	11.5	.0	17.0
		% within AVERAGE USE:MARIJUANA	35.3%	52.9%	11.8%	100.0%
		% within Gender	2.6%	1.8%	100.0%	2.3%
		% of Total	.8%	1.2%	.3%	2.3%
	Once/year	Count	18	37	0	55
		Expected Count	17.6	37.3	.1	55.0
		% within AVERAGE USE:MARIJUANA	32.7%	67.3%	.0%	100.0%
		% within Gender	7.7%	7.4%	.0%	7.5%
		% of Total	2.4%	5.0%	.0%	7.5%
	6 times/year	Count	34	43	0	77
		Expected Count	24.6	52.2	.2	77.0
		% within AVERAGE USE:MARIJUANA	44.2%	55.8%	.0%	100.0%
		% within Gender	14.5%	8.6%	.0%	10.5%
		% of Total	4.6%	5.9%	.0%	10.5%
	Once/month	Count	27	66	0	93
		Expected Count	29.7	63.0	.3	93.0
		% within AVERAGE USE:MARIJUANA	29.0%	71.0%	.0%	100.0%
		% within Gender	11.5%	13.3%	.0%	12.7%
		% of Total	3.7%	9.0%	.0%	12.7%
	Twice/month	Count	35	90	0	125
		Expected Count	40.0	84.7	.3	125.0
		% within AVERAGE USE:MARIJUANA	28.0%	72.0%	.0%	100.0%
		% within Gender	14.9%	18.1%	.0%	17.0%
		% of Total	4.8%	12.2%	.0%	17.0%
	Once/week	Count	66	121	0	187
		Expected Count	59.8	126.7	.5	187.0
		% within AVERAGE USE:MARIJUANA	35.3%	64.7%	.0%	100.0%
		% within Gender	28.1%	24.3%	.0%	25.4%
		% of Total	9.0%	16.5%	.0%	25.4%
	3 times/week	Count	35	90	0	125
		Expected Count	40.0	84.7	.3	125.0
		% within AVERAGE USE:MARIJUANA	28.0%	72.0%	.0%	100.0%
		% within Gender	14.9%	18.1%	.0%	17.0%
		% of Total	4.8%	12.2%	.0%	17.0%

		Count	8	26	0	34
		Expected Count	10.9	23.0	.1	34.0
	5 times/week	% within AVERAGE USE:MARIJUANA	23.5%	76.5%	.0%	100.0%
		% within Gender	3.4%	5.2%	.0%	4.6%
		% of Total	1.1%	3.5%	.0%	4.6%
		Count	6	16	0	22
		Expected Count	7.0	14.9	.1	22.0
	Every day	% within AVERAGE USE:MARIJUANA	27.3%	72.7%	.0%	100.0%
		% within Gender	2.6%	3.2%	.0%	3.0%
		% of Total	.8%	2.2%	.0%	3.0%
Total		Count	235	498	2	735
		Expected Count	235.0	498.0	2.0	735.0
		% within AVERAGE USE:MARIJUANA	32.0%	67.8%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	32.0%	67.8%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	94.844(a)	16	.000
Likelihood Ratio	25.301	16	.065
Linear-by-Linear Association	1.380	1	.240
N of Valid Cases	735		

a 9 cells (33.3%) have expected count less than 5. The minimum expected count is .05.

## AVERAGE USE:COCAINE \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
Never used		Count	84	182	2	268
		Expected Count	85.9	181.3	.7	268.0
		% within AVERAGE USE:COCAINE	31.3%	67.9%	.7%	100.0%
		% within Gender	35.7%	36.7%	100.0%	36.6%
		% of Total	11.5%	24.8%	.3%	36.6%
		Count	57	120	0	177
		Expected Count	56.7	119.8	.5	177.0
		% within AVERAGE				

AVERAGE USE:COCAINE	Once/year	USE:COCAINE	32.2%	67.8%	.0%	100.0%
		% within Gender	24.3%	24.2%	.0%	24.1%
		% of Total	7.8%	16.4%	.0%	24.1%
	6 times/year	Count	26	53	0	79
		Expected Count	25.3	53.5	.2	79.0
		% within AVERAGE USE:COCAINE	32.9%	67.1%	.0%	100.0%
		% within Gender	11.1%	10.7%	.0%	10.8%
		% of Total	3.5%	7.2%	.0%	10.8%
	Once/month	Count	17	66	0	83
		Expected Count	26.6	56.2	.2	83.0
		% within AVERAGE USE:COCAINE	20.5%	79.5%	.0%	100.0%
		% within Gender	7.2%	13.3%	.0%	11.3%
		% of Total	2.3%	9.0%	.0%	11.3%
	Twice/month	Count	19	31	0	50
		Expected Count	16.0	33.8	.1	50.0
		% within AVERAGE USE:COCAINE	38.0%	62.0%	.0%	100.0%
		% within Gender	8.1%	6.3%	.0%	6.8%
		% of Total	2.6%	4.2%	.0%	6.8%
	Once/week	Count	21	29	0	50
		Expected Count	16.0	33.8	.1	50.0
		% within AVERAGE USE:COCAINE	42.0%	58.0%	.0%	100.0%
		% within Gender	8.9%	5.8%	.0%	6.8%
		% of Total	2.9%	4.0%	.0%	6.8%
	3 times/week	Count	6	12	0	18
		Expected Count	5.8	12.2	.0	18.0
		% within AVERAGE USE:COCAINE	33.3%	66.7%	.0%	100.0%
		% within Gender	2.6%	2.4%	.0%	2.5%
		% of Total	.8%	1.6%	.0%	2.5%
	5 times/week	Count	2	2	0	4
		Expected Count	1.3	2.7	.0	4.0
		% within AVERAGE USE:COCAINE	50.0%	50.0%	.0%	100.0%
		% within Gender	.9%	.4%	.0%	.5%
		% of Total	.3%	.3%	.0%	.5%
	Every day	Count	3	1	0	4
		Expected Count	1.3	2.7	.0	4.0
		% within AVERAGE USE:COCAINE	75.0%	25.0%	.0%	100.0%

		% within Gender	1.3%	.2%	.0%	.5%
		% of Total	.4%	.1%	.0%	.5%
Total	Count	235	496	2	733	
	Expected Count	235.0	496.0	2.0	733.0	
	% within AVERAGE USE:COCAINE	32.1%	67.7%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	32.1%	67.7%	.3%	100.0%	

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	15.725(a)	16	.472
Likelihood Ratio	16.244	16	.436
Linear-by-Linear Association	2.013	1	.156
N of Valid Cases	733		

a 13 cells (48.1%) have expected count less than 5. The minimum expected count is .01.

## AVERAGE USE:AMPHETAMINES \* Gender

### Crosstab

			Gender			Total
			Male	Female	Unknown	
	Never used	Count	80	114	1	195
		Expected Count	62.1	132.4	.5	195.0
		% within AVERAGE USE:AMPHETAMINES	41.0%	58.5%	.5%	100.0%
		% within Gender	34.5%	23.0%	50.0%	26.7%
		% of Total	11.0%	15.6%	.1%	26.7%
	Once/year	Count	44	79	1	124
		Expected Count	39.5	84.2	.3	124.0
		% within AVERAGE USE:AMPHETAMINES	35.5%	63.7%	.8%	100.0%
		% within Gender	19.0%	16.0%	50.0%	17.0%
		% of Total	6.0%	10.8%	.1%	17.0%
	6 times/year	Count	23	76	0	99
		Expected Count	31.5	67.2	.3	99.0
		% within AVERAGE USE:AMPHETAMINES	23.2%	76.8%	.0%	100.0%
		% within Gender	9.9%	15.4%	.0%	13.6%
		% of Total	3.2%	10.4%	.0%	13.6%
	Count	23	52	0	75	

AVERAGE USE:AMPHETAMINES	Once/month	Expected Count	23.9	50.9	.2	75.0
		% within AVERAGE USE:AMPHETAMINES	30.7%	69.3%	.0%	100.0%
		% within Gender	9.9%	10.5%	.0%	10.3%
		% of Total	3.2%	7.1%	.0%	10.3%
	Twice/month	Count	19	47	0	66
		Expected Count	21.0	44.8	.2	66.0
		% within AVERAGE USE:AMPHETAMINES	28.8%	71.2%	.0%	100.0%
		% within Gender	8.2%	9.5%	.0%	9.1%
		% of Total	2.6%	6.4%	.0%	9.1%
	Once/week	Count	22	53	0	75
		Expected Count	23.9	50.9	.2	75.0
		% within AVERAGE USE:AMPHETAMINES	29.3%	70.7%	.0%	100.0%
		% within Gender	9.5%	10.7%	.0%	10.3%
		% of Total	3.0%	7.3%	.0%	10.3%
	3 times/week	Count	11	39	0	50
		Expected Count	15.9	34.0	.1	50.0
		% within AVERAGE USE:AMPHETAMINES	22.0%	78.0%	.0%	100.0%
		% within Gender	4.7%	7.9%	.0%	6.9%
		% of Total	1.5%	5.3%	.0%	6.9%
	5 times/week	Count	5	16	0	21
		Expected Count	6.7	14.3	.1	21.0
		% within AVERAGE USE:AMPHETAMINES	23.8%	76.2%	.0%	100.0%
		% within Gender	2.2%	3.2%	.0%	2.9%
		% of Total	.7%	2.2%	.0%	2.9%
	Every day	Count	5	19	0	24
		Expected Count	7.6	16.3	.1	24.0
		% within AVERAGE USE:AMPHETAMINES	20.8%	79.2%	.0%	100.0%
		% within Gender	2.2%	3.8%	.0%	3.3%
		% of Total	.7%	2.6%	.0%	3.3%
	Total	Count	232	495	2	729
		Expected Count	232.0	495.0	2.0	729.0
		% within AVERAGE USE:AMPHETAMINES	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	19.687(a)	16	.235
<b>Likelihood Ratio</b>	20.453	16	.201
<b>Linear-by-Linear Association</b>	9.477	1	.002
<b>N of Valid Cases</b>	729		

a 9 cells (33.3%) have expected count less than 5. The minimum expected count is .06.

## AVERAGE USE:SEDATIVES \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
AVERAGE USE:SEDATIVES	Never used	Count	95	179	2	276
		Expected Count	88.3	186.9	.8	276.0
		% within AVERAGE USE:SEDATIVES	34.4%	64.9%	.7%	100.0%
		% within Gender	40.8%	36.3%	100.0%	37.9%
		% of Total	13.0%	24.6%	.3%	37.9%
	Once/year	Count	56	106	0	162
		Expected Count	51.8	109.7	.4	162.0
		% within AVERAGE USE:SEDATIVES	34.6%	65.4%	.0%	100.0%
		% within Gender	24.0%	21.5%	.0%	22.3%
		% of Total	7.7%	14.6%	.0%	22.3%
	6 times/year	Count	19	66	0	85
		Expected Count	27.2	57.6	.2	85.0
		% within AVERAGE USE:SEDATIVES	22.4%	77.6%	.0%	100.0%
		% within Gender	8.2%	13.4%	.0%	11.7%
		% of Total	2.6%	9.1%	.0%	11.7%
	Once/month	Count	24	58	0	82
		Expected Count	26.2	55.5	.2	82.0
		% within AVERAGE USE:SEDATIVES	29.3%	70.7%	.0%	100.0%
		% within Gender	10.3%	11.8%	.0%	11.3%
		% of Total	3.3%	8.0%	.0%	11.3%
	Twice/month	Count	13	34	0	47
		Expected Count	15.0	31.8	.1	47.0
		% within AVERAGE USE:SEDATIVES	27.7%	72.3%	.0%	100.0%
		% within Gender	5.6%	6.9%	.0%	6.5%

	% of Total	1.8%	4.7%	.0%	6.5%
Once/week	Count	16	28	0	44
	Expected Count	14.1	29.8	.1	44.0
	% within AVERAGE USE:SEDATIVES	36.4%	63.6%	.0%	100.0%
	% within Gender	6.9%	5.7%	.0%	6.0%
	% of Total	2.2%	3.8%	.0%	6.0%
3 times/week	Count	5	15	0	20
	Expected Count	6.4	13.5	.1	20.0
	% within AVERAGE USE:SEDATIVES	25.0%	75.0%	.0%	100.0%
	% within Gender	2.1%	3.0%	.0%	2.7%
	% of Total	.7%	2.1%	.0%	2.7%
5 times/week	Count	1	2	0	3
	Expected Count	1.0	2.0	.0	3.0
	% within AVERAGE USE:SEDATIVES	33.3%	66.7%	.0%	100.0%
	% within Gender	.4%	.4%	.0%	.4%
	% of Total	.1%	.3%	.0%	.4%
Every day	Count	4	5	0	9
	Expected Count	2.9	6.1	.0	9.0
	% within AVERAGE USE:SEDATIVES	44.4%	55.6%	.0%	100.0%
	% within Gender	1.7%	1.0%	.0%	1.2%
	% of Total	.5%	.7%	.0%	1.2%
Total	Count	233	493	2	728
	Expected Count	233.0	493.0	2.0	728.0
	% within AVERAGE USE:SEDATIVES	32.0%	67.7%	.3%	100.0%
	% within Gender	100.0%	100.0%	100.0%	100.0%
	% of Total	32.0%	67.7%	.3%	100.0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	10.476(a)	16	.841
Likelihood Ratio	11.295	16	.791
Linear-by-Linear Association	.318	1	.573
N of Valid Cases	728		

a 12 cells (44.4%) have expected count less than 5. The minimum expected count is .01.

## AVERAGE USE:HALLUCINOGENS \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
AVERAGE USE:HALLUINOGENS	Never used	Count	80	210	2	292
		Expected Count	92.8	198.4	.8	292.0
		% within AVERAGE USE:HALLUINOGENS	27.4%	71.9%	.7%	100.0%
		% within Gender	34.5%	42.3%	100.0%	40.0%
		% of Total	11.0%	28.8%	.3%	40.0%
	Once/year	Count	60	118	0	178
		Expected Count	56.6	120.9	.5	178.0
		% within AVERAGE USE:HALLUINOGENS	33.7%	66.3%	.0%	100.0%
		% within Gender	25.9%	23.8%	.0%	24.4%
		% of Total	8.2%	16.2%	.0%	24.4%
	6 times/year	Count	38	70	0	108
		Expected Count	34.3	73.4	.3	108.0
		% within AVERAGE USE:HALLUINOGENS	35.2%	64.8%	.0%	100.0%
		% within Gender	16.4%	14.1%	.0%	14.8%
		% of Total	5.2%	9.6%	.0%	14.8%
	Once/month	Count	22	38	0	60
		Expected Count	19.1	40.8	.2	60.0
		% within AVERAGE USE:HALLUINOGENS	36.7%	63.3%	.0%	100.0%
		% within Gender	9.5%	7.7%	.0%	8.2%
		% of Total	3.0%	5.2%	.0%	8.2%
	Twice/month	Count	7	21	0	28
		Expected Count	8.9	19.0	.1	28.0
		% within AVERAGE USE:HALLUINOGENS	25.0%	75.0%	.0%	100.0%
		% within Gender	3.0%	4.2%	.0%	3.8%
		% of Total	1.0%	2.9%	.0%	3.8%
	Once/week	Count	16	29	0	45
		Expected Count	14.3	30.6	.1	45.0
		% within AVERAGE USE:HALLUINOGENS	35.6%	64.4%	.0%	100.0%
		% within Gender	6.9%	5.8%	.0%	6.2%
		% of Total	2.2%	4.0%	.0%	6.2%
	3	Count	4	9	0	13
		Expected Count	4.1	8.8	.0	13.0
3 % within AVERAGE						

	times/week	USE:HALUCINOGENS	30.8%	69.2%	.0%	100.0%
		% within Gender	1.7%	1.8%	.0%	1.8%
		% of Total	.5%	1.2%	.0%	1.8%
	5 times/week	Count	1	0	0	1
		Expected Count	.3	.7	.0	1.0
		% within AVERAGE USE:HALUCINOGENS	100.0%	.0%	.0%	100.0%
		% within Gender	.4%	.0%	.0%	.1%
		% of Total	.1%	.0%	.0%	.1%
	Every day	Count	4	1	0	5
		Expected Count	1.6	3.4	.0	5.0
		% within AVERAGE USE:HALUCINOGENS	80.0%	20.0%	.0%	100.0%
		% within Gender	1.7%	.2%	.0%	.7%
		% of Total	.5%	.1%	.0%	.7%
Total	Count	232	496	2	730	
	Expected Count	232.0	496.0	2.0	730.0	
	% within AVERAGE USE:HALUCINOGENS	31.8%	67.9%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	31.8%	67.9%	.3%	100.0%	

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	15.305(a)	16	.502
Likelihood Ratio	15.739	16	.471
Linear-by-Linear Association	4.687	1	.030
N of Valid Cases	730		

a 14 cells (51.9%) have expected count less than 5. The minimum expected count is .00.

## AVERAGE USE:OPIATES \* Gender

#### Crosstab

			Gender			Total
			Male	Female	Unknown	
	Never used	Count	117	263	2	382
		Expected Count	122.4	258.5	1.0	382.0
		% within AVERAGE USE:OPIATES	30.6%	68.8%	.5%	100.0%
		% within Gender	50.0%	53.2%	100.0%	52.3%
		% of Total	16.0%	36.0%	.3%	52.3%

AVERAGE USE:OPIATES	Once/year	Count	63	113	0	176
		Expected Count	56.4	119.1	.5	176.0
		% within AVERAGE USE:OPIATES	35.8%	64.2%	.0%	100.0%
		% within Gender	26.9%	22.9%	.0%	24.1%
		% of Total	8.6%	15.5%	.0%	24.1%
	6 times/year	Count	12	48	0	60
		Expected Count	19.2	40.6	.2	60.0
		% within AVERAGE USE:OPIATES	20.0%	80.0%	.0%	100.0%
		% within Gender	5.1%	9.7%	.0%	8.2%
		% of Total	1.6%	6.6%	.0%	8.2%
	Once/month	Count	13	34	0	47
		Expected Count	15.1	31.8	.1	47.0
		% within AVERAGE USE:OPIATES	27.7%	72.3%	.0%	100.0%
		% within Gender	5.6%	6.9%	.0%	6.4%
		% of Total	1.8%	4.7%	.0%	6.4%
	Twice/month	Count	7	11	0	18
		Expected Count	5.8	12.2	.0	18.0
		% within AVERAGE USE:OPIATES	38.9%	61.1%	.0%	100.0%
		% within Gender	3.0%	2.2%	.0%	2.5%
		% of Total	1.0%	1.5%	.0%	2.5%
	Once/week	Count	14	16	0	30
		Expected Count	9.6	20.3	.1	30.0
		% within AVERAGE USE:OPIATES	46.7%	53.3%	.0%	100.0%
		% within Gender	6.0%	3.2%	.0%	4.1%
		% of Total	1.9%	2.2%	.0%	4.1%
	3 times/week	Count	2	7	0	9
		Expected Count	2.9	6.1	.0	9.0
		% within AVERAGE USE:OPIATES	22.2%	77.8%	.0%	100.0%
		% within Gender	.9%	1.4%	.0%	1.2%
		% of Total	.3%	1.0%	.0%	1.2%
	5 times/week	Count	2	0	0	2
		Expected Count	.6	1.4	.0	2.0
		% within AVERAGE USE:OPIATES	100.0%	.0%	.0%	100.0%
		% within Gender	.9%	.0%	.0%	.3%
		% of Total	.3%	.0%	.0%	.3%
		Count	4	2	0	6

		Expected Count	1.9	4.1	.0	6.0
	Every day	% within AVERAGE USE:OPIATES	66.7%	33.3%	.0%	100.0%
		% within Gender	1.7%	.4%	.0%	.8%
		% of Total	.5%	.3%	.0%	.8%
		Count	234	494	2	730
Total		Expected Count	234.0	494.0	2.0	730.0
		% within AVERAGE USE:OPIATES	32.1%	67.7%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	32.1%	67.7%	.3%	100.0%

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	18.941(a)	16	.272
Likelihood Ratio	19.898	16	.225
Linear-by-Linear Association	3.078	1	.079
N of Valid Cases	730		

a 14 cells (51.9%) have expected count less than 5. The minimum expected count is .01.

## AVERAGE USE:INHALANTS \* Gender

### Crosstab

			Gender			Total
			Male	Female	Unknown	
	Never used	Count	120	245	2	367
		Expected Count	118.3	247.7	1.0	367.0
		% within AVERAGE USE:INHALANTS	32.7%	66.8%	.5%	100.0%
		% within Gender	51.1%	49.8%	100.0%	50.3%
		% of Total	16.5%	33.6%	.3%	50.3%
	Once/year	Count	49	110	0	159
		Expected Count	51.3	107.3	.4	159.0
		% within AVERAGE USE:INHALANTS	30.8%	69.2%	.0%	100.0%
		% within Gender	20.9%	22.4%	.0%	21.8%
		% of Total	6.7%	15.1%	.0%	21.8%
	6 times/year	Count	27	50	0	77
		Expected Count	24.8	52.0	.2	77.0
		% within AVERAGE USE:INHALANTS	35.1%	64.9%	.0%	100.0%

AVERAGE USE:INHALANTS		% within Gender	11.5%	10.2%	.0%	10.6%
		% of Total	3.7%	6.9%	.0%	10.6%
		Count	17	34	0	51
		Expected Count	16.4	34.4	.1	51.0
		% within AVERAGE USE:INHALANTS	33.3%	66.7%	.0%	100.0%
		% within Gender	7.2%	6.9%	.0%	7.0%
		% of Total	2.3%	4.7%	.0%	7.0%
		Count	6	19	0	25
		Expected Count	8.1	16.9	.1	25.0
		% within AVERAGE USE:INHALANTS	24.0%	76.0%	.0%	100.0%
		% within Gender	2.6%	3.9%	.0%	3.4%
		% of Total	.8%	2.6%	.0%	3.4%
		Count	8	21	0	29
		Expected Count	9.3	19.6	.1	29.0
		% within AVERAGE USE:INHALANTS	27.6%	72.4%	.0%	100.0%
		% within Gender	3.4%	4.3%	.0%	4.0%
		% of Total	1.1%	2.9%	.0%	4.0%
		Count	3	10	0	13
		Expected Count	4.2	8.8	.0	13.0
		% within AVERAGE USE:INHALANTS	23.1%	76.9%	.0%	100.0%
		% within Gender	1.3%	2.0%	.0%	1.8%
		% of Total	.4%	1.4%	.0%	1.8%
		Count	2	1	0	3
		Expected Count	1.0	2.0	.0	3.0
		% within AVERAGE USE:INHALANTS	66.7%	33.3%	.0%	100.0%
		% within Gender	.9%	.2%	.0%	.4%
		% of Total	.3%	.1%	.0%	.4%
		Count	3	2	0	5
		Expected Count	1.6	3.4	.0	5.0
		% within AVERAGE USE:INHALANTS	60.0%	40.0%	.0%	100.0%
		% within Gender	1.3%	.4%	.0%	.7%
		% of Total	.4%	.3%	.0%	.7%
		Count	235	492	2	729
		Expected Count	235.0	492.0	2.0	729.0
		% within AVERAGE USE:INHALANTS	32.2%	67.5%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%

	% of Total	32.2%	67.5%	.3%	100.0%
--	------------	-------	-------	-----	--------

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	7.463(a)	16	.963
Likelihood Ratio	8.020	16	.948
Linear-by-Linear Association	.012	1	.912
N of Valid Cases	729		

a 14 cells (51.9%) have expected count less than 5. The minimum expected count is .01.

## AVERAGE USE:DESIGNER \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
	Never used	Count	61	136	2	199
		Expected Count	63.1	135.4	.5	199.0
		% within AVERAGE USE:DESIGNER	30.7%	68.3%	1.0%	100.0%
		% within Gender	26.3%	27.3%	100.0%	27.2%
		% of Total	8.3%	18.6%	.3%	27.2%
	Once/year	Count	52	124	0	176
		Expected Count	55.8	119.7	.5	176.0
		% within AVERAGE USE:DESIGNER	29.5%	70.5%	.0%	100.0%
		% within Gender	22.4%	24.9%	.0%	24.0%
		% of Total	7.1%	16.9%	.0%	24.0%
	6 times/year	Count	34	100	0	134
		Expected Count	42.5	91.2	.4	134.0
		% within AVERAGE USE:DESIGNER	25.4%	74.6%	.0%	100.0%
		% within Gender	14.7%	20.1%	.0%	18.3%
		% of Total	4.6%	13.7%	.0%	18.3%
	Once/month	Count	32	45	0	77
		Expected Count	24.4	52.4	.2	77.0
		% within AVERAGE USE:DESIGNER	41.6%	58.4%	.0%	100.0%
		% within Gender	13.8%	9.0%	.0%	10.5%
		% of Total	4.4%	6.1%	.0%	10.5%
		Count	21	43	0	64
		Expected Count	20.3	43.5	.2	64.0

AVERAGE USE:DESIGNER	Twice/month	% within AVERAGE USE:DESIGNER	32.8%	67.2%	.0%	100.0%
		% within Gender	9.1%	8.6%	.0%	8.7%
		% of Total	2.9%	5.9%	.0%	8.7%
	Once/week	Count	21	32	0	53
		Expected Count	16.8	36.1	.1	53.0
		% within AVERAGE USE:DESIGNER	39.6%	60.4%	.0%	100.0%
		% within Gender	9.1%	6.4%	.0%	7.2%
		% of Total	2.9%	4.4%	.0%	7.2%
	3 times/week	Count	5	14	0	19
		Expected Count	6.0	12.9	.1	19.0
		% within AVERAGE USE:DESIGNER	26.3%	73.7%	.0%	100.0%
		% within Gender	2.2%	2.8%	.0%	2.6%
		% of Total	.7%	1.9%	.0%	2.6%
	5 times/week	Count	2	3	0	5
		Expected Count	1.6	3.4	.0	5.0
		% within AVERAGE USE:DESIGNER	40.0%	60.0%	.0%	100.0%
		% within Gender	.9%	.6%	.0%	.7%
		% of Total	.3%	.4%	.0%	.7%
	Every day	Count	4	1	0	5
		Expected Count	1.6	3.4	.0	5.0
		% within AVERAGE USE:DESIGNER	80.0%	20.0%	.0%	100.0%
		% within Gender	1.7%	.2%	.0%	.7%
		% of Total	.5%	.1%	.0%	.7%
Total		Count	232	498	2	732
		Expected Count	232.0	498.0	2.0	732.0
		% within AVERAGE USE:DESIGNER	31.7%	68.0%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.7%	68.0%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	19.122(a)	16	.262
Likelihood Ratio	18.415	16	.300
Linear-by-Linear Association	3.983	1	.046
N of Valid Cases	732		

a 13 cells (48.1%) have expected count less than 5. The minimum expected count is .01.

# AVERAGE USE:STEROIDS \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
AVERAGE USE:STEROIDS	Never used	Count	102	168	1	271
		Expected Count	86.6	183.6	.7	271.0
		% within AVERAGE USE:STEROIDS	37.6%	62.0%	.4%	100.0%
		% within Gender	43.6%	33.9%	50.0%	37.0%
		% of Total	13.9%	23.0%	.1%	37.0%
	Once/year	Count	40	100	1	141
		Expected Count	45.1	95.5	.4	141.0
		% within AVERAGE USE:STEROIDS	28.4%	70.9%	.7%	100.0%
		% within Gender	17.1%	20.2%	50.0%	19.3%
		% of Total	5.5%	13.7%	.1%	19.3%
	6 times/year	Count	28	69	0	97
		Expected Count	31.0	65.7	.3	97.0
		% within AVERAGE USE:STEROIDS	28.9%	71.1%	.0%	100.0%
		% within Gender	12.0%	13.9%	.0%	13.3%
		% of Total	3.8%	9.4%	.0%	13.3%
	Once/month	Count	16	52	0	68
		Expected Count	21.7	46.1	.2	68.0
		% within AVERAGE USE:STEROIDS	23.5%	76.5%	.0%	100.0%
		% within Gender	6.8%	10.5%	.0%	9.3%
		% of Total	2.2%	7.1%	.0%	9.3%
	Twice/month	Count	18	35	0	53
		Expected Count	16.9	35.9	.1	53.0
		% within AVERAGE USE:STEROIDS	34.0%	66.0%	.0%	100.0%
		% within Gender	7.7%	7.1%	.0%	7.2%
		% of Total	2.5%	4.8%	.0%	7.2%
	Once/week	Count	14	33	0	47
		Expected Count	15.0	31.8	.1	47.0
		% within AVERAGE USE:STEROIDS	29.8%	70.2%	.0%	100.0%
		% within Gender	6.0%	6.7%	.0%	6.4%
		% of Total	1.9%	4.5%	.0%	6.4%

		Count	9	25	0	34
		Expected Count	10.9	23.0	.1	34.0
		% within AVERAGE USE:STEROIDS	26.5%	73.5%	.0%	100.0%
		% within Gender	3.8%	5.0%	.0%	4.6%
		% of Total	1.2%	3.4%	.0%	4.6%
	3 times/week	Count	1	7	0	8
		Expected Count	2.6	5.4	.0	8.0
		% within AVERAGE USE:STEROIDS	12.5%	87.5%	.0%	100.0%
		% within Gender	.4%	1.4%	.0%	1.1%
		% of Total	.1%	1.0%	.0%	1.1%
	5 times/week	Count	6	7	0	13
		Expected Count	4.2	8.8	.0	13.0
		% within AVERAGE USE:STEROIDS	46.2%	53.8%	.0%	100.0%
		% within Gender	2.6%	1.4%	.0%	1.8%
		% of Total	.8%	1.0%	.0%	1.8%
	Every day	Count	234	496	2	732
		Expected Count	234.0	496.0	2.0	732.0
		% within AVERAGE USE:STEROIDS	32.0%	67.8%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	32.0%	67.8%	.3%	100.0%
Total		Count	234	496	2	732
		Expected Count	234.0	496.0	2.0	732.0
		% within AVERAGE USE:STEROIDS	32.0%	67.8%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	32.0%	67.8%	.3%	100.0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	12.801(a)	16	.687
Likelihood Ratio	13.598	16	.629
Linear-by-Linear Association	1.653	1	.199
N of Valid Cases	732		

a 11 cells (40.7%) have expected count less than 5. The minimum expected count is .02.

## AVERAGE USE:OTHER \* Gender

#### Crosstab

			Gender			Total
			Male	Female	Unknown	
Never used		Count	99	218	2	319
		Expected Count	102.3	215.9	.9	319.0
		% within AVERAGE USE:OTHER	31.0%	68.3%	.6%	100.0%

AVERAGE USE:OTHER	Once/year	% within Gender	42.1%	44.0%	100.0%	43.5%
		% of Total	13.5%	29.7%	.3%	43.5%
		Count	55	98	0	153
		Expected Count	49.1	103.5	.4	153.0
		% within AVERAGE USE:OTHER	35.9%	64.1%	.0%	100.0%
	6 times/year	% within Gender	23.4%	19.8%	.0%	20.9%
		% of Total	7.5%	13.4%	.0%	20.9%
		Count	23	65	0	88
		Expected Count	28.2	59.5	.2	88.0
		% within AVERAGE USE:OTHER	26.1%	73.9%	.0%	100.0%
	Once/month	% within Gender	9.8%	13.1%	.0%	12.0%
		% of Total	3.1%	8.9%	.0%	12.0%
		Count	20	41	0	61
		Expected Count	19.6	41.3	.2	61.0
		% within AVERAGE USE:OTHER	32.8%	67.2%	.0%	100.0%
	Twice/month	% within Gender	8.5%	8.3%	.0%	8.3%
		% of Total	2.7%	5.6%	.0%	8.3%
		Count	13	24	0	37
		Expected Count	11.9	25.0	.1	37.0
		% within AVERAGE USE:OTHER	35.1%	64.9%	.0%	100.0%
	Once/week	% within Gender	5.5%	4.8%	.0%	5.0%
		% of Total	1.8%	3.3%	.0%	5.0%
		Count	15	29	0	44
		Expected Count	14.1	29.8	.1	44.0
		% within AVERAGE USE:OTHER	34.1%	65.9%	.0%	100.0%
	3 times/week	% within Gender	6.4%	5.8%	.0%	6.0%
		% of Total	2.0%	4.0%	.0%	6.0%
		Count	5	13	0	18
		Expected Count	5.8	12.2	.0	18.0
		% within AVERAGE USE:OTHER	27.8%	72.2%	.0%	100.0%
	5 times/week	% within Gender	2.1%	2.6%	.0%	2.5%
		% of Total	.7%	1.8%	.0%	2.5%
		Count	2	1	0	3
		Expected Count	1.0	2.0	.0	3.0
		% within AVERAGE USE:OTHER	66.7%	33.3%	.0%	100.0%
	Every day	% within Gender	.9%	.2%	.0%	.4%
		% of Total	.3%	.1%	.0%	.4%
		Count	3	7	0	10
		Expected Count	3.2	6.8	.0	10.0
		% within AVERAGE USE:OTHER	30.0%	70.0%	.0%	100.0%

		% within Gender	1.3%	1.4%	.0%	1.4%
		% of Total	.4%	1.0%	.0%	1.4%
Total	Count	235	496	2	733	
	Expected Count	235.0	496.0	2.0	733.0	
	% within AVERAGE USE:OTHER	32.1%	67.7%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	32.1%	67.7%	.3%	100.0%	

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	7.264(a)	16	.968
Likelihood Ratio	7.879	16	.952
Linear-by-Linear Association	.170	1	.680
N of Valid Cases	733		

a 12 cells (44.4%) have expected count less than 5. The minimum expected count is .01.

## Crosstabs

### Notes

Output Created		16-JUN-2004 09:46:57
Comments		
Input	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	747
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.

Syntax	CROSSTABS /TABLES=q20a1 q20a2 q20a3 q20a4 q20a5 q20a6 q20a7 q20a8 q20a9 q20b1 q20b2 q20b3 q20b4 q20b5 q20b6 q20b7 q20b8 q20b9 q20c1 q20c2 q20c3 q20c4 q20c5 q20c6 q20c7 q20c8 q20c9 q20d1 q20d2 q20d3 q20d4 q20d5 q20d6 q20d7 q20d8 q20d9 q20e1 q20e2 q20e3 q20e4 q20e5 q20e6 q20e7 q20e8 q20e9 q20f1 q20f2 q20f3 q20f4 q20f5 q20f6 q20f7 q20f8 q20f9 q20g1 q20g2 q20g3 q20g4 q20g5 q20g6 q20g7 q20g8 q20g9 q20h1 q20h2 q20h3 q20h4 q20h5 q20h6 q20h7 q20h8 q20h9 q20i1 q20i2 q20i3 q20i4 q20i5 q20i6 q20i7 q20i8 q20i9 q20j1 q20j2 q20j3 q20j4 q20j5 q20j6 q20j7 q20j8 q20j9 q20k1 q20k2 q20k3 q20k4 q20k5 q20k6 q20k7 q20k8 q20k9 q20l1
--------	--

		q20I2 q20I3 q20I4 q20I5 q20I6 q20I7 q20I8 q20I9 BY Q05 /STATISTIC=CHISQ /CELLS= COUNT EXPECTED ROW COLUMN TOTAL .
Resources	Dimensions Requested	2
	Cells Available	116508
	Elapsed Time	0:00:02.21

## Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Use tobacco: never * Gender	732	98.0%	15	2.0%	747	100.0%
Use tobacco: on campus * Gender	732	98.0%	15	2.0%	747	100.0%
Use tobacco: res hall * Gender	732	98.0%	15	2.0%	747	100.0%
Use tobacco: frat/sor * Gender	732	98.0%	15	2.0%	747	100.0%
Use tobacco: bar/restaurant * Gender	732	98.0%	15	2.0%	747	100.0%
Use tobacco: where live * Gender	732	98.0%	15	2.0%	747	100.0%
Use tobacco: in car * Gender	732	98.0%	15	2.0%	747	100.0%
Use tobacco: private parties * Gender	732	98.0%	15	2.0%	747	100.0%
Use tobacco: other * Gender	732	98.0%	15	2.0%	747	100.0%
Use alcohol: never * Gender	732	98.0%	15	2.0%	747	100.0%
Use alcohol: on campus * Gender	732	98.0%	15	2.0%	747	100.0%
Use alcohol: res hall * Gender	732	98.0%	15	2.0%	747	100.0%
Use alcohol: frat/sor * Gender	732	98.0%	15	2.0%	747	100.0%
Use alcohol: bar/restaurant * Gender	732	98.0%	15	2.0%	747	100.0%
Use alcohol: where live * Gender	732	98.0%	15	2.0%	747	100.0%
Use alcohol: in car * Gender	732	98.0%	15	2.0%	747	100.0%
Use alcohol: private parties * Gender	732	98.0%	15	2.0%	747	100.0%
Use alcohol: other * Gender	732	98.0%	15	2.0%	747	100.0%
Use marijuana: never * Gender	732	98.0%	15	2.0%	747	100.0%
Use marijuana: on campus * Gender	732	98.0%	15	2.0%	747	100.0%
Use marijuana: res hall * Gender	732	98.0%	15	2.0%	747	100.0%
Use marijuana: frat/sor * Gender	732	98.0%	15	2.0%	747	100.0%
Use marijuana: bar/restaurant * Gender	732	98.0%	15	2.0%	747	100.0%
Use marijuana: where live * Gender	732	98.0%	15	2.0%	747	100.0%
Use marijuana: in car * Gender	732	98.0%	15	2.0%	747	100.0%
Use marijuana: private parties * Gender	732	98.0%	15	2.0%	747	100.0%
Use marijuana: other * Gender	732	98.0%	15	2.0%	747	100.0%
Use cocaine: never * Gender	732	98.0%	15	2.0%	747	100.0%
Use cocaine: on campus * Gender	732	98.0%	15	2.0%	747	100.0%
Use cocaine: res hall * Gender	732	98.0%	15	2.0%	747	100.0%

<b>Use cocaine: frat/sor * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use cocaine: bar/restaurant * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use cocaine: where live * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use cocaine: in car * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use cocaine: private parties * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use cocaine: other * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use amphetamines: never * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use amphetamines: on campus * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use amphetamines: res hall * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use amphetamines: frat/sor * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use amphetamines: bar/restaurant * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use amphetamines: where live * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use amphetamines: in car * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use amphetamines: private parties * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use amphetamines: other * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use sedatives: never * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use sedatives: on campus * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use sedatives: res hall * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use sedatives: frat/sor * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use sedatives: bar/restaurant * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use sedatives: where live * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use sedatives: in car * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use sedatives: private parties * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use sedatives: other * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use hallucinogens: never * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use hallucinogens: on campus * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use hallucinogens: res hall * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use hallucinogens: frat/sor * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use hallucinogens: bar/restaurant * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use hallucinogens: where live * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use hallucinogens: in car * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use hallucinogens: private parties * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use hallucinogens: other * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use opiates: never * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use opiates: on campus * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use opiates: res hall * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use opiates: frat/sor * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use opiates: bar/restaurant * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use opiates: where live * Gender</b>	732	98.0%	15	2.0%	747	100.0%
<b>Use opiates: in car * Gender</b>	732	98.0%	15	2.0%	747	100.0%

Use opiates: private parties * Gender	732	98.0%	15	2.0%	747	100.0%
Use opiates: other * Gender	732	98.0%	15	2.0%	747	100.0%
Use inhalants: never * Gender	732	98.0%	15	2.0%	747	100.0%
Use inhalants: on campus * Gender	732	98.0%	15	2.0%	747	100.0%
Use inhalants: res hall * Gender	732	98.0%	15	2.0%	747	100.0%
Use inhalants: frat/sor * Gender	732	98.0%	15	2.0%	747	100.0%
Use inhalants: bar/restaurant * Gender	732	98.0%	15	2.0%	747	100.0%
Use inhalants: where live * Gender	732	98.0%	15	2.0%	747	100.0%
Use inhalants: in car * Gender	732	98.0%	15	2.0%	747	100.0%
Use inhalants: private parties * Gender	732	98.0%	15	2.0%	747	100.0%
Use inhalants: other * Gender	732	98.0%	15	2.0%	747	100.0%
Use designer drugs: never * Gender	732	98.0%	15	2.0%	747	100.0%
Use designer drugs: on campus * Gender	732	98.0%	15	2.0%	747	100.0%
Use designer drugs: res hall * Gender	732	98.0%	15	2.0%	747	100.0%
Use designer drugs: frat/sor * Gender	732	98.0%	15	2.0%	747	100.0%
Use designer drugs: bar/restaurant * Gender	732	98.0%	15	2.0%	747	100.0%
Use designer drugs: where live * Gender	732	98.0%	15	2.0%	747	100.0%
Use designer drugs: in car * Gender	732	98.0%	15	2.0%	747	100.0%
Use designer drugs: private parties * Gender	732	98.0%	15	2.0%	747	100.0%
Use designer drugs: other * Gender	732	98.0%	15	2.0%	747	100.0%
Use steroids: never * Gender	732	98.0%	15	2.0%	747	100.0%
Use steroids: on campus * Gender	732	98.0%	15	2.0%	747	100.0%
Use steroids: res hall * Gender	732	98.0%	15	2.0%	747	100.0%
Use steroids: frat/sor * Gender	732	98.0%	15	2.0%	747	100.0%
Use steroids: bar/restaurant * Gender	732	98.0%	15	2.0%	747	100.0%
Use steroids: where live * Gender	732	98.0%	15	2.0%	747	100.0%
Use steroids: in car * Gender	732	98.0%	15	2.0%	747	100.0%
Use steroids: private parties * Gender	732	98.0%	15	2.0%	747	100.0%
Use steroids: other * Gender	732	98.0%	15	2.0%	747	100.0%
Use other drugs: never * Gender	732	98.0%	15	2.0%	747	100.0%
Use other drugs: on campus * Gender	732	98.0%	15	2.0%	747	100.0%
Use other drugs: res hall * Gender	732	98.0%	15	2.0%	747	100.0%
Use other drugs: frat/sor * Gender	732	98.0%	15	2.0%	747	100.0%
Use other drugs: bar/restaurant * Gender	732	98.0%	15	2.0%	747	100.0%
Use other drugs: where live * Gender	732	98.0%	15	2.0%	747	100.0%
Use other drugs: in car * Gender	732	98.0%	15	2.0%	747	100.0%
Use other drugs: private parties * Gender	732	98.0%	15	2.0%	747	100.0%
Use other drugs: other * Gender	732	98.0%	15	2.0%	747	100.0%

## Use tobacco: never \* Gender

Crosstab

		Gender			Total	
		Male	Female	Unknown		
Use tobacco: never	No	Count	152	278	1	431
		Expected Count	137.2	292.6	1.2	431.0
		% within Use tobacco: never	35.3%	64.5%	.2%	100.0%
		% within Gender	65.2%	55.9%	50.0%	58.9%
		% of Total	20.8%	38.0%	.1%	58.9%
	Yes	Count	81	219	1	301
		Expected Count	95.8	204.4	.8	301.0
		% within Use tobacco: never	26.9%	72.8%	.3%	100.0%
		% within Gender	34.8%	44.1%	50.0%	41.1%
		% of Total	11.1%	29.9%	.1%	41.1%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use tobacco: never	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	5.733(a)	2	.057
Likelihood Ratio	5.793	2	.055
Linear-by-Linear Association	5.716	1	.017
N of Valid Cases	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .82.

## Use tobacco: on campus \* Gender

Crosstab

		Gender			Total	
		Male	Female	Unknown		
Use tobacco: on campus	No	Count	196	454	1	651
		Expected Count	207.2	442.0	1.8	651.0
		% within Use tobacco: on campus	30.1%	69.7%	.2%	100.0%
		% within Gender	84.1%	91.3%	50.0%	88.9%
		% of Total	26.8%	62.0%	.1%	88.9%
		Count	37	43	1	81
		Expected Count	25.8	55.0	.2	81.0
		% within Use tobacco: on campus	45.1%	51.9%	.1%	50.0%
		% within Gender	44.4%	50.0%	50.0%	48.9%
		% of Total	45.8%	53.1%	.1%	50.0%

		% within Use tobacco: on campus	45.7%	53.1%	1.2%	100.0%
	Yes	% within Gender	15.9%	8.7%	50.0%	11.1%
		% of Total	5.1%	5.9%	.1%	11.1%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use tobacco: on campus	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	11.510(a)	2	.003
Likelihood Ratio	9.932	2	.007
Linear-by-Linear Association	6.821	1	.009
N of Valid Cases	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .22.

## Use tobacco: res hall \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
Use tobacco: res hall	No	Count	193	440	2	635
		Expected Count	202.1	431.1	1.7	635.0
		% within Use tobacco: res hall	30.4%	69.3%	.3%	100.0%
		% within Gender	82.8%	88.5%	100.0%	86.7%
		% of Total	26.4%	60.1%	.3%	86.7%
	Yes	Count	40	57	0	97
		Expected Count	30.9	65.9	.3	97.0
		% within Use tobacco: res hall	41.2%	58.8%	.0%	100.0%
		% within Gender	17.2%	11.5%	.0%	13.3%
		% of Total	5.5%	7.8%	.0%	13.3%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use tobacco: res hall	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

--	--	--	--

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	4.788(a)	2	.091
<b>Likelihood Ratio</b>	4.881	2	.087
<b>Linear-by-Linear Association</b>	4.725	1	.030
<b>N of Valid Cases</b>	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .27.

## Use tobacco: frat/sor \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Use tobacco: frat/sor	No	Count	193	437	1	631
		Expected Count	200.9	428.4	1.7	631.0
		% within Use tobacco: frat/sor	30.6%	69.3%	.2%	100.0%
		% within Gender	82.8%	87.9%	50.0%	86.2%
		% of Total	26.4%	59.7%	.1%	86.2%
	Yes	Count	40	60	1	101
		Expected Count	32.1	68.6	.3	101.0
		% within Use tobacco: frat/sor	39.6%	59.4%	1.0%	100.0%
		% within Gender	17.2%	12.1%	50.0%	13.8%
		% of Total	5.5%	8.2%	.1%	13.8%
	Total	Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use tobacco: frat/sor	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	5.672(a)	2	.059
<b>Likelihood Ratio</b>	4.865	2	.088
<b>Linear-by-Linear Association</b>	2.631	1	.105
<b>N of Valid Cases</b>	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .28.

## Use tobacco: bar/restaurant \* Gender

Crosstab

			Gender			Total		
			Male	Female	Unknown			
Use tobacco: bar/restaurant	No	Count	146	340	1	487		
		Expected Count	155.0	330.7	1.3	487.0		
		% within Use tobacco: bar/restaurant	30.0%	69.8%	.2%	100.0%		
		% within Gender	62.7%	68.4%	50.0%	66.5%		
	Yes	% of Total	19.9%	46.4%	.1%	66.5%		
		Count	87	157	1	245		
		Expected Count	78.0	166.3	.7	245.0		
		% within Use tobacco: bar/restaurant	35.5%	64.1%	.4%	100.0%		
	Total	% within Gender	37.3%	31.6%	50.0%	33.5%		
		% of Total	11.9%	21.4%	.1%	33.5%		
			Count	233	497	2	732	
			Expected Count	233.0	497.0	2.0	732.0	
			% within Use tobacco: bar/restaurant	31.8%	67.9%	.3%	100.0%	
			% within Gender	100.0%	100.0%	100.0%	100.0%	
			% of Total	31.8%	67.9%	.3%	100.0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	2.601(a)	2	.272
Likelihood Ratio	2.567	2	.277
Linear-by-Linear Association	2.087	1	.149
N of Valid Cases	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .67.

## Use tobacco: where live \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Use tobacco: where live	No	Count	144	342	1	487
		Expected Count	155.0	330.7	1.3	487.0
		% within Use tobacco: where live	29.6%	70.2%	.2%	100.0%
		% within Gender	61.8%	68.8%	50.0%	66.5%
	Yes	% of Total	19.7%	46.7%	.1%	66.5%
		Count	89	155	1	245
		Expected Count	78.0	166.3	.7	245.0
		% of Total	36.7%	62.7%	.4%	66.5%

		% within Use tobacco: where live	36.3%	63.3%	.4%	100.0%
	Yes	% within Gender	38.2%	31.2%	50.0%	33.5%
		% of Total	12.2%	21.2%	.1%	33.5%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use tobacco: where live	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	3.747(a)	2	.154
Likelihood Ratio	3.698	2	.157
Linear-by-Linear Association	3.158	1	.076
N of Valid Cases	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .67.

## Use tobacco: in car \* Gender

## Crosstab

			Gender			Total		
			Male	Female	Unknown			
Use tobacco: in car	No	Count	144	336	1	481		
		Expected Count	153.1	326.6	1.3	481.0		
		% within Use tobacco: in car	29.9%	69.9%	.2%	100.0%		
		% within Gender	61.8%	67.6%	50.0%	65.7%		
		% of Total	19.7%	45.9%	.1%	65.7%		
	Yes	Count	89	161	1	251		
		Expected Count	79.9	170.4	.7	251.0		
		% within Use tobacco: in car	35.5%	64.1%	.4%	100.0%		
		% within Gender	38.2%	32.4%	50.0%	34.3%		
		% of Total	12.2%	22.0%	.1%	34.3%		
				233	497	2	732	
				233.0	497.0	2.0	732.0	
				% within Use tobacco: in car	31.8%	67.9%	.3%	100.0%
				% within Gender	100.0%	100.0%	100.0%	100.0%
				% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

--	--	--	--

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	2.591(a)	2	.274
<b>Likelihood Ratio</b>	2.560	2	.278
<b>Linear-by-Linear Association</b>	2.113	1	.146
<b>N of Valid Cases</b>	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .69.

## Use tobacco: private parties \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Use tobacco: private parties	No	Count	125	321	1	447
		Expected Count	142.3	303.5	1.2	447.0
		% within Use tobacco: private parties	28.0%	71.8%	.2%	100.0%
		% within Gender	53.6%	64.6%	50.0%	61.1%
		% of Total	17.1%	43.9%	.1%	61.1%
	Yes	Count	108	176	1	285
		Expected Count	90.7	193.5	.8	285.0
		% within Use tobacco: private parties	37.9%	61.8%	.4%	100.0%
		% within Gender	46.4%	35.4%	50.0%	38.9%
		% of Total	14.8%	24.0%	.1%	38.9%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use tobacco: private parties	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	8.088(a)	2	.018
<b>Likelihood Ratio</b>	8.016	2	.018
<b>Linear-by-Linear Association</b>	7.543	1	.006
<b>N of Valid Cases</b>	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .78.

## Use tobacco: other \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Use tobacco: other	No	Count	185	446	1	632
		Expected Count	201.2	429.1	1.7	632.0
		% within Use tobacco: other	29.3%	70.6%	.2%	100.0%
		% within Gender	79.4%	89.7%	50.0%	86.3%
		% of Total	25.3%	60.9%	.1%	86.3%
	Yes	Count	48	51	1	100
		Expected Count	31.8	67.9	.3	100.0
		% within Use tobacco: other	48.0%	51.0%	1.0%	100.0%
		% within Gender	20.6%	10.3%	50.0%	13.7%
		% of Total	6.6%	7.0%	.1%	13.7%
Total	Count	233	497	2	732	
	Expected Count	233.0	497.0	2.0	732.0	
	% within Use tobacco: other	31.8%	67.9%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	31.8%	67.9%	.3%	100.0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	16.623(a)	2	.000
Likelihood Ratio	15.195	2	.001
Linear-by-Linear Association	12.456	1	.000
N of Valid Cases	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .27.

## Use alcohol: never \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Use alcohol: never	No	Count	221	463	2	686
		Expected Count	218.4	465.8	1.9	686.0
		% within Use alcohol: never	32.2%	67.5%	.3%	100.0%
		% within Gender	94.8%	93.2%	100.0%	93.7%
		% of Total	30.2%	63.3%	.3%	93.7%
	Yes	Count	12	34	0	46
		Expected Count	14.6	31.2	.1	46.0
		% within Use alcohol: never	26.1%	73.9%	.2%	100.0%
		% within Gender	26.1%	73.9%	100.0%	93.7%
		% of Total	26.1%	73.9%	.2%	93.7%

		% within Use alcohol: never	26.1%	73.9%	.0%	100.0%
	Yes	% within Gender	5.2%	6.8%	.0%	6.3%
		% of Total	1.6%	4.6%	.0%	6.3%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use alcohol: never	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.905(a)	2	.636
Likelihood Ratio	1.055	2	.590
Linear-by-Linear Association	.662	1	.416
N of Valid Cases	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .13.

## Use alcohol: on campus \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
Use alcohol: on campus	No	Count	187	445	1	633
		Expected Count	201.5	429.8	1.7	633.0
		% within Use alcohol: on campus	29.5%	70.3%	.2%	100.0%
		% within Gender	80.3%	89.5%	50.0%	86.5%
		% of Total	25.5%	60.8%	.1%	86.5%
	Yes	Count	46	52	1	99
		Expected Count	31.5	67.2	.3	99.0
		% within Use alcohol: on campus	46.5%	52.5%	1.0%	100.0%
		% within Gender	19.7%	10.5%	50.0%	13.5%
		% of Total	6.3%	7.1%	.1%	13.5%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use alcohol: on campus	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

--	--	--	--

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	13.961(a)	2	.001
<b>Likelihood Ratio</b>	12.683	2	.002
<b>Linear-by-Linear Association</b>	9.971	1	.002
<b>N of Valid Cases</b>	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .27.

## Use alcohol: res hall \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Use alcohol: res hall	No	Count	110	305	2	417
		Expected Count	132.7	283.1	1.1	417.0
		% within Use alcohol: res hall	26.4%	73.1%	.5%	100.0%
		% within Gender	47.2%	61.4%	100.0%	57.0%
		% of Total	15.0%	41.7%	.3%	57.0%
	Yes	Count	123	192	0	315
		Expected Count	100.3	213.9	.9	315.0
		% within Use alcohol: res hall	39.0%	61.0%	.0%	100.0%
		% within Gender	52.8%	38.6%	.0%	43.0%
		% of Total	16.8%	26.2%	.0%	43.0%
Total	Count		233	497	2	732
	Expected Count		233.0	497.0	2.0	732.0
	% within Use alcohol: res hall		31.8%	67.9%	.3%	100.0%
	% within Gender		100.0%	100.0%	100.0%	100.0%
	% of Total		31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	14.486(a)	2	.001
<b>Likelihood Ratio</b>	15.157	2	.001
<b>Linear-by-Linear Association</b>	13.990	1	.000
<b>N of Valid Cases</b>	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .86.

## Use alcohol: frat/sor \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Use alcohol: frat/sor	No	Count	125	329	1	455
		Expected Count	144.8	308.9	1.2	455.0
		% within Use alcohol: frat/sor	27.5%	72.3%	.2%	100.0%
		% within Gender	53.6%	66.2%	50.0%	62.2%
	Yes	% of Total	17.1%	44.9%	.1%	62.2%
		Count	108	168	1	277
		Expected Count	88.2	188.1	.8	277.0
		% within Use alcohol: frat/sor	39.0%	60.6%	.4%	100.0%
Total	Yes	% within Gender	46.4%	33.8%	50.0%	37.8%
		% of Total	14.8%	23.0%	.1%	37.8%
		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use alcohol: frat/sor	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	10.747(a)	2	.005
Likelihood Ratio	10.628	2	.005
Linear-by-Linear Association	10.047	1	.002
N of Valid Cases	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .76.

## Use alcohol: bar/restaurant \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Use alcohol: bar/restaurant	No	Count	87	226	0	313
		Expected Count	99.6	212.5	.9	313.0
		% within Use alcohol: bar/restaurant	27.8%	72.2%	.0%	100.0%
		% within Gender	37.3%	45.5%	.0%	42.8%
	Yes	% of Total	11.9%	30.9%	.0%	42.8%
		Count	146	271	2	419
		Expected Count	133.4	284.5	1.1	419.0

		% within Use alcohol: bar/restaurant	34.8%	64.7%	.5%	100.0%
	Yes	% within Gender	62.7%	54.5%	100.0%	57.2%
		% of Total	19.9%	37.0%	.3%	57.2%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use alcohol: bar/restaurant	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	5.786(a)	2	.055
Likelihood Ratio	6.553	2	.038
Linear-by-Linear Association	3.490	1	.062
N of Valid Cases	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .86.

## Use alcohol: where live \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
Use alcohol: where live	No	Count	71	206	1	278
		Expected Count	88.5	188.8	.8	278.0
		% within Use alcohol: where live	25.5%	74.1%	.4%	100.0%
		% within Gender	30.5%	41.4%	50.0%	38.0%
		% of Total	9.7%	28.1%	.1%	38.0%
	Yes	Count	162	291	1	454
		Expected Count	144.5	308.2	1.2	454.0
		% within Use alcohol: where live	35.7%	64.1%	.2%	100.0%
		% within Gender	69.5%	58.6%	50.0%	62.0%
		% of Total	22.1%	39.8%	.1%	62.0%
				233	497	2
				233.0	497.0	2.0
				% within Use alcohol: where live	31.8%	67.9%
				% within Gender	100.0%	100.0%
				% of Total	31.8%	67.9%

## Chi-Square Tests

--	--	--	--

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	8.237(a)	2	.016
<b>Likelihood Ratio</b>	8.379	2	.015
<b>Linear-by-Linear Association</b>	8.221	1	.004
<b>N of Valid Cases</b>	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .76.

## Use alcohol: in car \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Use alcohol: in car	No	Count	183	442	2	627
		Expected Count	199.6	425.7	1.7	627.0
		% within Use alcohol: in car	29.2%	70.5%	.3%	100.0%
		% within Gender	78.5%	88.9%	100.0%	85.7%
		% of Total	25.0%	60.4%	.3%	85.7%
	Yes	Count	50	55	0	105
		Expected Count	33.4	71.3	.3	105.0
		% within Use alcohol: in car	47.6%	52.4%	.0%	100.0%
		% within Gender	21.5%	11.1%	.0%	14.3%
		% of Total	6.8%	7.5%	.0%	14.3%
	Total	Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use alcohol: in car	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	14.281(a)	2	.001
<b>Likelihood Ratio</b>	13.820	2	.001
<b>Linear-by-Linear Association</b>	14.261	1	.000
<b>N of Valid Cases</b>	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .29.

## Use alcohol: private parties \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Use alcohol: private parties	No	Count	57	132	0	189
		Expected Count	60.2	128.3	.5	189.0
		% within Use alcohol: private parties	30.2%	69.8%	.0%	100.0%
		% within Gender	24.5%	26.6%	.0%	25.8%
	Yes	% of Total	7.8%	18.0%	.0%	25.8%
		Count	176	365	2	543
		Expected Count	172.8	368.7	1.5	543.0
		% within Use alcohol: private parties	32.4%	67.2%	.4%	100.0%
Total	No	% within Gender	75.5%	73.4%	100.0%	74.2%
		% of Total	24.0%	49.9%	.3%	74.2%
		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use alcohol: private parties	31.8%	67.9%	.3%	100.0%
	Yes	% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1.062(a)	2	.588
Likelihood Ratio	1.562	2	.458
Linear-by-Linear Association	.225	1	.635
N of Valid Cases	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .52.

## Use alcohol: other \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Use alcohol: other	No	Count	172	432	1	605
		Expected Count	192.6	410.8	1.7	605.0
		% within Use alcohol: other	28.4%	71.4%	.2%	100.0%
		% within Gender	73.8%	86.9%	50.0%	82.7%
	Yes	% of Total	23.5%	59.0%	.1%	82.7%
		Count	61	65	1	127
		Expected Count	40.4	86.2	.3	127.0

		% within Use alcohol: other	48.0%	51.2%	.8%	100.0%
	Yes	% within Gender	26.2%	13.1%	50.0%	17.3%
		% of Total	8.3%	8.9%	.1%	17.3%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use alcohol: other	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	20.480(a)	2	.000
Likelihood Ratio	19.232	2	.000
Linear-by-Linear Association	17.051	1	.000
N of Valid Cases	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .35.

## Use marijuana: never \* Gender

### Crosstab

			Gender			Total
			Male	Female	Unknown	
Use marijuana: never	No	Count	120	219	0	339
		Expected Count	107.9	230.2	.9	339.0
		% within Use marijuana: never	35.4%	64.6%	.0%	100.0%
		% within Gender	51.5%	44.1%	.0%	46.3%
		% of Total	16.4%	29.9%	.0%	46.3%
	Yes	Count	113	278	2	393
		Expected Count	125.1	266.8	1.1	393.0
		% within Use marijuana: never	28.8%	70.7%	.5%	100.0%
		% within Gender	48.5%	55.9%	100.0%	53.7%
		% of Total	15.4%	38.0%	.3%	53.7%
Total						

### Chi-Square Tests

--	--	--	--

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	5.259(a)	2	.072
<b>Likelihood Ratio</b>	6.016	2	.049
<b>Linear-by-Linear Association</b>	4.201	1	.040
<b>N of Valid Cases</b>	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .93.

## Use marijuana: on campus \* Gender

Crosstab

		Gender			Total
		Male	Female	Unknown	
Use marijuana: on campus	No	Count	218	490	2 710
	No	Expected Count	226.0	482.1	1.9 710.0
	No	% within Use marijuana: on campus	30.7%	69.0%	.3% 100.0%
	No	% within Gender	93.6%	98.6%	100.0% 97.0%
	No	% of Total	29.8%	66.9%	.3% 97.0%
	Yes	Count	15	7	0 22
	Yes	Expected Count	7.0	14.9	.1 22.0
	Yes	% within Use marijuana: on campus	68.2%	31.8%	.0% 100.0%
	Yes	% within Gender	6.4%	1.4%	.0% 3.0%
	Yes	% of Total	2.0%	1.0%	.0% 3.0%
Total		Count	233	497	2 732
		Expected Count	233.0	497.0	2.0 732.0
		% within Use marijuana: on campus	31.8%	67.9%	.3% 100.0%
		% within Gender	100.0%	100.0%	100.0% 100.0%
		% of Total	31.8%	67.9%	.3% 100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	13.826(a)	2	.001
<b>Likelihood Ratio</b>	12.660	2	.002
<b>Linear-by-Linear Association</b>	13.720	1	.000
<b>N of Valid Cases</b>	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .06.

## Use marijuana: res hall \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Use marijuana: res hall	No	Count	200	459	2	661
		Expected Count	210.4	448.8	1.8	661.0
		% within Use marijuana: res hall	30.3%	69.4%	.3%	100.0%
		% within Gender	85.8%	92.4%	100.0%	90.3%
	Yes	% of Total	27.3%	62.7%	.3%	90.3%
		Count	33	38	0	71
		Expected Count	22.6	48.2	.2	71.0
		% within Use marijuana: res hall	46.5%	53.5%	.0%	100.0%
Total	No	% within Gender	14.2%	7.6%	.0%	9.7%
		% of Total	4.5%	5.2%	.0%	9.7%
		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use marijuana: res hall	31.8%	67.9%	.3%	100.0%
	Yes	% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	7.908(a)	2	.019
Likelihood Ratio	7.678	2	.022
Linear-by-Linear Association	7.894	1	.005
N of Valid Cases	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .19.

## Use marijuana: frat/sor \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Use marijuana: frat/sor	No	Count	209	468	2	679
		Expected Count	216.1	461.0	1.9	679.0
		% within Use marijuana: frat/sor	30.8%	68.9%	.3%	100.0%
		% within Gender	89.7%	94.2%	100.0%	92.8%
	Yes	% of Total	28.6%	63.9%	.3%	92.8%
		Count	24	29	0	53
		Expected Count	16.9	36.0	.1	53.0

Total	Yes	% within Use marijuana: frat/sor	45.3%	54.7%	.0%	100.0%	
		% within Gender	10.3%	5.8%	.0%	7.2%	
		% of Total	3.3%	4.0%	.0%	7.2%	
	Total	Count	233	497	2	732	
		Expected Count	233.0	497.0	2.0	732.0	
		% within Use marijuana: frat/sor	31.8%	67.9%	.3%	100.0%	
		% within Gender	100.0%	100.0%	100.0%	100.0%	
		% of Total	31.8%	67.9%	.3%	100.0%	

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	4.866(a)	2	.088
Likelihood Ratio	4.758	2	.093
Linear-by-Linear Association	4.854	1	.028
N of Valid Cases	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .14.

## Use marijuana: bar/restaurant \* Gender

**Crosstab**

		Gender			Total	
		Male	Female	Unknown		
Use marijuana: bar/restaurant	No	Count	222	487	2	711
		Expected Count	226.3	482.7	1.9	711.0
		% within Use marijuana: bar/restaurant	31.2%	68.5%	.3%	100.0%
		% within Gender	95.3%	98.0%	100.0%	97.1%
		% of Total	30.3%	66.5%	.3%	97.1%
	Yes	Count	11	10	0	21
		Expected Count	6.7	14.3	.1	21.0
		% within Use marijuana: bar/restaurant	52.4%	47.6%	.0%	100.0%
		% within Gender	4.7%	2.0%	.0%	2.9%
		% of Total	1.5%	1.4%	.0%	2.9%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use marijuana: bar/restaurant	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	4.237(a)	2	.120
<b>Likelihood Ratio</b>	3.986	2	.136
<b>Linear-by-Linear Association</b>	4.228	1	.040
<b>N of Valid Cases</b>	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .06.

**Use marijuana: where live \* Gender****Crosstab**

			Gender			Total
			Male	Female	Unknown	
Use marijuana: where live	No	Count	166	390	2	558
		Expected Count	177.6	378.9	1.5	558.0
		% within Use marijuana: where live	29.7%	69.9%	.4%	100.0%
		% within Gender	71.2%	78.5%	100.0%	76.2%
		% of Total	22.7%	53.3%	.3%	76.2%
	Yes	Count	67	107	0	174
		Expected Count	55.4	118.1	.5	174.0
		% within Use marijuana: where live	38.5%	61.5%	.0%	100.0%
		% within Gender	28.8%	21.5%	.0%	23.8%
		% of Total	9.2%	14.6%	.0%	23.8%
		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use marijuana: where live	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	5.197(a)	2	.074
<b>Likelihood Ratio</b>	5.553	2	.062
<b>Linear-by-Linear Association</b>	4.970	1	.026
<b>N of Valid Cases</b>	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .48.

**Use marijuana: in car \* Gender**

Crosstab

		Gender			Total	
		Male	Female	Unknown		
Use marijuana: in car	No	Count	178	416	2	596
		Expected Count	189.7	404.7	1.6	596.0
		% within Use marijuana: in car	29.9%	69.8%	.3%	100.0%
		% within Gender	76.4%	83.7%	100.0%	81.4%
		% of Total	24.3%	56.8%	.3%	81.4%
	Yes	Count	55	81	0	136
		Expected Count	43.3	92.3	.4	136.0
		% within Use marijuana: in car	40.4%	59.6%	.0%	100.0%
		% within Gender	23.6%	16.3%	.0%	18.6%
		% of Total	7.5%	11.1%	.0%	18.6%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use marijuana: in car	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	6.057(a)	2	.048
Likelihood Ratio	6.244	2	.044
Linear-by-Linear Association	5.945	1	.015
N of Valid Cases	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .37.

## Use marijuana: private parties \* Gender

Crosstab

		Gender			Total	
		Male	Female	Unknown		
Use marijuana: private parties	No	Count	150	348	2	500
		Expected Count	159.2	339.5	1.4	500.0
		% within Use marijuana: private parties	30.0%	69.6%	.4%	100.0%
		% within Gender	64.4%	70.0%	100.0%	68.3%
		% of Total	20.5%	47.5%	.3%	68.3%
		Count	83	149	0	232

		Expected Count	73.8	157.5	.6	232.0
	Yes	% within Use marijuana: private parties	35.8%	64.2%	.0%	100.0%
		% within Gender	35.6%	30.0%	.0%	31.7%
		% of Total	11.3%	20.4%	.0%	31.7%
		Count	233	497	2	732
Total		Expected Count	233.0	497.0	2.0	732.0
		% within Use marijuana: private parties	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	3.263(a)	2	.196
Likelihood Ratio	3.833	2	.147
Linear-by-Linear Association	2.726	1	.099
N of Valid Cases	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .63.

## Use marijuana: other \* Gender

### Crosstab

		Gender			Total	
		Male	Female	Unknown		
Use marijuana: other	No	Count	192	463	2	657
		Expected Count	209.1	446.1	1.8	657.0
		% within Use marijuana: other	29.2%	70.5%	.3%	100.0%
		% within Gender	82.4%	93.2%	100.0%	89.8%
		% of Total	26.2%	63.3%	.3%	89.8%
	Yes	Count	41	34	0	75
		Expected Count	23.9	50.9	.2	75.0
		% within Use marijuana: other	54.7%	45.3%	.0%	100.0%
		% within Gender	17.6%	6.8%	.0%	10.2%
		% of Total	5.6%	4.6%	.0%	10.2%
	Total	Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use marijuana: other	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	20.184(a)	2	.000
<b>Likelihood Ratio</b>	18.979	2	.000
<b>Linear-by-Linear Association</b>	20.124	1	.000
<b>N of Valid Cases</b>	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .20.

**Use cocaine: never \* Gender****Crosstab**

		Gender			Total	
		Male	Female	Unknown		
Use cocaine: never	No	Count	20	38	0 58	
		Expected Count	18.5	39.4	.2 58.0	
		% within Use cocaine: never	34.5%	65.5%	.0% 100.0%	
		% within Gender	8.6%	7.6%	.0% 7.9%	
	Yes	% of Total	2.7%	5.2%	.0% 7.9%	
		Count	213	459	2 674	
		Expected Count	214.5	457.6	1.8 674.0	
		% within Use cocaine: never	31.6%	68.1%	.3% 100.0%	
	Total	% within Gender	91.4%	92.4%	100.0% 92.1%	
		% of Total	29.1%	62.7%	.3% 92.1%	
		Count	233	497	2 732	
		Expected Count	233.0	497.0	2.0 732.0	
		% within Use cocaine: never	31.8%	67.9%	.3% 100.0%	
		% within Gender	100.0%	100.0%	100.0% 100.0%	
		% of Total	31.8%	67.9%	.3% 100.0%	

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	.364(a)	2	.834
<b>Likelihood Ratio</b>	.519	2	.771
<b>Linear-by-Linear Association</b>	.243	1	.622
<b>N of Valid Cases</b>	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .16.

**Use cocaine: on campus \* Gender**

Crosstab

			Gender			Total
			Male	Female	Unknown	
Use cocaine: on campus	No	Count	232	497	2	731
		Expected Count	232.7	496.3	2.0	731.0
		% within Use cocaine: on campus	31.7%	68.0%	.3%	100.0%
		% within Gender	99.6%	100.0%	100.0%	99.9%
		% of Total	31.7%	67.9%	.3%	99.9%
	Yes	Count	1	0	0	1
		Expected Count	.3	.7	.0	1.0
		% within Use cocaine: on campus	100.0%	.0%	.0%	100.0%
		% within Gender	.4%	.0%	.0%	.1%
		% of Total	.1%	.0%	.0%	.1%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use cocaine: on campus	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	2.145(a)	2	.342
Likelihood Ratio	2.292	2	.318
Linear-by-Linear Association	2.115	1	.146
N of Valid Cases	732		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .00.

## Use cocaine: res hall \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Use cocaine: res hall	No	Count	231	493	2	726
		Expected Count	231.1	492.9	2.0	726.0
		% within Use cocaine: res hall	31.8%	67.9%	.3%	100.0%
		% within Gender	99.1%	99.2%	100.0%	99.2%
		% of Total	31.6%	67.3%	.3%	99.2%
		Count	2	4	0	6
		Expected Count	1.9	4.1	.0	6.0
		% within Use cocaine: res hall	33.3%	66.7%	.0%	100.0%
		% within Gender	50.0%	50.0%	0.0%	100.0%
		% of Total	33.3%	66.7%	.0%	100.0%

		% within Use cocaine: res hall	33.3%	66.7%	.0%	100.0%
	Yes	% within Gender	.9%	.8%	.0%	.8%
		% of Total	.3%	.5%	.0%	.8%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use cocaine: res hall	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.022(a)	2	.989
Likelihood Ratio	.039	2	.981
Linear-by-Linear Association	.009	1	.926
N of Valid Cases	732		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .02.

## Use cocaine: frat/sor \* Gender

### Crosstab

			Gender			Total
			Male	Female	Unknown	
Use cocaine: frat/sor	No	Count	229	496	2	727
		Expected Count	231.4	493.6	2.0	727.0
		% within Use cocaine: frat/sor	31.5%	68.2%	.3%	100.0%
		% within Gender	98.3%	99.8%	100.0%	99.3%
		% of Total	31.3%	67.8%	.3%	99.3%
	Yes	Count	4	1	0	5
		Expected Count	1.6	3.4	.0	5.0
		% within Use cocaine: frat/sor	80.0%	20.0%	.0%	100.0%
		% within Gender	1.7%	.2%	.0%	.7%
		% of Total	.5%	.1%	.0%	.7%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use cocaine: frat/sor	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

### Chi-Square Tests

--	--	--	--

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	5.385(a)	2	.068
<b>Likelihood Ratio</b>	4.965	2	.084
<b>Linear-by-Linear Association</b>	5.328	1	.021
<b>N of Valid Cases</b>	732		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .01.

## Use cocaine: bar/restaurant \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Use cocaine: bar/restaurant	No	Count	228	490	2	720
		Expected Count	229.2	488.9	2.0	720.0
		% within Use cocaine: bar/restaurant	31.7%	68.1%	.3%	100.0%
		% within Gender	97.9%	98.6%	100.0%	98.4%
		% of Total	31.1%	66.9%	.3%	98.4%
	Yes	Count	5	7	0	12
		Expected Count	3.8	8.1	.0	12.0
		% within Use cocaine: bar/restaurant	41.7%	58.3%	.0%	100.0%
		% within Gender	2.1%	1.4%	.0%	1.6%
		% of Total	.7%	1.0%	.0%	1.6%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use cocaine: bar/restaurant	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	.568(a)	2	.753
<b>Likelihood Ratio</b>	.577	2	.749
<b>Linear-by-Linear Association</b>	.562	1	.453
<b>N of Valid Cases</b>	732		

a 3 cells (50.0%) have expected count less than 5. The minimum expected count is .03.

## Use cocaine: where live \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Use cocaine: where live	No	Count	221	486	2	709
		Expected Count	225.7	481.4	1.9	709.0
		% within Use cocaine: where live	31.2%	68.5%	.3%	100.0%
		% within Gender	94.8%	97.8%	100.0%	96.9%
	Yes	% of Total	30.2%	66.4%	.3%	96.9%
		Count	12	11	0	23
		Expected Count	7.3	15.6	.1	23.0
		% within Use cocaine: where live	52.2%	47.8%	.0%	100.0%
Total	No	% within Gender	5.2%	2.2%	.0%	3.1%
		% of Total	1.6%	1.5%	.0%	3.1%
		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use cocaine: where live	31.8%	67.9%	.3%	100.0%
	Yes	% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	4.561(a)	2	.102
Likelihood Ratio	4.295	2	.117
Linear-by-Linear Association	4.551	1	.033
N of Valid Cases	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .06.

## Use cocaine: in car \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Use cocaine: in car	No	Count	227	491	2	720
		Expected Count	229.2	488.9	2.0	720.0
		% within Use cocaine: in car	31.5%	68.2%	.3%	100.0%
		% within Gender	97.4%	98.8%	100.0%	98.4%
	Yes	% of Total	31.0%	67.1%	.3%	98.4%
		Count	6	6	0	12
		Expected Count	3.8	8.1	.0	12.0

		% within Use cocaine: in car	50.0%	50.0%	.0%	100.0%
		% within Gender	2.6%	1.2%	.0%	1.6%
		% of Total	.8%	.8%	.0%	1.6%
Total	Yes	Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use cocaine: in car	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1.874(a)	2	.392
Likelihood Ratio	1.778	2	.411
Linear-by-Linear Association	1.871	1	.171
N of Valid Cases	732		

a 3 cells (50.0%) have expected count less than 5. The minimum expected count is .03.

## Use cocaine: private parties \* Gender

## Crosstab

		Gender			Total	
		Male	Female	Unknown		
Use cocaine: private parties	No	Count	221	469	2	692
		Expected Count	220.3	469.8	1.9	692.0
		% within Use cocaine: private parties	31.9%	67.8%	.3%	100.0%
		% within Gender	94.8%	94.4%	100.0%	94.5%
		% of Total	30.2%	64.1%	.3%	94.5%
	Yes	Count	12	28	0	40
		Expected Count	12.7	27.2	.1	40.0
		% within Use cocaine: private parties	30.0%	70.0%	.0%	100.0%
		% within Gender	5.2%	5.6%	.0%	5.5%
		% of Total	1.6%	3.8%	.0%	5.5%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use cocaine: private parties	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

--	--	--	--

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	.188(a)	2	.910
<b>Likelihood Ratio</b>	.297	2	.862
<b>Linear-by-Linear Association</b>	.046	1	.830
<b>N of Valid Cases</b>	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .11.

## Use cocaine: other \* Gender

Crosstab

			Gender			Total	
			Male	Female	Unknown		
Use cocaine: other	No	Count	228	488	2	718	
		Expected Count	228.5	487.5	2.0	718.0	
		% within Use cocaine: other	31.8%	68.0%	.3%	100.0%	
		% within Gender	97.9%	98.2%	100.0%	98.1%	
		% of Total	31.1%	66.7%	.3%	98.1%	
	Yes	Count	5	9	0	14	
		Expected Count	4.5	9.5	.0	14.0	
		% within Use cocaine: other	35.7%	64.3%	.0%	100.0%	
		% within Gender	2.1%	1.8%	.0%	1.9%	
		% of Total	.7%	1.2%	.0%	1.9%	
Total		Count	233	497	2	732	
		Expected Count	233.0	497.0	2.0	732.0	
		% within Use cocaine: other	31.8%	67.9%	.3%	100.0%	
		% within Gender	100.0%	100.0%	100.0%	100.0%	
		% of Total	31.8%	67.9%	.3%	100.0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	.134(a)	2	.935
<b>Likelihood Ratio</b>	.170	2	.918
<b>Linear-by-Linear Association</b>	.111	1	.739
<b>N of Valid Cases</b>	732		

a 3 cells (50.0%) have expected count less than 5. The minimum expected count is .04.

## Use amphetamines: never \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Use amphetamines: never	No	Count	16	74	0	90
		Expected Count	28.6	61.1	.2	90.0
		% within Use amphetamines: never	17.8%	82.2%	.0%	100.0%
		% within Gender	6.9%	14.9%	.0%	12.3%
	Yes	% of Total	2.2%	10.1%	.0%	12.3%
		Count	217	423	2	642
		Expected Count	204.4	435.9	1.8	642.0
		% within Use amphetamines: never	33.8%	65.9%	.3%	100.0%
Total	No	% within Gender	93.1%	85.1%	100.0%	87.7%
		% of Total	29.6%	57.8%	.3%	87.7%
		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use amphetamines: never	31.8%	67.9%	.3%	100.0%
	Yes	% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	9.749(a)	2	.008
Likelihood Ratio	10.880	2	.004
Linear-by-Linear Association	8.787	1	.003
N of Valid Cases	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .25.

## Use amphetamines: on campus \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Use amphetamines: on campus	No	Count	231	494	2	727
		Expected Count	231.4	493.6	2.0	727.0
		% within Use amphetamines: on campus	31.8%	68.0%	.3%	100.0%
		% within Gender	99.1%	99.4%	100.0%	99.3%
	Yes	% of Total	31.6%	67.5%	.3%	99.3%
		Count	2	3	0	5
		Expected Count	1.6	3.4	.0	5.0

		% within Use amphetamines: on campus	40.0%	60.0%	.0%	100.0%
	Yes	% within Gender	.9%	.6%	.0%	.7%
		% of Total	.3%	.4%	.0%	.7%
		Count	233	497	2	732
Total		Expected Count	233.0	497.0	2.0	732.0
		% within Use amphetamines: on campus	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.166(a)	2	.921
Likelihood Ratio	.173	2	.917
Linear-by-Linear Association	.162	1	.687
N of Valid Cases	732		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .01.

## Use amphetamines: res hall \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
Use amphetamines: res hall	No	Count	230	483	2	715
		Expected Count	227.6	485.5	2.0	715.0
		% within Use amphetamines: res hall	32.2%	67.6%	.3%	100.0%
		% within Gender	98.7%	97.2%	100.0%	97.7%
		% of Total	31.4%	66.0%	.3%	97.7%
	Yes	Count	3	14	0	17
		Expected Count	5.4	11.5	.0	17.0
		% within Use amphetamines: res hall	17.6%	82.4%	.0%	100.0%
		% within Gender	1.3%	2.8%	.0%	2.3%
		% of Total	.4%	1.9%	.0%	2.3%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use amphetamines: res hall	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1.683(a)	2	.431
Likelihood Ratio	1.905	2	.386
Linear-by-Linear Association	1.519	1	.218
N of Valid Cases	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .05.

**Use amphetamines: frat/sor \* Gender**

## Crosstab

			Gender			Total
			Male	Female	Unknown	
Use amphetamines: frat/sor	No	Count	232	497	2	731
		Expected Count	232.7	496.3	2.0	731.0
		% within Use amphetamines: frat/sor	31.7%	68.0%	.3%	100.0%
		% within Gender	99.6%	100.0%	100.0%	99.9%
		% of Total	31.7%	67.9%	.3%	99.9%
	Yes	Count	1	0	0	1
		Expected Count	.3	.7	.0	1.0
		% within Use amphetamines: frat/sor	100.0%	.0%	.0%	100.0%
		% within Gender	.4%	.0%	.0%	.1%
		% of Total	.1%	.0%	.0%	.1%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use amphetamines: frat/sor	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	2.145(a)	2	.342
Likelihood Ratio	2.292	2	.318
Linear-by-Linear Association	2.115	1	.146
N of Valid Cases	732		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .00.

**Use amphetamines: bar/restaurant \* Gender**

Crosstab

			Gender			Total
			Male	Female	Unknown	
Use amphetamines: bar/restaurant	No	Count	231	495	2	728
		Expected Count	231.7	494.3	2.0	728.0
		% within Use amphetamines: bar/restaurant	31.7%	68.0%	.3%	100.0%
		% within Gender	99.1%	99.6%	100.0%	99.5%
		% of Total	31.6%	67.6%	.3%	99.5%
	Yes	Count	2	2	0	4
		Expected Count	1.3	2.7	.0	4.0
		% within Use amphetamines: bar/restaurant	50.0%	50.0%	.0%	100.0%
		% within Gender	.9%	.4%	.0%	.5%
		% of Total	.3%	.3%	.0%	.5%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use amphetamines: bar/restaurant	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.618(a)	2	.734
Likelihood Ratio	.586	2	.746
Linear-by-Linear Association	.617	1	.432
N of Valid Cases	732		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .01.

## Use amphetamines: where live \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
No	No	Count	224	441	2	667
		Expected Count	212.3	452.9	1.8	667.0
		% within Use amphetamines: where live	33.6%	66.1%	.3%	100.0%
		% within Gender	96.1%	88.7%	100.0%	91.1%

Use amphetamines: where live	Yes	% of Total	30.6%	60.2%	.3%	91.1%
		Count	9	56	0	65
		Expected Count	20.7	44.1	.2	65.0
		% within Use amphetamines: where live	13.8%	86.2%	.0%	100.0%
		% within Gender	3.9%	11.3%	.0%	8.9%
		% of Total	1.2%	7.7%	.0%	8.9%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use amphetamines: where live	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	10.946(a)	2	.004
Likelihood Ratio	12.652	2	.002
Linear-by-Linear Association	10.091	1	.001
N of Valid Cases	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .18.

## Use amphetamines: in car \* Gender

### Crosstab

			Gender			Total
			Male	Female	Unknown	
Use amphetamines: in car	No	Count	232	494	2	728
		Expected Count	231.7	494.3	2.0	728.0
		% within Use amphetamines: in car	31.9%	67.9%	.3%	100.0%
		% within Gender	99.6%	99.4%	100.0%	99.5%
		% of Total	31.7%	67.5%	.3%	99.5%
	Yes	Count	1	3	0	4
		Expected Count	1.3	2.7	.0	4.0
		% within Use amphetamines: in car	25.0%	75.0%	.0%	100.0%
		% within Gender	.4%	.6%	.0%	.5%
		% of Total	.1%	.4%	.0%	.5%
	Total	Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use amphetamines: in car	31.8%	67.9%	.3%	100.0%

% within Gender	100.0%	100.0%	100.0%	100.0%
% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.100(a)	2	.951
Likelihood Ratio	.114	2	.944
Linear-by-Linear Association	.078	1	.780
N of Valid Cases	732		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .01.

## Use amphetamines: private parties \* Gender

## Crosstab

		Gender			Total	
		Male	Female	Unknown		
Use amphetamines: private parties	No	Count	231	489	2	722
		Expected Count	229.8	490.2	2.0	722.0
		% within Use amphetamines: private parties	32.0%	67.7%	.3%	100.0%
		% within Gender	99.1%	98.4%	100.0%	98.6%
		% of Total	31.6%	66.8%	.3%	98.6%
	Yes	Count	2	8	0	10
		Expected Count	3.2	6.8	.0	10.0
		% within Use amphetamines: private parties	20.0%	80.0%	.0%	100.0%
		% within Gender	.9%	1.6%	.0%	1.4%
		% of Total	.3%	1.1%	.0%	1.4%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use amphetamines: private parties	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.692(a)	2	.707
Likelihood Ratio	.775	2	.679

Linear-by-Linear Association	.611	1	.435
N of Valid Cases	732		
a 3 cells (50.0%) have expected count less than 5. The minimum expected count is .03.			

## Use amphetamines: other \* Gender

		Crosstab			Total	
		Gender				
		Male	Female	Unknown		
Use amphetamines: other	No	Count	229	487	2	718
		Expected Count	228.5	487.5	2.0	718.0
		% within Use amphetamines: other	31.9%	67.8%	.3%	100.0%
		% within Gender	98.3%	98.0%	100.0%	98.1%
		% of Total	31.3%	66.5%	.3%	98.1%
	Yes	Count	4	10	0	14
		Expected Count	4.5	9.5	.0	14.0
		% within Use amphetamines: other	28.6%	71.4%	.0%	100.0%
		% within Gender	1.7%	2.0%	.0%	1.9%
		% of Total	.5%	1.4%	.0%	1.9%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use amphetamines: other	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests			
	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.113(a)	2	.945
Likelihood Ratio	.152	2	.927
Linear-by-Linear Association	.057	1	.811
N of Valid Cases	732		
a 3 cells (50.0%) have expected count less than 5. The minimum expected count is .04.			

## Use sedatives: never \* Gender

		Crosstab			Total	
		Gender				
		Male	Female	Unknown		

		Count	10	20	0	30
	No	Expected Count	9.5	20.4	.1	30.0
Use sedatives: never		% within Use sedatives: never	33.3%	66.7%	.0%	100.0%
		% within Gender	4.3%	4.0%	.0%	4.1%
		% of Total	1.4%	2.7%	.0%	4.1%
	Yes	Count	223	477	2	702
		Expected Count	223.5	476.6	1.9	702.0
		% within Use sedatives: never	31.8%	67.9%	.3%	100.0%
		% within Gender	95.7%	96.0%	100.0%	95.9%
		% of Total	30.5%	65.2%	.3%	95.9%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use sedatives: never	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.115(a)	2	.944
Likelihood Ratio	.196	2	.907
Linear-by-Linear Association	.044	1	.833
N of Valid Cases	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .08.

## Use sedatives: on campus \* Gender

## Crosstab

		Gender			Total	
		Male	Female	Unknown		
Use sedatives: on campus	No	Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use sedatives: on campus	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use sedatives: on campus	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value
Pearson Chi-Square	.(a)
N of Valid Cases	732

a No statistics are computed because Use sedatives: on campus is a constant.

## Use sedatives: res hall \* Gender

## Crosstab

		No	Gender			Total
			Male	Female	Unknown	
Use sedatives: res hall	No	Count	231	494	2	727
		Expected Count	231.4	493.6	2.0	727.0
		% within Use sedatives: res hall	31.8%	68.0%	.3%	100.0%
		% within Gender	99.1%	99.4%	100.0%	99.3%
	Yes	% of Total	31.6%	67.5%	.3%	99.3%
		Count	2	3	0	5
		Expected Count	1.6	3.4	.0	5.0
		% within Use sedatives: res hall	40.0%	60.0%	.0%	100.0%
Total	Yes	% within Gender	.9%	.6%	.0%	.7%
		% of Total	.3%	.4%	.0%	.7%
		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use sedatives: res hall	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.166(a)	2	.921
Likelihood Ratio	.173	2	.917
Linear-by-Linear Association	.162	1	.687
N of Valid Cases	732		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .01.

## Use sedatives: frat/sor \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Use sedatives: frat/sor	No	Count	233	496	2	731
		Expected Count	232.7	496.3	2.0	731.0
		% within Use sedatives: frat/sor	31.9%	67.9%	.3%	100.0%
		% within Gender	100.0%	99.8%	100.0%	99.9%
	Yes	% of Total	31.8%	67.8%	.3%	99.9%
		Count	0	1	0	1
		Expected Count	.3	.7	.0	1.0
		% within Use sedatives: frat/sor	.0%	100.0%	.0%	100.0%
Total	No	% within Gender	.0%	.2%	.0%	.1%
		% of Total	.0%	.1%	.0%	.1%
		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use sedatives: frat/sor	31.8%	67.9%	.3%	100.0%
	Yes	% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.473(a)	2	.789
Likelihood Ratio	.775	2	.679
Linear-by-Linear Association	.450	1	.502
N of Valid Cases	732		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .00.

## Use sedatives: bar/restaurant \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Use sedatives: bar/restaurant	No	Count	232	496	2	730
		Expected Count	232.4	495.6	2.0	730.0
		% within Use sedatives: bar/restaurant	31.8%	67.9%	.3%	100.0%
		% within Gender	99.6%	99.8%	100.0%	99.7%
	Yes	% of Total	31.7%	67.8%	.3%	99.7%
		Count	1	1	0	2
		Expected Count	.6	1.4	.0	2.0

		% within Use sedatives: bar/restaurant	50.0%	50.0%	.0%	100.0%
	Yes	% within Gender	.4%	.2%	.0%	.3%
		% of Total	.1%	.1%	.0%	.3%
		Count	233	497	2	732
Total		Expected Count	233.0	497.0	2.0	732.0
		% within Use sedatives: bar/restaurant	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.308(a)	2	.857
Likelihood Ratio	.292	2	.864
Linear-by-Linear Association	.308	1	.579
N of Valid Cases	732		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .01.

## Use sedatives: where live \* Gender

## Crosstab

		Gender			Total	
		Male	Female	Unknown		
Use sedatives: where live	No	Count	228	482	2	712
		Expected Count	226.6	483.4	1.9	712.0
		% within Use sedatives: where live	32.0%	67.7%	.3%	100.0%
		% within Gender	97.9%	97.0%	100.0%	97.3%
		% of Total	31.1%	65.8%	.3%	97.3%
	Yes	Count	5	15	0	20
		Expected Count	6.4	13.6	.1	20.0
		% within Use sedatives: where live	25.0%	75.0%	.0%	100.0%
		% within Gender	2.1%	3.0%	.0%	2.7%
		% of Total	.7%	2.0%	.0%	2.7%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use sedatives: where live	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.510(a)	2	.775
Likelihood Ratio	.584	2	.747
Linear-by-Linear Association	.399	1	.528
N of Valid Cases	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .05.

## Use sedatives: in car \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Use sedatives: in car	No	Count	232	496	2	730
		Expected Count	232.4	495.6	2.0	730.0
		% within Use sedatives: in car	31.8%	67.9%	.3%	100.0%
		% within Gender	99.6%	99.8%	100.0%	99.7%
		% of Total	31.7%	67.8%	.3%	99.7%
	Yes	Count	1	1	0	2
		Expected Count	.6	1.4	.0	2.0
		% within Use sedatives: in car	50.0%	50.0%	.0%	100.0%
		% within Gender	.4%	.2%	.0%	.3%
		% of Total	.1%	.1%	.0%	.3%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use sedatives: in car	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.308(a)	2	.857
Likelihood Ratio	.292	2	.864
Linear-by-Linear Association	.308	1	.579
N of Valid Cases	732		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .01.

## Use sedatives: private parties \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Use sedatives: private parties	No	Count	228	490	2	720
		Expected Count	229.2	488.9	2.0	720.0
		% within Use sedatives: private parties	31.7%	68.1%	.3%	100.0%
		% within Gender	97.9%	98.6%	100.0%	98.4%
		% of Total	31.1%	66.9%	.3%	98.4%
	Yes	Count	5	7	0	12
		Expected Count	3.8	8.1	.0	12.0
		% within Use sedatives: private parties	41.7%	58.3%	.0%	100.0%
		% within Gender	2.1%	1.4%	.0%	1.6%
		% of Total	.7%	1.0%	.0%	1.6%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use sedatives: private parties	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.568(a)	2	.753
Likelihood Ratio	.577	2	.749
Linear-by-Linear Association	.562	1	.453
N of Valid Cases	732		

a 3 cells (50.0%) have expected count less than 5. The minimum expected count is .03.

## Use sedatives: other \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Use sedatives: other	No	Count	230	495	2	727
		Expected Count	231.4	493.6	2.0	727.0
		% within Use sedatives: other	31.6%	68.1%	.3%	100.0%
		% within Gender	98.7%	99.6%	100.0%	99.3%
		% of Total	31.4%	67.6%	.3%	99.3%
	Yes	Count	3	1	0	4
		Expected Count	.75	.25	.0	1.0
		% within Use sedatives: other	75.0%	25.0%	.0%	100.0%
		% within Gender	75.0%	25.0%	.0%	100.0%
		% of Total	.7%	.2%	.0%	1.0%

Use sedatives: other	Yes	Count	3	2	0	5
		Expected Count	1.6	3.4	.0	5.0
		% within Use sedatives: other	60.0%	40.0%	.0%	100.0%
		% within Gender	1.3%	.4%	.0%	.7%
		% of Total	.4%	.3%	.0%	.7%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use sedatives: other	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1.846(a)	2	.397
Likelihood Ratio	1.700	2	.427
Linear-by-Linear Association	1.837	1	.175
N of Valid Cases	732		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .01.

## Use hallucinogens: never \* Gender

## Crosstab

		Gender			Total	
		Male	Female	Unknown		
Use hallucinogens: never	No	Count	20	49	0	69
		Expected Count	22.0	46.8	.2	69.0
		% within Use hallucinogens: never	29.0%	71.0%	.0%	100.0%
		% within Gender	8.6%	9.9%	.0%	9.4%
		% of Total	2.7%	6.7%	.0%	9.4%
	Yes	Count	213	448	2	663
		Expected Count	211.0	450.2	1.8	663.0
		% within Use hallucinogens: never	32.1%	67.6%	.3%	100.0%
		% within Gender	91.4%	90.1%	100.0%	90.6%
		% of Total	29.1%	61.2%	.3%	90.6%
	Total	Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use hallucinogens: never	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	.511(a)	2	.775
<b>Likelihood Ratio</b>	.703	2	.704
<b>Linear-by-Linear Association</b>	.227	1	.634
<b>N of Valid Cases</b>	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .19.

**Use hallucinogens: on campus \* Gender****Crosstab**

		Gender			Total
		Male	Female	Unknown	
Use hallucinogens: on campus	No	Count	230	497	2 729
		Expected Count	232.0	495.0	2.0 729.0
		% within Use hallucinogens: on campus	31.6%	68.2%	.3% 100.0%
		% within Gender	98.7%	100.0%	100.0% 99.6%
		% of Total	31.4%	67.9%	.3% 99.6%
	Yes	Count	3	0	0 3
		Expected Count	1.0	2.0	.0 3.0
		% within Use hallucinogens: on campus	100.0%	.0%	.0% 100.0%
		% within Gender	1.3%	.0%	.0% .4%
		% of Total	.4%	.0%	.0% .4%
Total		Count	233	497	2 732
		Expected Count	233.0	497.0	2.0 732.0
		% within Use hallucinogens: on campus	31.8%	67.9%	.3% 100.0%
		% within Gender	100.0%	100.0%	100.0% 100.0%
		% of Total	31.8%	67.9%	.3% 100.0%

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	6.451(a)	2	.040
<b>Likelihood Ratio</b>	6.895	2	.032
<b>Linear-by-Linear Association</b>	6.363	1	.012
<b>N of Valid Cases</b>	732		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .01.

# Use hallucinogens: res hall \* Gender

Crosstab

		Gender			Total	
		Male	Female	Unknown		
Use hallucinogens: res hall	No	Count	231	493	2	726
		Expected Count	231.1	492.9	2.0	726.0
		% within Use hallucinogens: res hall	31.8%	67.9%	.3%	100.0%
		% within Gender	99.1%	99.2%	100.0%	99.2%
		% of Total	31.6%	67.3%	.3%	99.2%
	Yes	Count	2	4	0	6
		Expected Count	1.9	4.1	.0	6.0
		% within Use hallucinogens: res hall	33.3%	66.7%	.0%	100.0%
		% within Gender	.9%	.8%	.0%	.8%
		% of Total	.3%	.5%	.0%	.8%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use hallucinogens: res hall	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.022(a)	2	.989
Likelihood Ratio	.039	2	.981
Linear-by-Linear Association	.009	1	.926
N of Valid Cases	732		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .02.

# Use hallucinogens: frat/sor \* Gender

Crosstab

		Gender			Total	
		Male	Female	Unknown		
Use hallucinogens: frat/sor	No	Count	230	496	2	728
		Expected Count	231.7	494.3	2.0	728.0
		% within Use hallucinogens: frat/sor	31.6%	68.1%	.3%	100.0%
		% within Gender	98.7%	99.8%	100.0%	99.5%
	Yes	Count	2	4	0	6
		Expected Count	1.9	4.1	.0	6.0
		% within Use hallucinogens: frat/sor	33.3%	66.7%	.0%	100.0%
		% within Gender	.9%	.8%	.0%	.8%
		% of Total	.3%	.5%	.0%	.8%

		% of Total	31.4%	67.8%	.3%	99.5%
Use hallucinogens: frat/sor	Yes	Count	3	1	0	4
		Expected Count	1.3	2.7	.0	4.0
		% within Use hallucinogens: frat/sor	75.0%	25.0%	.0%	100.0%
		% within Gender	1.3%	.2%	.0%	.5%
		% of Total	.4%	.1%	.0%	.5%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use hallucinogens: frat/sor	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	3.456(a)	2	.178
Likelihood Ratio	3.163	2	.206
Linear-by-Linear Association	3.423	1	.064
N of Valid Cases	732		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .01.

## Use hallucinogens: bar/restaurant \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
Use hallucinogens: bar/restaurant	No	Count	230	496	2	728
		Expected Count	231.7	494.3	2.0	728.0
		% within Use hallucinogens: bar/restaurant	31.6%	68.1%	.3%	100.0%
		% within Gender	98.7%	99.8%	100.0%	99.5%
		% of Total	31.4%	67.8%	.3%	99.5%
	Yes	Count	3	1	0	4
		Expected Count	1.3	2.7	.0	4.0
		% within Use hallucinogens: bar/restaurant	75.0%	25.0%	.0%	100.0%
		% within Gender	1.3%	.2%	.0%	.5%
		% of Total	.4%	.1%	.0%	.5%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use hallucinogens:	31.8%	67.9%	.3%	100.0%

	bar/restaurant				
	% within Gender	100.0%	100.0%	100.0%	100.0%
	% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	3.456(a)	2	.178
Likelihood Ratio	3.163	2	.206
Linear-by-Linear Association	3.423	1	.064
N of Valid Cases	732		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .01.

## Use hallucinogens: where live \* Gender

## Crosstab

		Gender			Total
		Male	Female	Unknown	
Use hallucinogens: where live	No	Count	222	480	2 704
		Expected Count	224.1	478.0	1.9 704.0
		% within Use hallucinogens: where live	31.5%	68.2%	.3% 100.0%
		% within Gender	95.3%	96.6%	100.0% 96.2%
		% of Total	30.3%	65.6%	.3% 96.2%
	Yes	Count	11	17	0 28
		Expected Count	8.9	19.0	.1 28.0
		% within Use hallucinogens: where live	39.3%	60.7%	.0% 100.0%
		% within Gender	4.7%	3.4%	.0% 3.8%
		% of Total	1.5%	2.3%	.0% 3.8%
Total		Count	233	497	2 732
		Expected Count	233.0	497.0	2.0 732.0
		% within Use hallucinogens: where live	31.8%	67.9%	.3% 100.0%
		% within Gender	100.0%	100.0%	100.0% 100.0%
		% of Total	31.8%	67.9%	.3% 100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.809(a)	2	.667
Likelihood Ratio	.859	2	.651
Linear-by-Linear Association	.784	1	.376

N of Valid Cases	732	
a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .08.		

## Use hallucinogens: in car \* Gender

Crosstab

		Gender			Total	
		Male	Female	Unknown		
Use hallucinogens: in car	No	Count	226	494	2	722
		Expected Count	229.8	490.2	2.0	722.0
		% within Use hallucinogens: in car	31.3%	68.4%	.3%	100.0%
		% within Gender	97.0%	99.4%	100.0%	98.6%
		% of Total	30.9%	67.5%	.3%	98.6%
	Yes	Count	7	3	0	10
		Expected Count	3.2	6.8	.0	10.0
		% within Use hallucinogens: in car	70.0%	30.0%	.0%	100.0%
		% within Gender	3.0%	.6%	.0%	1.4%
		% of Total	1.0%	.4%	.0%	1.4%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use hallucinogens: in car	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	6.813(a)	2	.033
Likelihood Ratio	6.226	2	.044
Linear-by-Linear Association	6.757	1	.009
N of Valid Cases	732		

a 3 cells (50.0%) have expected count less than 5. The minimum expected count is .03.

## Use hallucinogens: private parties \* Gender

Crosstab

		Gender			Total	
		Male	Female	Unknown		
		Count	221	471	2	694

Use hallucinogens: private parties	No	Expected Count	220.9	471.2	1.9	694.0
		% within Use hallucinogens: private parties	31.8%	67.9%	.3%	100.0%
		% within Gender	94.8%	94.8%	100.0%	94.8%
		% of Total	30.2%	64.3%	.3%	94.8%
	Yes	Count	12	26	0	38
		Expected Count	12.1	25.8	.1	38.0
		% within Use hallucinogens: private parties	31.6%	68.4%	.0%	100.0%
		% within Gender	5.2%	5.2%	.0%	5.2%
		% of Total	1.6%	3.6%	.0%	5.2%
	Total	Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use hallucinogens: private parties	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.112(a)	2	.946
Likelihood Ratio	.216	2	.898
Linear-by-Linear Association	.000	1	.998
N of Valid Cases	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .10.

## Use hallucinogens: other \* Gender

#### Crosstab

			Gender			Total
			Male	Female	Unknown	
Use hallucinogens: other	No	Count	225	484	2	711
		Expected Count	226.3	482.7	1.9	711.0
		% within Use hallucinogens: other	31.6%	68.1%	.3%	100.0%
		% within Gender	96.6%	97.4%	100.0%	97.1%
		% of Total	30.7%	66.1%	.3%	97.1%
	Yes	Count	8	13	0	21
		Expected Count	6.7	14.3	.1	21.0
		% within Use hallucinogens: other	38.1%	61.9%	.0%	100.0%
		% within Gender	3.4%	2.6%	.0%	2.9%

	% of Total	1.1%	1.8%	.0%	2.9%
Total	Count	233	497	2	732
	Expected Count	233.0	497.0	2.0	732.0
	% within Use hallucinogens: other	31.8%	67.9%	.3%	100.0%
	% within Gender	100.0%	100.0%	100.0%	100.0%
	% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.440(a)	2	.803
Likelihood Ratio	.485	2	.785
Linear-by-Linear Association	.417	1	.519
N of Valid Cases	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .06.

## Use opiates: never \* Gender

## Crosstab

		No	Gender			Total
			Male	Female	Unknown	
Use opiates: never	No	Count	9	15	0	24
		Expected Count	7.6	16.3	.1	24.0
		% within Use opiates: never	37.5%	62.5%	.0%	100.0%
		% within Gender	3.9%	3.0%	.0%	3.3%
		% of Total	1.2%	2.0%	.0%	3.3%
	Yes	Count	224	482	2	708
		Expected Count	225.4	480.7	1.9	708.0
		% within Use opiates: never	31.6%	68.1%	.3%	100.0%
		% within Gender	96.1%	97.0%	100.0%	96.7%
		% of Total	30.6%	65.8%	.3%	96.7%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use opiates: never	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.425(a)	2	.809

Likelihood Ratio	.480	2	.787
Linear-by-Linear Association	.395	1	.530
N of Valid Cases	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .07.

## Use opiates: on campus \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Use opiates: on campus	No	Count	232	497	2	731
		Expected Count	232.7	496.3	2.0	731.0
		% within Use opiates: on campus	31.7%	68.0%	.3%	100.0%
		% within Gender	99.6%	100.0%	100.0%	99.9%
	Yes	% of Total	31.7%	67.9%	.3%	99.9%
		Count	1	0	0	1
		Expected Count	.3	.7	.0	1.0
		% within Use opiates: on campus	100.0%	.0%	.0%	100.0%
Total	Yes	% within Gender	.4%	.0%	.0%	.1%
		% of Total	.1%	.0%	.0%	.1%
		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use opiates: on campus	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	2.145(a)	2	.342
Likelihood Ratio	2.292	2	.318
Linear-by-Linear Association	2.115	1	.146
N of Valid Cases	732		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .00.

## Use opiates: res hall \* Gender

Crosstab

	Gender		Total
	Male	Female	

			Male	Female	Unknown	
Use opiates: res hall	No	Count	232	496	2	730
		Expected Count	232.4	495.6	2.0	730.0
		% within Use opiates: res hall	31.8%	67.9%	.3%	100.0%
		% within Gender	99.6%	99.8%	100.0%	99.7%
		% of Total	31.7%	67.8%	.3%	99.7%
	Yes	Count	1	1	0	2
		Expected Count	.6	1.4	.0	2.0
		% within Use opiates: res hall	50.0%	50.0%	.0%	100.0%
		% within Gender	.4%	.2%	.0%	.3%
		% of Total	.1%	.1%	.0%	.3%
Total	Count		233	497	2	732
	Expected Count		233.0	497.0	2.0	732.0
	% within Use opiates: res hall		31.8%	67.9%	.3%	100.0%
	% within Gender		100.0%	100.0%	100.0%	100.0%
	% of Total		31.8%	67.9%	.3%	100.0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.308(a)	2	.857
Likelihood Ratio	.292	2	.864
Linear-by-Linear Association	.308	1	.579
N of Valid Cases	732		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .01.

## Use opiates: frat/sor \* Gender

#### Crosstab

			Gender			Total
			Male	Female	Unknown	
Use opiates: frat/sor	No	Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use opiates: frat/sor	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%
	Total	Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use opiates: frat/sor	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

	% of Total	31.8%	67.9%	.3%	100.0%
--	------------	-------	-------	-----	--------

## Chi-Square Tests

	Value
Pearson Chi-Square	.(a)
N of Valid Cases	732

a No statistics are computed because Use opiates: frat/sor is a constant.

## Use opiates: bar/restaurant \* Gender

## Crosstab

		Count	Gender			Total
			Male	Female	Unknown	
Use opiates: bar/restaurant	No	Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use opiates: bar/restaurant	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use opiates: bar/restaurant	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value
Pearson Chi-Square	.(a)
N of Valid Cases	732

a No statistics are computed because Use opiates: bar/restaurant is a constant.

## Use opiates: where live \* Gender

## Crosstab

		Count	Gender			Total
			Male	Female	Unknown	
	No	Count	228	492	2	722
		Expected Count	229.8	490.2	2.0	722.0
		% within Use opiates: where live	31.6%	68.1%	.3%	100.0%

		% within Gender	97.9%	99.0%	100.0%	98.6%
		% of Total	31.1%	67.2%	.3%	98.6%
Use opiates: where live	Yes	Count	5	5	0	10
		Expected Count	3.2	6.8	.0	10.0
		% within Use opiates: where live	50.0%	50.0%	.0%	100.0%
		% within Gender	2.1%	1.0%	.0%	1.4%
		% of Total	.7%	.7%	.0%	1.4%
		Count	233	497	2	732
Total		Expected Count	233.0	497.0	2.0	732.0
		% within Use opiates: where live	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1.557(a)	2	.459
Likelihood Ratio	1.478	2	.478
Linear-by-Linear Association	1.555	1	.212
N of Valid Cases	732		

a 3 cells (50.0%) have expected count less than 5. The minimum expected count is .03.

## Use opiates: in car \* Gender

### Crosstab

		Gender			Total	
		Male	Female	Unknown		
Use opiates: in car	No	Count	232	495	2	729
		Expected Count	232.0	495.0	2.0	729.0
		% within Use opiates: in car	31.8%	67.9%	.3%	100.0%
		% within Gender	99.6%	99.6%	100.0%	99.6%
		% of Total	31.7%	67.6%	.3%	99.6%
	Yes	Count	1	2	0	3
		Expected Count	1.0	2.0	.0	3.0
		% within Use opiates: in car	33.3%	66.7%	.0%	100.0%
		% within Gender	.4%	.4%	.0%	.4%
		% of Total	.1%	.3%	.0%	.4%
	Total	Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use opiates: in car	31.8%	67.9%	.3%	100.0%

% within Gender	100.0%	100.0%	100.0%	100.0%
% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.011(a)	2	.994
Likelihood Ratio	.019	2	.990
Linear-by-Linear Association	.004	1	.948
N of Valid Cases	732		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .01.

## Use opiates: private parties \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
Use opiates: private parties	No	Count	228	490	2	720
		Expected Count	229.2	488.9	2.0	720.0
		% within Use opiates: private parties	31.7%	68.1%	.3%	100.0%
	Yes	% within Gender	97.9%	98.6%	100.0%	98.4%
		% of Total	31.1%	66.9%	.3%	98.4%
		Count	5	7	0	12
		Expected Count	3.8	8.1	.0	12.0
		% within Use opiates: private parties	41.7%	58.3%	.0%	100.0%
		% within Gender	2.1%	1.4%	.0%	1.6%
		% of Total	.7%	1.0%	.0%	1.6%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use opiates: private parties	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.568(a)	2	.753
Likelihood Ratio	.577	2	.749
Linear-by-Linear Association	.562	1	.453
N of Valid Cases	732		

a 3 cells (50.0%) have expected count less than 5. The minimum expected count is .03.

# Use opiates: other \* Gender

Crosstab

		Gender			Total	
		Male	Female	Unknown		
Use opiates: other	No	Count	230	493	2	725
		Expected Count	230.8	492.2	2.0	725.0
		% within Use opiates: other	31.7%	68.0%	.3%	100.0%
		% within Gender	98.7%	99.2%	100.0%	99.0%
		% of Total	31.4%	67.3%	.3%	99.0%
	Yes	Count	3	4	0	7
		Expected Count	2.2	4.8	.0	7.0
		% within Use opiates: other	42.9%	57.1%	.0%	100.0%
		% within Gender	1.3%	.8%	.0%	1.0%
		% of Total	.4%	.5%	.0%	1.0%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use opiates: other	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.410(a)	2	.815
Likelihood Ratio	.409	2	.815
Linear-by-Linear Association	.407	1	.524
N of Valid Cases	732		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .02.

# Use inhalants: never \* Gender

Crosstab

		Gender			Total	
		Male	Female	Unknown		
Use inhalants: never	No	Count	9	20	0	29
		Expected Count	9.2	19.7	.1	29.0
		% within Use inhalants: never	31.0%	69.0%	.0%	100.0%
		% within Gender	3.9%	4.0%	.0%	4.0%
	Yes	Count	1	1	0	2
		Expected Count	1.0	1.0	.0	2.0
		% within Use inhalants: never	50.0%	50.0%	.0%	100.0%
		% within Gender	.5%	.5%	.0%	1.0%

		% of Total	1.2%	2.7%	.0%	4.0%
Use inhalants: never	Yes	Count	224	477	2	703
		Expected Count	223.8	477.3	1.9	703.0
		% within Use inhalants: never	31.9%	67.9%	.3%	100.0%
		% within Gender	96.1%	96.0%	100.0%	96.0%
		% of Total	30.6%	65.2%	.3%	96.0%
		Count	233	497	2	732
Total		Expected Count	233.0	497.0	2.0	732.0
		% within Use inhalants: never	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.094(a)	2	.954
Likelihood Ratio	.173	2	.917
Linear-by-Linear Association	.004	1	.951
N of Valid Cases	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .08.

## Use inhalants: on campus \* Gender

## Crosstab

		Gender			Total	
		Male	Female	Unknown		
Use inhalants: on campus	No	Count	232	497	2	731
		Expected Count	232.7	496.3	2.0	731.0
		% within Use inhalants: on campus	31.7%	68.0%	.3%	100.0%
		% within Gender	99.6%	100.0%	100.0%	99.9%
		% of Total	31.7%	67.9%	.3%	99.9%
	Yes	Count	1	0	0	1
		Expected Count	.3	.7	.0	1.0
		% within Use inhalants: on campus	100.0%	.0%	.0%	100.0%
		% within Gender	.4%	.0%	.0%	.1%
		% of Total	.1%	.0%	.0%	.1%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use inhalants: on campus	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%

	% of Total	31.8%	67.9%	.3%	100.0%
--	------------	-------	-------	-----	--------

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	2.145(a)	2	.342
<b>Likelihood Ratio</b>	2.292	2	.318
<b>Linear-by-Linear Association</b>	2.115	1	.146
<b>N of Valid Cases</b>	732		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .00.

## Use inhalants: res hall \* Gender

## Crosstab

		Gender			Total	
		Male	Female	Unknown		
Use inhalants: res hall	No	<b>Count</b>	231	494	2	727
		<b>Expected Count</b>	231.4	493.6	2.0	727.0
		<b>% within Use inhalants: res hall</b>	31.8%	68.0%	.3%	100.0%
		<b>% within Gender</b>	99.1%	99.4%	100.0%	99.3%
		<b>% of Total</b>	31.6%	67.5%	.3%	99.3%
	Yes	<b>Count</b>	2	3	0	5
		<b>Expected Count</b>	1.6	3.4	.0	5.0
		<b>% within Use inhalants: res hall</b>	40.0%	60.0%	.0%	100.0%
		<b>% within Gender</b>	.9%	.6%	.0%	.7%
		<b>% of Total</b>	.3%	.4%	.0%	.7%
Total		<b>Count</b>	233	497	2	732
		<b>Expected Count</b>	233.0	497.0	2.0	732.0
		<b>% within Use inhalants: res hall</b>	31.8%	67.9%	.3%	100.0%
		<b>% within Gender</b>	100.0%	100.0%	100.0%	100.0%
		<b>% of Total</b>	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	.166(a)	2	.921
<b>Likelihood Ratio</b>	.173	2	.917
<b>Linear-by-Linear Association</b>	.162	1	.687
<b>N of Valid Cases</b>	732		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .01.

# Use inhalants: frat/sor \* Gender

Crosstab

		Gender			Total	
		Male	Female	Unknown		
Use inhalants: frat/sor	No	Count	232	497	2	731
		Expected Count	232.7	496.3	2.0	731.0
		% within Use inhalants: frat/sor	31.7%	68.0%	.3%	100.0%
		% within Gender	99.6%	100.0%	100.0%	99.9%
		% of Total	31.7%	67.9%	.3%	99.9%
	Yes	Count	1	0	0	1
		Expected Count	.3	.7	.0	1.0
		% within Use inhalants: frat/sor	100.0%	.0%	.0%	100.0%
		% within Gender	.4%	.0%	.0%	.1%
		% of Total	.1%	.0%	.0%	.1%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use inhalants: frat/sor	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	2.145(a)	2	.342
Likelihood Ratio	2.292	2	.318
Linear-by-Linear Association	2.115	1	.146
N of Valid Cases	732		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .00.

# Use inhalants: bar/restaurant \* Gender

Crosstab

		Gender			Total	
		Male	Female	Unknown		
	No	Count	232	497	2	731
		Expected Count	232.7	496.3	2.0	731.0
		% within Use inhalants: bar/restaurant	31.7%	68.0%	.3%	100.0%

Use inhalants: bar/restaurant	Yes	% within Gender	99.6%	100.0%	100.0%	99.9%
		% of Total	31.7%	67.9%	.3%	99.9%
		Count	1	0	0	1
		Expected Count	.3	.7	.0	1.0
		% within Use inhalants: bar/restaurant	100.0%	.0%	.0%	100.0%
	No	% within Gender	.4%	.0%	.0%	.1%
		% of Total	.1%	.0%	.0%	.1%
	Total	Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use inhalants: bar/restaurant	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	2.145(a)	2	.342
Likelihood Ratio	2.292	2	.318
Linear-by-Linear Association	2.115	1	.146
N of Valid Cases	732		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .00.

## Use inhalants: where live \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
Use inhalants: where live	No	Count	229	491	2	722
		Expected Count	229.8	490.2	2.0	722.0
		% within Use inhalants: where live	31.7%	68.0%	.3%	100.0%
		% within Gender	98.3%	98.8%	100.0%	98.6%
		% of Total	31.3%	67.1%	.3%	98.6%
	Yes	Count	4	6	0	10
		Expected Count	3.2	6.8	.0	10.0
		% within Use inhalants: where live	40.0%	60.0%	.0%	100.0%
		% within Gender	1.7%	1.2%	.0%	1.4%
		% of Total	.5%	.8%	.0%	1.4%
		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0

Total	% within Use inhalants: where live	31.8%	67.9%	.3%	100.0%
	% within Gender	100.0%	100.0%	100.0%	100.0%
	% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.333(a)	2	.846
Likelihood Ratio	.349	2	.840
Linear-by-Linear Association	.326	1	.568
N of Valid Cases	732		

a 3 cells (50.0%) have expected count less than 5. The minimum expected count is .03.

## Use inhalants: in car \* Gender

## Crosstab

		Gender			Total	
		Male	Female	Unknown		
Use inhalants: in car	No	Count	231	497	2	730
		Expected Count	232.4	495.6	2.0	730.0
		% within Use inhalants: in car	31.6%	68.1%	.3%	100.0%
		% within Gender	99.1%	100.0%	100.0%	99.7%
		% of Total	31.6%	67.9%	.3%	99.7%
	Yes	Count	2	0	0	2
		Expected Count	.6	1.4	.0	2.0
		% within Use inhalants: in car	100.0%	.0%	.0%	100.0%
		% within Gender	.9%	.0%	.0%	.3%
		% of Total	.3%	.0%	.0%	.3%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use inhalants: in car	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	4.295(a)	2	.117
Likelihood Ratio	4.591	2	.101
Linear-by-Linear Association	4.236	1	.040
N of Valid Cases	732		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .01.

## Use inhalants: private parties \* Gender

Crosstab

		Gender			Total	
		Male	Female	Unknown		
Use inhalants: private parties	No	Count	228	491	2	721
		Expected Count	229.5	489.5	2.0	721.0
		% within Use inhalants: private parties	31.6%	68.1%	.3%	100.0%
		% within Gender	97.9%	98.8%	100.0%	98.5%
		% of Total	31.1%	67.1%	.3%	98.5%
	Yes	Count	5	6	0	11
		Expected Count	3.5	7.5	.0	11.0
		% within Use inhalants: private parties	45.5%	54.5%	.0%	100.0%
		% within Gender	2.1%	1.2%	.0%	1.5%
		% of Total	.7%	.8%	.0%	1.5%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use inhalants: private parties	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.975(a)	2	.614
Likelihood Ratio	.950	2	.622
Linear-by-Linear Association	.973	1	.324
N of Valid Cases	732		

a 3 cells (50.0%) have expected count less than 5. The minimum expected count is .03.

## Use inhalants: other \* Gender

Crosstab

	Gender			Total
	Male	Female	Unknown	

Use inhalants: other	No	Count	230	492	2	724
		Expected Count	230.5	491.6	2.0	724.0
		% within Use inhalants: other	31.8%	68.0%	.3%	100.0%
		% within Gender	98.7%	99.0%	100.0%	98.9%
		% of Total	31.4%	67.2%	.3%	98.9%
	Yes	Count	3	5	0	8
		Expected Count	2.5	5.4	.0	8.0
		% within Use inhalants: other	37.5%	62.5%	.0%	100.0%
		% within Gender	1.3%	1.0%	.0%	1.1%
		% of Total	.4%	.7%	.0%	1.1%
	Total		233	497	2	732
	Expected Count		233.0	497.0	2.0	732.0
	% within Use inhalants: other		31.8%	67.9%	.3%	100.0%
	% within Gender		100.0%	100.0%	100.0%	100.0%
	% of Total		31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.138(a)	2	.933
Likelihood Ratio	.157	2	.925
Linear-by-Linear Association	.129	1	.720
N of Valid Cases	732		

a 3 cells (50.0%) have expected count less than 5. The minimum expected count is .02.

## Use designer drugs: never \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
Use designer drugs: never	No	Count	16	43	0	59
		Expected Count	18.8	40.1	.2	59.0
		% within Use designer drugs: never	27.1%	72.9%	.0%	100.0%
		% within Gender	6.9%	8.7%	.0%	8.1%
		% of Total	2.2%	5.9%	.0%	8.1%
	Yes	Count	217	454	2	673
		Expected Count	214.2	456.9	1.8	673.0
		% within Use designer drugs: never	32.2%	67.5%	.3%	100.0%
		% within Gender	93.1%	91.3%	100.0%	91.9%
		% of Total	29.6%	62.0%	.3%	91.9%

Total	Count	233	497	2	732
	Expected Count	233.0	497.0	2.0	732.0
	% within Use designer drugs: never	31.8%	67.9%	.3%	100.0%
	% within Gender	100.0%	100.0%	100.0%	100.0%
	% of Total	31.8%	67.9%	.3%	100.0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.858(a)	2	.651
Likelihood Ratio	1.035	2	.596
Linear-by-Linear Association	.570	1	.450
N of Valid Cases	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .16.

## Use designer drugs: on campus \* Gender

#### Crosstab

		Gender			Total
		Male	Female	Unknown	
Use designer drugs: on campus	No	Count	233	497	2 732
		Expected Count	233.0	497.0	2.0 732.0
		% within Use designer drugs: on campus	31.8%	67.9%	.3% 100.0%
		% within Gender	100.0%	100.0%	100.0% 100.0%
		% of Total	31.8%	67.9%	.3% 100.0%
Total		Count	233	497	2 732
		Expected Count	233.0	497.0	2.0 732.0
		% within Use designer drugs: on campus	31.8%	67.9%	.3% 100.0%
		% within Gender	100.0%	100.0%	100.0% 100.0%
		% of Total	31.8%	67.9%	.3% 100.0%

#### Chi-Square Tests

	Value
Pearson Chi-Square	.(a)
N of Valid Cases	732

a No statistics are computed because Use designer drugs: on campus is a constant.

## Use designer drugs: res hall \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Use designer drugs: res hall	No	Count	233	490	2	725
		Expected Count	230.8	492.2	2.0	725.0
		% within Use designer drugs: res hall	32.1%	67.6%	.3%	100.0%
		% within Gender	100.0%	98.6%	100.0%	99.0%
		% of Total	31.8%	66.9%	.3%	99.0%
	Yes	Count	0	7	0	7
		Expected Count	2.2	4.8	.0	7.0
		% within Use designer drugs: res hall	.0%	100.0%	.0%	100.0%
		% within Gender	.0%	1.4%	.0%	1.0%
		% of Total	.0%	1.0%	.0%	1.0%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use designer drugs: res hall	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	3.342(a)	2	.188
Likelihood Ratio	5.453	2	.065
Linear-by-Linear Association	3.174	1	.075
N of Valid Cases	732		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .02.

## Use designer drugs: frat/sor \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Use designer drugs: frat/sor	No	Count	232	495	2	729
		Expected Count	232.0	495.0	2.0	729.0
		% within Use designer drugs: frat/sor	31.8%	67.9%	.3%	100.0%
		% within Gender	99.6%	99.6%	100.0%	99.6%
		% of Total	31.7%	67.6%	.3%	99.6%
		Count	1	2	0	3
		Expected Count	1.0	2.0	.0	3.0
		% within Use designer drugs: frat/sor	.3%	67.9%	.3%	99.6%
		% within Gender	99.6%	99.6%	100.0%	99.6%
		% of Total	.3%	67.6%	.3%	99.6%

		% within Use designer drugs: frat/sor	33.3%	66.7%	.0%	100.0%
	Yes	% within Gender	.4%	.4%	.0%	.4%
		% of Total	.1%	.3%	.0%	.4%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use designer drugs: frat/sor	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.011(a)	2	.994
Likelihood Ratio	.019	2	.990
Linear-by-Linear Association	.004	1	.948
N of Valid Cases	732		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .01.

## Use designer drugs: bar/restaurant \* Gender

#### Crosstab

			Gender			Total
			Male	Female	Unknown	
Use designer drugs: bar/restaurant	No	Count	230	492	2	724
		Expected Count	230.5	491.6	2.0	724.0
		% within Use designer drugs: bar/restaurant	31.8%	68.0%	.3%	100.0%
		% within Gender	98.7%	99.0%	100.0%	98.9%
		% of Total	31.4%	67.2%	.3%	98.9%
	Yes	Count	3	5	0	8
		Expected Count	2.5	5.4	.0	8.0
		% within Use designer drugs: bar/restaurant	37.5%	62.5%	.0%	100.0%
		% within Gender	1.3%	1.0%	.0%	1.1%
		% of Total	.4%	.7%	.0%	1.1%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use designer drugs: bar/restaurant	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%

	% of Total	31.8%	67.9%	.3%	100.0%
--	------------	-------	-------	-----	--------

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.138(a)	2	.933
Likelihood Ratio	.157	2	.925
Linear-by-Linear Association	.129	1	.720
N of Valid Cases	732		

a 3 cells (50.0%) have expected count less than 5. The minimum expected count is .02.

# Use designer drugs: where live \* Gender

## Crosstab

		Gender			Total
		Male	Female	Unknown	
Use designer drugs: where live	No	Count	227	482	2 711
		Expected Count	226.3	482.7	1.9 711.0
		% within Use designer drugs: where live	31.9%	67.8%	.3% 100.0%
		% within Gender	97.4%	97.0%	100.0% 97.1%
		% of Total	31.0%	65.8%	.3% 97.1%
	Yes	Count	6	15	0 21
		Expected Count	6.7	14.3	.1 21.0
		% within Use designer drugs: where live	28.6%	71.4%	.0% 100.0%
		% within Gender	2.6%	3.0%	.0% 2.9%
		% of Total	.8%	2.0%	.0% 2.9%
Total		Count	233	497	2 732
		Expected Count	233.0	497.0	2.0 732.0
		% within Use designer drugs: where live	31.8%	67.9%	.3% 100.0%
		% within Gender	100.0%	100.0%	100.0% 100.0%
		% of Total	31.8%	67.9%	.3% 100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.171(a)	2	.918
Likelihood Ratio	.230	2	.891
Linear-by-Linear Association	.087	1	.768
N of Valid Cases	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .06.

# Use designer drugs: in car \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Use designer drugs: in car	No	Count	231	491	2	724
		Expected Count	230.5	491.6	2.0	724.0
		% within Use designer drugs: in car	31.9%	67.8%	.3%	100.0%
		% within Gender	99.1%	98.8%	100.0%	98.9%
		% of Total	31.6%	67.1%	.3%	98.9%
	Yes	Count	2	6	0	8
		Expected Count	2.5	5.4	.0	8.0
		% within Use designer drugs: in car	25.0%	75.0%	.0%	100.0%
		% within Gender	.9%	1.2%	.0%	1.1%
		% of Total	.3%	.8%	.0%	1.1%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use designer drugs: in car	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.201(a)	2	.904
Likelihood Ratio	.230	2	.891
Linear-by-Linear Association	.157	1	.692
N of Valid Cases	732		

a 3 cells (50.0%) have expected count less than 5. The minimum expected count is .02.

# Use designer drugs: private parties \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
		Count	224	471	2	697
		Expected Count	221.9	473.2	1.9	697.0

Use designer drugs: private parties	No	% within Use designer drugs: private parties	32.1%	67.6%	.3%	100.0%	
		% within Gender	96.1%	94.8%	100.0%	95.2%	
		% of Total	30.6%	64.3%	.3%	95.2%	
	Yes	Count	9	26	0	35	
		Expected Count	11.1	23.8	.1	35.0	
		% within Use designer drugs: private parties	25.7%	74.3%	.0%	100.0%	
		% within Gender	3.9%	5.2%	.0%	4.8%	
		% of Total	1.2%	3.6%	.0%	4.8%	
			Count	233	497	2	732
			Expected Count	233.0	497.0	2.0	732.0
			% within Use designer drugs: private parties	31.8%	67.9%	.3%	100.0%
			% within Gender	100.0%	100.0%	100.0%	100.0%
			% of Total	31.8%	67.9%	.3%	100.0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.753(a)	2	.686
Likelihood Ratio	.872	2	.647
Linear-by-Linear Association	.566	1	.452
N of Valid Cases	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .10.

## Use designer drugs: other \* Gender

#### Crosstab

	No		Gender			Total
			Male	Female	Unknown	
Use designer drugs: other	No	Count	230	492	2	724
		Expected Count	230.5	491.6	2.0	724.0
		% within Use designer drugs: other	31.8%	68.0%	.3%	100.0%
		% within Gender	98.7%	99.0%	100.0%	98.9%
		% of Total	31.4%	67.2%	.3%	98.9%
	Yes	Count	3	5	0	8
		Expected Count	2.5	5.4	.0	8.0
		% within Use designer drugs: other	37.5%	62.5%	.0%	100.0%
		% within Gender	1.3%	1.0%	.0%	1.1%
		% of Total	.4%	.7%	.0%	1.1%

Total	Count	233	497	2	732
	Expected Count	233.0	497.0	2.0	732.0
	% within Use designer drugs: other	31.8%	67.9%	.3%	100.0%
	% within Gender	100.0%	100.0%	100.0%	100.0%
	% of Total	31.8%	67.9%	.3%	100.0%

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.138(a)	2	.933
Likelihood Ratio	.157	2	.925
Linear-by-Linear Association	.129	1	.720
N of Valid Cases	732		

a 3 cells (50.0%) have expected count less than 5. The minimum expected count is .02.

## Use steroids: never \* Gender

### Crosstab

		Gender			Total	
		Male	Female	Unknown		
Use steroids: never	No	Count	3	3	0	6
		Expected Count	1.9	4.1	.0	6.0
		% within Use steroids: never	50.0%	50.0%	.0%	100.0%
		% within Gender	1.3%	.6%	.0%	.8%
		% of Total	.4%	.4%	.0%	.8%
	Yes	Count	230	494	2	726
		Expected Count	231.1	492.9	2.0	726.0
		% within Use steroids: never	31.7%	68.0%	.3%	100.0%
		% within Gender	98.7%	99.4%	100.0%	99.2%
		% of Total	31.4%	67.5%	.3%	99.2%
	Total	Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use steroids: never	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.929(a)	2	.628
Likelihood Ratio	.881	2	.644

Linear-by-Linear Association	.928	1	.335
N of Valid Cases	732		
a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .02.			

## Use steroids: on campus \* Gender

			Crosstab			Total
			Male	Female	Unknown	
Use steroids: on campus	No	Count	232	497	2	731
		Expected Count	232.7	496.3	2.0	731.0
		% within Use steroids: on campus	31.7%	68.0%	.3%	100.0%
		% within Gender	99.6%	100.0%	100.0%	99.9%
		% of Total	31.7%	67.9%	.3%	99.9%
	Yes	Count	1	0	0	1
		Expected Count	.3	.7	.0	1.0
		% within Use steroids: on campus	100.0%	.0%	.0%	100.0%
		% within Gender	.4%	.0%	.0%	.1%
		% of Total	.1%	.0%	.0%	.1%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use steroids: on campus	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests			
	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	2.145(a)	2	.342
Likelihood Ratio	2.292	2	.318
Linear-by-Linear Association	2.115	1	.146
N of Valid Cases	732		
a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .00.			

## Use steroids: res hall \* Gender

			Crosstab			Total
			Male	Female	Unknown	

Use steroids: res hall	No	Count	233	496	2	731
		Expected Count	232.7	496.3	2.0	731.0
		% within Use steroids: res hall	31.9%	67.9%	.3%	100.0%
		% within Gender	100.0%	99.8%	100.0%	99.9%
		% of Total	31.8%	67.8%	.3%	99.9%
	Yes	Count	0	1	0	1
		Expected Count	.3	.7	.0	1.0
		% within Use steroids: res hall	.0%	100.0%	.0%	100.0%
		% within Gender	.0%	.2%	.0%	.1%
		% of Total	.0%	.1%	.0%	.1%
	Total		233	497	2	732
	Expected Count		233.0	497.0	2.0	732.0
	% within Use steroids: res hall		31.8%	67.9%	.3%	100.0%
	% within Gender		100.0%	100.0%	100.0%	100.0%
	% of Total		31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.473(a)	2	.789
Likelihood Ratio	.775	2	.679
Linear-by-Linear Association	.450	1	.502
N of Valid Cases	732		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .00.

## Use steroids: frat/sor \* Gender

## Crosstab

		Gender			Total	
		Male	Female	Unknown		
Use steroids: frat/sor	No	Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use steroids: frat/sor	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use steroids: frat/sor	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

**Chi-Square Tests**

	Value
Pearson Chi-Square	.(a)
N of Valid Cases	732

a No statistics are computed because Use steroids: frat/sor is a constant.

**Use steroids: bar/restaurant \* Gender****Crosstab**

		Count	Gender			Total
			Male	Female	Unknown	
Use steroids: bar/restaurant	No	Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use steroids: bar/restaurant	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use steroids: bar/restaurant	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

**Chi-Square Tests**

	Value
Pearson Chi-Square	.(a)
N of Valid Cases	732

a No statistics are computed because Use steroids: bar/restaurant is a constant.

**Use steroids: where live \* Gender****Crosstab**

		Count	Gender			Total
			Male	Female	Unknown	
Use steroids: where live	No	Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use steroids: where live	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

	% of Total	31.8%	67.9%	.3%	100.0%
Total	<b>Count</b>	233	497	2	732
	<b>Expected Count</b>	233.0	497.0	2.0	732.0
	<b>% within Use steroids: where live</b>	31.8%	67.9%	.3%	100.0%
	<b>% within Gender</b>	100.0%	100.0%	100.0%	100.0%
	<b>% of Total</b>	31.8%	67.9%	.3%	100.0%

**Chi-Square Tests**

	Value
<b>Pearson Chi-Square</b>	.(a)
<b>N of Valid Cases</b>	732

a No statistics are computed because Use steroids: where live is a constant.

**Use steroids: in car \* Gender****Crosstab**

		Gender			Total	
		Male	Female	Unknown		
Use steroids: in car	No	<b>Count</b>	233	497	2	732
		<b>Expected Count</b>	233.0	497.0	2.0	732.0
		<b>% within Use steroids: in car</b>	31.8%	67.9%	.3%	100.0%
		<b>% within Gender</b>	100.0%	100.0%	100.0%	100.0%
		<b>% of Total</b>	31.8%	67.9%	.3%	100.0%
Total		<b>Count</b>	233	497	2	732
		<b>Expected Count</b>	233.0	497.0	2.0	732.0
		<b>% within Use steroids: in car</b>	31.8%	67.9%	.3%	100.0%
		<b>% within Gender</b>	100.0%	100.0%	100.0%	100.0%
		<b>% of Total</b>	31.8%	67.9%	.3%	100.0%

**Chi-Square Tests**

	Value
<b>Pearson Chi-Square</b>	.(a)
<b>N of Valid Cases</b>	732

a No statistics are computed because Use steroids: in car is a constant.

**Use steroids: private parties \* Gender****Crosstab**


		Gender			Total
		Male	Female	Unknown	
Use steroids: private parties	No	Count	233	497	2 732
		Expected Count	233.0	497.0	2.0 732.0
		% within Use steroids: private parties	31.8%	67.9%	.3% 100.0%
		% within Gender	100.0%	100.0%	100.0% 100.0%
		% of Total	31.8%	67.9%	.3% 100.0%
Total		Count	233	497	2 732
		Expected Count	233.0	497.0	2.0 732.0
		% within Use steroids: private parties	31.8%	67.9%	.3% 100.0%
		% within Gender	100.0%	100.0%	100.0% 100.0%
		% of Total	31.8%	67.9%	.3% 100.0%

## Chi-Square Tests

	Value
Pearson Chi-Square	.(a)
N of Valid Cases	732

a No statistics are computed because Use steroids: private parties is a constant.

## Use steroids: other \* Gender

## Crosstab

		Gender			Total
		Male	Female	Unknown	
Use steroids: other	No	Count	233	497	2 732
		Expected Count	233.0	497.0	2.0 732.0
		% within Use steroids: other	31.8%	67.9%	.3% 100.0%
		% within Gender	100.0%	100.0%	100.0% 100.0%
		% of Total	31.8%	67.9%	.3% 100.0%
Total		Count	233	497	2 732
		Expected Count	233.0	497.0	2.0 732.0
		% within Use steroids: other	31.8%	67.9%	.3% 100.0%
		% within Gender	100.0%	100.0%	100.0% 100.0%
		% of Total	31.8%	67.9%	.3% 100.0%

## Chi-Square Tests

	Value
Pearson Chi-Square	.(a)
N of Valid Cases	732

a No statistics are computed because Use steroids: other is a constant.

## Use other drugs: never \* Gender

Crosstab

		No	Gender			Total
			Male	Female	Unknown	
Use other drugs: never	No	Count	8	21	0	29
		Expected Count	9.2	19.7	.1	29.0
		% within Use other drugs: never	27.6%	72.4%	.0%	100.0%
		% within Gender	3.4%	4.2%	.0%	4.0%
		% of Total	1.1%	2.9%	.0%	4.0%
	Yes	Count	225	476	2	703
		Expected Count	223.8	477.3	1.9	703.0
		% within Use other drugs: never	32.0%	67.7%	.3%	100.0%
		% within Gender	96.6%	95.8%	100.0%	96.0%
		% of Total	30.7%	65.0%	.3%	96.0%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use other drugs: never	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.344(a)	2	.842
Likelihood Ratio	.429	2	.807
Linear-by-Linear Association	.215	1	.643
N of Valid Cases	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .08.

## Use other drugs: on campus \* Gender

Crosstab

		Count	Gender			Total
			Male	Female	Unknown	
		Expected Count	233.0	497.0	2.0	732.0
		233	497	2	732	

Use other drugs: on campus	No	% within Use other drugs: on campus	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use other drugs: on campus	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value
Pearson Chi-Square	.(a)
N of Valid Cases	732

a No statistics are computed because Use other drugs: on campus is a constant.

## Use other drugs: res hall \* Gender

## Crosstab

		No	Gender			Total
			Male	Female	Unknown	
Use other drugs: res hall	No	Count	233	494	2	729
		Expected Count	232.0	495.0	2.0	729.0
		% within Use other drugs: res hall	32.0%	67.8%	.3%	100.0%
		% within Gender	100.0%	99.4%	100.0%	99.6%
		% of Total	31.8%	67.5%	.3%	99.6%
	Yes	Count	0	3	0	3
		Expected Count	1.0	2.0	.0	3.0
		% within Use other drugs: res hall	.0%	100.0%	.0%	100.0%
		% within Gender	.0%	.6%	.0%	.4%
		% of Total	.0%	.4%	.0%	.4%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use other drugs: res hall	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1.424(a)	2	.491

Likelihood Ratio	2.329	2	.312
Linear-by-Linear Association	1.353	1	.245
N of Valid Cases	732		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .01.

## Use other drugs: frat/sor \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Use other drugs: frat/sor	No	Count	232	497	2	731
		Expected Count	232.7	496.3	2.0	731.0
		% within Use other drugs: frat/sor	31.7%	68.0%	.3%	100.0%
		% within Gender	99.6%	100.0%	100.0%	99.9%
	Yes	% of Total	31.7%	67.9%	.3%	99.9%
		Count	1	0	0	1
		Expected Count	.3	.7	.0	1.0
		% within Use other drugs: frat/sor	100.0%	.0%	.0%	100.0%
Total	No	% within Gender	.4%	.0%	.0%	.1%
		% of Total	.1%	.0%	.0%	.1%
		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use other drugs: frat/sor	31.8%	67.9%	.3%	100.0%
	Yes	% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	2.145(a)	2	.342
Likelihood Ratio	2.292	2	.318
Linear-by-Linear Association	2.115	1	.146
N of Valid Cases	732		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .00.

## Use other drugs: bar/restaurant \* Gender

Crosstab

			Gender		Total
			Male	Female	

			Male	Female	Unknown	
Use other drugs: bar/restaurant	No	Count	232	497	2	731
		Expected Count	232.7	496.3	2.0	731.0
		% within Use other drugs: bar/restaurant	31.7%	68.0%	.3%	100.0%
		% within Gender	99.6%	100.0%	100.0%	99.9%
		% of Total	31.7%	67.9%	.3%	99.9%
	Yes	Count	1	0	0	1
		Expected Count	.3	.7	.0	1.0
		% within Use other drugs: bar/restaurant	100.0%	.0%	.0%	100.0%
		% within Gender	.4%	.0%	.0%	.1%
		% of Total	.1%	.0%	.0%	.1%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use other drugs: bar/restaurant	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	2.145(a)	2	.342
Likelihood Ratio	2.292	2	.318
Linear-by-Linear Association	2.115	1	.146
N of Valid Cases	732		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .00.

## Use other drugs: where live \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
Use other drugs: where live	No	Count	227	487	2	716
		Expected Count	227.9	486.1	2.0	716.0
		% within Use other drugs: where live	31.7%	68.0%	.3%	100.0%
		% within Gender	97.4%	98.0%	100.0%	97.8%
		% of Total	31.0%	66.5%	.3%	97.8%
		Count	6	10	0	16
		Expected Count	5.1	10.9	.0	16.0
		% within Use other drugs: where live	37.5%	62.5%	0.0%	100.0%
		% within Gender	30.0%	50.0%	20.0%	100.0%
		% of Total	37.5%	62.5%	0.0%	100.0%

		% within Use other drugs: where live	37.5%	62.5%	.0%	100.0%
	Yes	% within Gender	2.6%	2.0%	.0%	2.2%
		% of Total	.8%	1.4%	.0%	2.2%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use other drugs: where live	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.280(a)	2	.869
Likelihood Ratio	.317	2	.854
Linear-by-Linear Association	.260	1	.610
N of Valid Cases	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .04.

## Use other drugs: in car \* Gender

### Crosstab

			Gender			Total
			Male	Female	Unknown	
Use other drugs: in car	No	Count	232	494	2	728
		Expected Count	231.7	494.3	2.0	728.0
		% within Use other drugs: in car	31.9%	67.9%	.3%	100.0%
		% within Gender	99.6%	99.4%	100.0%	99.5%
		% of Total	31.7%	67.5%	.3%	99.5%
	Yes	Count	1	3	0	4
		Expected Count	1.3	2.7	.0	4.0
		% within Use other drugs: in car	25.0%	75.0%	.0%	100.0%
		% within Gender	.4%	.6%	.0%	.5%
		% of Total	.1%	.4%	.0%	.5%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use other drugs: in car	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

### Chi-Square Tests

--	--	--	--

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	.100(a)	2	.951
<b>Likelihood Ratio</b>	.114	2	.944
<b>Linear-by-Linear Association</b>	.078	1	.780
<b>N of Valid Cases</b>	732		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .01.

## Use other drugs: private parties \* Gender

Crosstab

		Gender			Total	
		Male	Female	Unknown		
Use other drugs: private parties	No	Count	228	486	2	716
		Expected Count	227.9	486.1	2.0	716.0
		% within Use other drugs: private parties	31.8%	67.9%	.3%	100.0%
		% within Gender	97.9%	97.8%	100.0%	97.8%
		% of Total	31.1%	66.4%	.3%	97.8%
	Yes	Count	5	11	0	16
		Expected Count	5.1	10.9	.0	16.0
		% within Use other drugs: private parties	31.3%	68.8%	.0%	100.0%
		% within Gender	2.1%	2.2%	.0%	2.2%
		% of Total	.7%	1.5%	.0%	2.2%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use other drugs: private parties	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	.048(a)	2	.976
<b>Likelihood Ratio</b>	.092	2	.955
<b>Linear-by-Linear Association</b>	.001	1	.979
<b>N of Valid Cases</b>	732		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .04.

## Use other drugs: other \* Gender

## Crosstab

		Gender			Total	
		Male	Female	Unknown		
Use other drugs: other	No	Count	227	491	2	720
		Expected Count	229.2	488.9	2.0	720.0
		% within Use other drugs: other	31.5%	68.2%	.3%	100.0%
		% within Gender	97.4%	98.8%	100.0%	98.4%
		% of Total	31.0%	67.1%	.3%	98.4%
	Yes	Count	6	6	0	12
		Expected Count	3.8	8.1	.0	12.0
		% within Use other drugs: other	50.0%	50.0%	.0%	100.0%
		% within Gender	2.6%	1.2%	.0%	1.6%
		% of Total	.8%	.8%	.0%	1.6%
Total		Count	233	497	2	732
		Expected Count	233.0	497.0	2.0	732.0
		% within Use other drugs: other	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1.874(a)	2	.392
Likelihood Ratio	1.778	2	.411
Linear-by-Linear Association	1.871	1	.171
N of Valid Cases	732		

a 3 cells (50.0%) have expected count less than 5. The minimum expected count is .03.

## Crosstabs

## Notes

Output Created		16-JUN-2004 09:46:59
Comments		
Input	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	747
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.

Syntax	CROSSTABS /TABLES= q21a q21b q21c q21d q21e q21f q21g q21h q21i q21j q21k q21l q21m q21n q21o q21p q21q q21r q21s q22a q22b q22c q22d q22e q22f q22g q22h q22i q22j q22k q23 q24a q24b q24c q24d q24e q24f q24g q24h q24i q25a1 q25a2 q25a3 q25a4 q25a5 q25a6 BY q05 /STATISTIC=CHISQ /CELLS= COUNT EXPECTED ROW COLUMN TOTAL .
Resources	Dimensions Requested
	Cells Available
	Elapsed Time

## Case Processing Summary

		Cases					
		Valid		Missing		Total	
		N	Percent	N	Percent	N	Percent
Hangover * Gender		731	97.9%	16	2.1%	747	100.0%
Poor test score * Gender		728	97.5%	19	2.5%	747	100.0%
Trouble w/ police, etc * Gender		729	97.6%	18	2.4%	747	100.0%
Damaged prop, fire alarm * Gender		729	97.6%	18	2.4%	747	100.0%
Argument or fight * Gender		728	97.5%	19	2.5%	747	100.0%
Nauseated or vomited * Gender		724	96.9%	23	3.1%	747	100.0%
Driven under influence * Gender		726	97.2%	21	2.8%	747	100.0%
Missed class * Gender		730	97.7%	17	2.3%	747	100.0%
Been criticized * Gender		730	97.7%	17	2.3%	747	100.0%
Thought I had a problem * Gender		729	97.6%	18	2.4%	747	100.0%
Had a memory loss * Gender		724	96.9%	23	3.1%	747	100.0%
Later regretted action * Gender		728	97.5%	19	2.5%	747	100.0%
Arrested for DWI/DUI * Gender		726	97.2%	21	2.8%	747	100.0%
HAVE BEEN TAKEN ADVANTAGE SEXUALLY * Gender		724	96.9%	23	3.1%	747	100.0%
HAVE TAKEN ADVANTAGE OF SOMEONE SEXUALLY * Gender		725	97.1%	22	2.9%	747	100.0%
TRIED/FAILED TO STOP * Gender		730	97.7%	17	2.3%	747	100.0%
THOUGHT ABOUT SUICIDE * Gender		727	97.3%	20	2.7%	747	100.0%
TRIED TO COMMIT SUICIDE * Gender		728	97.5%	19	2.5%	747	100.0%
BEEN HURT/INJURED * Gender		730	97.7%	17	2.3%	747	100.0%
A/D problems: mother * Gender		720	96.4%	27	3.6%	747	100.0%
A/D problems: father * Gender		720	96.4%	27	3.6%	747	100.0%
A/D problems: stepmother * Gender		720	96.4%	27	3.6%	747	100.0%
A/D problems: stepfather * Gender		720	96.4%	27	3.6%	747	100.0%
A/D problems: br/sister * Gender		720	96.4%	27	3.6%	747	100.0%
A/D problems: mother's parents * Gender		720	96.4%	27	3.6%	747	100.0%
A/D problems: father's parents * Gender		720	96.4%	27	3.6%	747	100.0%

A/D problems: aunts/uncles * Gender	720	96.4%	27	3.6%	747	100.0%
A/D problems: spouse * Gender	720	96.4%	27	3.6%	747	100.0%
A/D problems: children * Gender	720	96.4%	27	3.6%	747	100.0%
A/D problems: none * Gender	720	96.4%	27	3.6%	747	100.0%
VOLUNTEERING * Gender	656	87.8%	91	12.2%	747	100.0%
PARTICIPATED-INTERCOLLEGIATE ATHLETICS * Gender	726	97.2%	21	2.8%	747	100.0%
PARTICIPATED-INTRAMURAL OR CLUB SPORTS * Gender	726	97.2%	21	2.8%	747	100.0%
PARTICIPATED-SOCIAL FRATERNITIES OR SOR * Gender	724	96.9%	23	3.1%	747	100.0%
PARTICIPATED-RELIGIOUS GROUPS * Gender	726	97.2%	21	2.8%	747	100.0%
PARTICIPATED-INTERNATIONAL AND LANGUAGE GPS * Gender	726	97.2%	21	2.8%	747	100.0%
PARTICIPATED-MINORITY AND ETHNIC GROUPS * Gender	726	97.2%	21	2.8%	747	100.0%
PARTICIPATED-POLITICAL AND SOCIAL ACTION GPS * Gender	725	97.1%	22	2.9%	747	100.0%
PARTICIPATED-MUSIC,PERFORMING ARTS GROUPS * Gender	726	97.2%	21	2.8%	747	100.0%
PARTICIPATED-NEWSPAPER, RADIO, TV * Gender	723	96.8%	24	3.2%	747	100.0%
HAPPENED - ETHNIC HARASSMENT * Gender	728	97.5%	19	2.5%	747	100.0%
HAPPENED - THREATS OF PHYSICAL VIOLENCE * Gender	725	97.1%	22	2.9%	747	100.0%
HAPPENED - ACTUAL PHYSICAL VIOLENCE * Gender	727	97.3%	20	2.7%	747	100.0%
HAPPENED -THEFT INVOLVING FORCE * Gender	727	97.3%	20	2.7%	747	100.0%
HAPPENED -FORCED SEXUAL TOUCHING * Gender	729	97.6%	18	2.4%	747	100.0%
HAPPENED - UNWANTED SEXUAL INTERCOURSE * Gender	727	97.3%	20	2.7%	747	100.0%

## Hangover \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Hangover	Never	Count	50	126	0	176
		Expected Count	55.9	119.7	.5	176.0
		% within Hangover	28.4%	71.6%	.0%	100.0%
		% within Gender	21.6%	25.4%	.0%	24.1%
		% of Total	6.8%	17.2%	.0%	24.1%
	Once	Count	26	76	0	102
		Expected Count	32.4	69.3	.3	102.0
		% within Hangover	25.5%	74.5%	.0%	100.0%
		% within Gender	11.2%	15.3%	.0%	14.0%
		% of Total	3.6%	10.4%	.0%	14.0%
	Twice	Count	17	55	0	72
		Expected Count	22.9	49.0	.2	72.0
		% within Hangover	23.6%	76.4%	.0%	100.0%

		% within Gender	7.3%	11.1%	.0%	9.8%
		% of Total	2.3%	7.5%	.0%	9.8%
Hangover	3-5 times	Count	52	99	0	151
		Expected Count	47.9	102.7	.4	151.0
		% within Hangover	34.4%	65.6%	.0%	100.0%
		% within Gender	22.4%	19.9%	.0%	20.7%
		% of Total	7.1%	13.5%	.0%	20.7%
	6-9 times	Count	31	47	1	79
		Expected Count	25.1	53.7	.2	79.0
		% within Hangover	39.2%	59.5%	1.3%	100.0%
		% within Gender	13.4%	9.5%	50.0%	10.8%
		% of Total	4.2%	6.4%	.1%	10.8%
	10+ times	Count	56	94	1	151
		Expected Count	47.9	102.7	.4	151.0
		% within Hangover	37.1%	62.3%	.7%	100.0%
		% within Gender	24.1%	18.9%	50.0%	20.7%
		% of Total	7.7%	12.9%	.1%	20.7%
Total		Count	232	497	2	731
		Expected Count	232.0	497.0	2.0	731.0
		% within Hangover	31.7%	68.0%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.7%	68.0%	.3%	100.0%

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	14.944(a)	10	.134
Likelihood Ratio	14.817	10	.139
Linear-by-Linear Association	5.281	1	.022
N of Valid Cases	731		

a 6 cells (33.3%) have expected count less than 5. The minimum expected count is .20.

## Poor test score \* Gender

### Crosstab

			Gender			Total
			Male	Female	Unknown	
Never	Count	157	341	2	500	
		159.3	339.3	1.4	500.0	
		31.4%	68.2%	.4%	100.0%	

		% within Gender	67.7%	69.0%	100.0%	68.7%
		% of Total	21.6%	46.8%	.3%	68.7%
Poor test score	Once	Count	32	79	0	111
		Expected Count	35.4	75.3	.3	111.0
		% within Poor test score	28.8%	71.2%	.0%	100.0%
		% within Gender	13.8%	16.0%	.0%	15.2%
		% of Total	4.4%	10.9%	.0%	15.2%
	Twice	Count	17	38	0	55
		Expected Count	17.5	37.3	.2	55.0
		% within Poor test score	30.9%	69.1%	.0%	100.0%
		% within Gender	7.3%	7.7%	.0%	7.6%
		% of Total	2.3%	5.2%	.0%	7.6%
	3-5 times	Count	19	27	0	46
		Expected Count	14.7	31.2	.1	46.0
		% within Poor test score	41.3%	58.7%	.0%	100.0%
		% within Gender	8.2%	5.5%	.0%	6.3%
		% of Total	2.6%	3.7%	.0%	6.3%
	6-9 times	Count	6	6	0	12
		Expected Count	3.8	8.1	.0	12.0
		% within Poor test score	50.0%	50.0%	.0%	100.0%
		% within Gender	2.6%	1.2%	.0%	1.6%
		% of Total	.8%	.8%	.0%	1.6%
	10+ times	Count	1	3	0	4
		Expected Count	1.3	2.7	.0	4.0
		% within Poor test score	25.0%	75.0%	.0%	100.0%
		% within Gender	.4%	.6%	.0%	.5%
		% of Total	.1%	.4%	.0%	.5%
Total		Count	232	494	2	728
		Expected Count	232.0	494.0	2.0	728.0
		% within Poor test score	31.9%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.9%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	5.230(a)	10	.875
Likelihood Ratio	5.622	10	.846
Linear-by-Linear Association	1.674	1	.196
N of Valid Cases	728		

a 9 cells (50.0%) have expected count less than 5. The minimum expected count is .01.

# Trouble w/ police, etc \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Trouble w/ police, etc	Never	Count	170	407	2	579
		Expected Count	184.3	393.1	1.6	579.0
		% within Trouble w/ police, etc	29.4%	70.3%	.3%	100.0%
		% within Gender	73.3%	82.2%	100.0%	79.4%
		% of Total	23.3%	55.8%	.3%	79.4%
	Once	Count	44	64	0	108
		Expected Count	34.4	73.3	.3	108.0
		% within Trouble w/ police, etc	40.7%	59.3%	.0%	100.0%
		% within Gender	19.0%	12.9%	.0%	14.8%
		% of Total	6.0%	8.8%	.0%	14.8%
	Twice	Count	12	17	0	29
		Expected Count	9.2	19.7	.1	29.0
		% within Trouble w/ police, etc	41.4%	58.6%	.0%	100.0%
		% within Gender	5.2%	3.4%	.0%	4.0%
		% of Total	1.6%	2.3%	.0%	4.0%
	3-5 times	Count	3	7	0	10
		Expected Count	3.2	6.8	.0	10.0
		% within Trouble w/ police, etc	30.0%	70.0%	.0%	100.0%
		% within Gender	1.3%	1.4%	.0%	1.4%
		% of Total	.4%	1.0%	.0%	1.4%
	6-9 times	Count	1	0	0	1
		Expected Count	.3	.7	.0	1.0
		% within Trouble w/ police, etc	100.0%	.0%	.0%	100.0%
		% within Gender	.4%	.0%	.0%	.1%
		% of Total	.1%	.0%	.0%	.1%
	10+ times	Count	2	0	0	2
		Expected Count	.6	1.4	.0	2.0
		% within Trouble w/ police, etc	100.0%	.0%	.0%	100.0%
		% within Gender	.9%	.0%	.0%	.3%
		% of Total	.3%	.0%	.0%	.3%
Total		Count	232	495	2	729
		Expected Count	232.0	495.0	2.0	729.0
		% within Trouble w/ police, etc	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%

	% of Total	31.8%	67.9%	.3%	100.0%
--	------------	-------	-------	-----	--------

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	13.631(a)	10	.190
Likelihood Ratio	14.271	10	.161
Linear-by-Linear Association	8.642	1	.003
N of Valid Cases	729		

a 11 cells (61.1%) have expected count less than 5. The minimum expected count is .00.

## Damaged prop, fire alarm \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
Damaged prop, fire alarm	Never	Count	183	475	2	660
		Expected Count	209.1	449.1	1.8	660.0
		% within Damaged prop, fire alarm	27.7%	72.0%	.3%	100.0%
		% within Gender	79.2%	95.8%	100.0%	90.5%
		% of Total	25.1%	65.2%	.3%	90.5%
	Once	Count	21	14	0	35
		Expected Count	11.1	23.8	.1	35.0
		% within Damaged prop, fire alarm	60.0%	40.0%	.0%	100.0%
		% within Gender	9.1%	2.8%	.0%	4.8%
		% of Total	2.9%	1.9%	.0%	4.8%
	Twice	Count	11	3	0	14
		Expected Count	4.4	9.5	.0	14.0
		% within Damaged prop, fire alarm	78.6%	21.4%	.0%	100.0%
		% within Gender	4.8%	.6%	.0%	1.9%
		% of Total	1.5%	.4%	.0%	1.9%
	3-5 times	Count	13	4	0	17
		Expected Count	5.4	11.6	.0	17.0
		% within Damaged prop, fire alarm	76.5%	23.5%	.0%	100.0%
		% within Gender	5.6%	.8%	.0%	2.3%
		% of Total	1.8%	.5%	.0%	2.3%
	10+ times	Count	3	0	0	3
		Expected Count	1.0	2.0	.0	3.0
		% within Damaged prop, fire alarm	100.0%	.0%	.0%	100.0%
		% within Gender	1.3%	.0%	.0%	.4%

		<b>% of Total</b>	.4%	.0%	.0%	.4%
<b>Total</b>	<b>Count</b>	231	496	2	729	
	<b>Expected Count</b>	231.0	496.0	2.0	729.0	
	<b>% within Damaged prop, fire alarm</b>	31.7%	68.0%	.3%	100.0%	
	<b>% within Gender</b>	100.0%	100.0%	100.0%	100.0%	
	<b>% of Total</b>	31.7%	68.0%	.3%	100.0%	

**Chi-Square Tests**

	<b>Value</b>	<b>df</b>	<b>Asymp. Sig. (2-sided)</b>
<b>Pearson Chi-Square</b>	54.223(a)	8	.000
<b>Likelihood Ratio</b>	51.195	8	.000
<b>Linear-by-Linear Association</b>	48.678	1	.000
<b>N of Valid Cases</b>	729		
a 8 cells (53.3%) have expected count less than 5. The minimum expected count is .01.			

**Argument or fight \* Gender****Crosstab**

			<b>Gender</b>			<b>Total</b>
			<b>Male</b>	<b>Female</b>	<b>Unknown</b>	
<b>Argument or fight</b>	<b>Never</b>	<b>Count</b>	120	263	2	385
		<b>Expected Count</b>	122.7	261.3	1.1	385.0
		<b>% within Argument or fight</b>	31.2%	68.3%	.5%	100.0%
		<b>% within Gender</b>	51.7%	53.2%	100.0%	52.9%
		<b>% of Total</b>	16.5%	36.1%	.3%	52.9%
	<b>Once</b>	<b>Count</b>	39	81	0	120
		<b>Expected Count</b>	38.2	81.4	.3	120.0
		<b>% within Argument or fight</b>	32.5%	67.5%	.0%	100.0%
		<b>% within Gender</b>	16.8%	16.4%	.0%	16.5%
		<b>% of Total</b>	5.4%	11.1%	.0%	16.5%
	<b>Twice</b>	<b>Count</b>	30	62	0	92
		<b>Expected Count</b>	29.3	62.4	.3	92.0
		<b>% within Argument or fight</b>	32.6%	67.4%	.0%	100.0%
		<b>% within Gender</b>	12.9%	12.6%	.0%	12.6%
		<b>% of Total</b>	4.1%	8.5%	.0%	12.6%
	<b>3-5 times</b>	<b>Count</b>	25	59	0	84
		<b>Expected Count</b>	26.8	57.0	.2	84.0
		<b>% within Argument or fight</b>	29.8%	70.2%	.0%	100.0%
		<b>% within Gender</b>	10.8%	11.9%	.0%	11.5%

		% of Total	3.4%	8.1%	.0%	11.5%
6-9 times	Count	5	16	0	21	
	Expected Count	6.7	14.3	.1	21.0	
	% within Argument or fight	23.8%	76.2%	.0%	100.0%	
	% within Gender	2.2%	3.2%	.0%	2.9%	
	% of Total	.7%	2.2%	.0%	2.9%	
10+ times	Count	13	13	0	26	
	Expected Count	8.3	17.6	.1	26.0	
	% within Argument or fight	50.0%	50.0%	.0%	100.0%	
	% within Gender	5.6%	2.6%	.0%	3.6%	
	% of Total	1.8%	1.8%	.0%	3.6%	
Total	Count	232	494	2	728	
	Expected Count	232.0	494.0	2.0	728.0	
	% within Argument or fight	31.9%	67.9%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	31.9%	67.9%	.3%	100.0%	

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	6.623(a)	10	.761
Likelihood Ratio	7.148	10	.711
Linear-by-Linear Association	.788	1	.375
N of Valid Cases	728		

a 6 cells (33.3%) have expected count less than 5. The minimum expected count is .06.

## Nauseated or vomited \* Gender

#### Crosstab

			Gender			Total
			Male	Female	Unknown	
Never	Never	Count	65	151	0	216
		Expected Count	68.3	147.1	.6	216.0
		% within Nauseated or vomited	30.1%	69.9%	.0%	100.0%
		% within Gender	28.4%	30.6%	.0%	29.8%
		% of Total	9.0%	20.9%	.0%	29.8%
Once	Once	Count	42	114	0	156
		Expected Count	49.3	106.2	.4	156.0
		% within Nauseated or vomited	26.9%	73.1%	.0%	100.0%
		% within Gender	18.3%	23.1%	.0%	21.5%

		% of Total	5.8%	15.7%	.0%	21.5%
Nauseated or vomited	Twice	Count	44	73	0	117
		Expected Count	37.0	79.7	.3	117.0
		% within Nauseated or vomited	37.6%	62.4%	.0%	100.0%
		% within Gender	19.2%	14.8%	.0%	16.2%
		% of Total	6.1%	10.1%	.0%	16.2%
	3-5 times	Count	48	91	1	140
		Expected Count	44.3	95.3	.4	140.0
		% within Nauseated or vomited	34.3%	65.0%	.7%	100.0%
		% within Gender	21.0%	18.5%	50.0%	19.3%
		% of Total	6.6%	12.6%	.1%	19.3%
	6-9 times	Count	14	34	0	48
		Expected Count	15.2	32.7	.1	48.0
		% within Nauseated or vomited	29.2%	70.8%	.0%	100.0%
		% within Gender	6.1%	6.9%	.0%	6.6%
		% of Total	1.9%	4.7%	.0%	6.6%
	10+ times	Count	16	30	1	47
		Expected Count	14.9	32.0	.1	47.0
		% within Nauseated or vomited	34.0%	63.8%	2.1%	100.0%
		% within Gender	7.0%	6.1%	50.0%	6.5%
		% of Total	2.2%	4.1%	.1%	6.5%
Total		Count	229	493	2	724
		Expected Count	229.0	493.0	2.0	724.0
		% within Nauseated or vomited	31.6%	68.1%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.6%	68.1%	.3%	100.0%

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	12.961(a)	10	.226
Likelihood Ratio	10.640	10	.386
Linear-by-Linear Association	.529	1	.467
N of Valid Cases	724		

a 6 cells (33.3%) have expected count less than 5. The minimum expected count is .13.

## Driven under influence \* Gender

### Crosstab

	Gender		Total

			Male	Female	Unknown		
Driven under influence	Never	Count	143	347	2	492	
		Expected Count	157.2	333.4	1.4	492.0	
		% within Driven under influence	29.1%	70.5%	.4%	100.0%	
		% within Gender	61.6%	70.5%	100.0%	67.8%	
		% of Total	19.7%	47.8%	.3%	67.8%	
	Once	Count	22	49	0	71	
		Expected Count	22.7	48.1	.2	71.0	
		% within Driven under influence	31.0%	69.0%	.0%	100.0%	
		% within Gender	9.5%	10.0%	.0%	9.8%	
		% of Total	3.0%	6.7%	.0%	9.8%	
	Twice	Count	15	21	0	36	
		Expected Count	11.5	24.4	.1	36.0	
		% within Driven under influence	41.7%	58.3%	.0%	100.0%	
		% within Gender	6.5%	4.3%	.0%	5.0%	
		% of Total	2.1%	2.9%	.0%	5.0%	
	3-5 times	Count	23	42	0	65	
		Expected Count	20.8	44.0	.2	65.0	
		% within Driven under influence	35.4%	64.6%	.0%	100.0%	
		% within Gender	9.9%	8.5%	.0%	9.0%	
		% of Total	3.2%	5.8%	.0%	9.0%	
	6-9 times	Count	11	10	0	21	
		Expected Count	6.7	14.2	.1	21.0	
		% within Driven under influence	52.4%	47.6%	.0%	100.0%	
		% within Gender	4.7%	2.0%	.0%	2.9%	
		% of Total	1.5%	1.4%	.0%	2.9%	
	10+ times	Count	18	23	0	41	
		Expected Count	13.1	27.8	.1	41.0	
		% within Driven under influence	43.9%	56.1%	.0%	100.0%	
		% within Gender	7.8%	4.7%	.0%	5.6%	
		% of Total	2.5%	3.2%	.0%	5.6%	
Total			Count	232	492	2	726
			Expected Count	232.0	492.0	2.0	726.0
			% within Driven under influence	32.0%	67.8%	.3%	100.0%
			% within Gender	100.0%	100.0%	100.0%	100.0%
			% of Total	32.0%	67.8%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	11.353(a)	10	.331

<b>Likelihood Ratio</b>	11.472	10	.322
<b>Linear-by-Linear Association</b>	8.826	1	.003
<b>N of Valid Cases</b>	726		

a 6 cells (33.3%) have expected count less than 5. The minimum expected count is .06.

## Missed class \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Missed class	Never	Count	121	290	1	412
		Expected Count	130.9	279.9	1.1	412.0
		% within Missed class	29.4%	70.4%	.2%	100.0%
		% within Gender	52.2%	58.5%	50.0%	56.4%
		% of Total	16.6%	39.7%	.1%	56.4%
	Once	Count	21	53	0	74
		Expected Count	23.5	50.3	.2	74.0
		% within Missed class	28.4%	71.6%	.0%	100.0%
		% within Gender	9.1%	10.7%	.0%	10.1%
		% of Total	2.9%	7.3%	.0%	10.1%
	Twice	Count	23	46	0	69
		Expected Count	21.9	46.9	.2	69.0
		% within Missed class	33.3%	66.7%	.0%	100.0%
		% within Gender	9.9%	9.3%	.0%	9.5%
		% of Total	3.2%	6.3%	.0%	9.5%
	3-5 times	Count	30	63	0	93
		Expected Count	29.6	63.2	.3	93.0
		% within Missed class	32.3%	67.7%	.0%	100.0%
		% within Gender	12.9%	12.7%	.0%	12.7%
		% of Total	4.1%	8.6%	.0%	12.7%
	6-9 times	Count	14	22	0	36
		Expected Count	11.4	24.5	.1	36.0
		% within Missed class	38.9%	61.1%	.0%	100.0%
		% within Gender	6.0%	4.4%	.0%	4.9%
		% of Total	1.9%	3.0%	.0%	4.9%
	10+ times	Count	23	22	1	46
		Expected Count	14.6	31.3	.1	46.0
		% within Missed class	50.0%	47.8%	2.2%	100.0%
		% within Gender	9.9%	4.4%	50.0%	6.3%

		<b>% of Total</b>	3.2%	3.0%	.1%	6.3%
<b>Total</b>	<b>Count</b>	232	496	2	730	
	<b>Expected Count</b>	232.0	496.0	2.0	730.0	
	<b>% within Missed class</b>	31.8%	67.9%	.3%	100.0%	
	<b>% within Gender</b>	100.0%	100.0%	100.0%	100.0%	
	<b>% of Total</b>	31.8%	67.9%	.3%	100.0%	

**Chi-Square Tests**

	<b>Value</b>	<b>df</b>	<b>Asymp. Sig. (2-sided)</b>
<b>Pearson Chi-Square</b>	16.794(a)	10	.079
<b>Likelihood Ratio</b>	13.465	10	.199
<b>Linear-by-Linear Association</b>	6.088	1	.014
<b>N of Valid Cases</b>	730		

a 6 cells (33.3%) have expected count less than 5. The minimum expected count is .10.

**Been criticized \* Gender****Crosstab**

			<b>Gender</b>			<b>Total</b>
			<b>Male</b>	<b>Female</b>	<b>Unknown</b>	
<b>Been criticized</b>	<b>Never</b>	<b>Count</b>	145	301	1	447
		<b>Expected Count</b>	142.1	303.7	1.2	447.0
		<b>% within Been criticized</b>	32.4%	67.3%	.2%	100.0%
		<b>% within Gender</b>	62.5%	60.7%	50.0%	61.2%
		<b>% of Total</b>	19.9%	41.2%	.1%	61.2%
	<b>Once</b>	<b>Count</b>	21	74	0	95
		<b>Expected Count</b>	30.2	64.5	.3	95.0
		<b>% within Been criticized</b>	22.1%	77.9%	.0%	100.0%
		<b>% within Gender</b>	9.1%	14.9%	.0%	13.0%
		<b>% of Total</b>	2.9%	10.1%	.0%	13.0%
	<b>Twice</b>	<b>Count</b>	23	50	0	73
		<b>Expected Count</b>	23.2	49.6	.2	73.0
		<b>% within Been criticized</b>	31.5%	68.5%	.0%	100.0%
		<b>% within Gender</b>	9.9%	10.1%	.0%	10.0%
		<b>% of Total</b>	3.2%	6.8%	.0%	10.0%
	<b>3-5 times</b>	<b>Count</b>	23	51	1	75
		<b>Expected Count</b>	23.8	51.0	.2	75.0
		<b>% within Been criticized</b>	30.7%	68.0%	1.3%	100.0%
		<b>% within Gender</b>	9.9%	10.3%	50.0%	10.3%

		% of Total	3.2%	7.0%	.1%	10.3%
6-9 times	Count	10	13	0	23	
	Expected Count	7.3	15.6	.1	23.0	
	% within Been criticized	43.5%	56.5%	.0%	100.0%	
	% within Gender	4.3%	2.6%	.0%	3.2%	
	% of Total	1.4%	1.8%	.0%	3.2%	
10+ times	Count	10	7	0	17	
	Expected Count	5.4	11.6	.0	17.0	
	% within Been criticized	58.8%	41.2%	.0%	100.0%	
	% within Gender	4.3%	1.4%	.0%	2.3%	
	% of Total	1.4%	1.0%	.0%	2.3%	
Total	Count	232	496	2	730	
	Expected Count	232.0	496.0	2.0	730.0	
	% within Been criticized	31.8%	67.9%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	31.8%	67.9%	.3%	100.0%	

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	15.122(a)	10	.128
Likelihood Ratio	13.900	10	.178
Linear-by-Linear Association	1.459	1	.227
N of Valid Cases	730		

a 6 cells (33.3%) have expected count less than 5. The minimum expected count is .05.

## Thought I had a problem \* Gender

#### Crosstab

			Gender			Total
			Male	Female	Unknown	
Never	Never	Count	201	452	1	654
		Expected Count	208.1	444.1	1.8	654.0
		% within Thought I had a problem	30.7%	69.1%	.2%	100.0%
		% within Gender	86.6%	91.3%	50.0%	89.7%
		% of Total	27.6%	62.0%	.1%	89.7%
Once	Once	Count	7	20	0	27
		Expected Count	8.6	18.3	.1	27.0
		% within Thought I had a problem	25.9%	74.1%	.0%	100.0%
		% within Gender	3.0%	4.0%	.0%	3.7%

		% of Total	1.0%	2.7%	.0%	3.7%
Thought I had a problem	Twice	Count	9	8	0	17
	Expected Count	5.4	11.5	.0	17.0	
	% within Thought I had a problem	52.9%	47.1%	.0%	100.0%	
	% within Gender	3.9%	1.6%	.0%	2.3%	
	% of Total	1.2%	1.1%	.0%	2.3%	
	3-5 times	Count	9	10	0	19
	Expected Count	6.0	12.9	.1	19.0	
	% within Thought I had a problem	47.4%	52.6%	.0%	100.0%	
	% within Gender	3.9%	2.0%	.0%	2.6%	
	% of Total	1.2%	1.4%	.0%	2.6%	
	6-9 times	Count	2	0	0	2
	Expected Count	.6	1.4	.0	2.0	
	% within Thought I had a problem	100.0%	.0%	.0%	100.0%	
	% within Gender	.9%	.0%	.0%	.3%	
	% of Total	.3%	.0%	.0%	.3%	
	10+ times	Count	4	5	1	10
	Expected Count	3.2	6.8	.0	10.0	
	% within Thought I had a problem	40.0%	50.0%	10.0%	100.0%	
	% within Gender	1.7%	1.0%	50.0%	1.4%	
	% of Total	.5%	.7%	.1%	1.4%	
Total		Count	232	495	2	729
		Expected Count	232.0	495.0	2.0	729.0
		% within Thought I had a problem	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	46.365(a)	10	.000
Likelihood Ratio	17.317	10	.068
Linear-by-Linear Association	3.382	1	.066
N of Valid Cases	729		

a 9 cells (50.0%) have expected count less than 5. The minimum expected count is .01.

## Had a memory loss \* Gender

### Crosstab

	Gender		Total

			Male	Female	Unknown	
Had a memory loss	Never	Count	120	286	1	407
		Expected Count	129.3	276.6	1.1	407.0
		% within Had a memory loss	29.5%	70.3%	.2%	100.0%
		% within Gender	52.2%	58.1%	50.0%	56.2%
		% of Total	16.6%	39.5%	.1%	56.2%
	Once	Count	28	55	0	83
		Expected Count	26.4	56.4	.2	83.0
		% within Had a memory loss	33.7%	66.3%	.0%	100.0%
		% within Gender	12.2%	11.2%	.0%	11.5%
		% of Total	3.9%	7.6%	.0%	11.5%
	Twice	Count	35	56	0	91
		Expected Count	28.9	61.8	.3	91.0
		% within Had a memory loss	38.5%	61.5%	.0%	100.0%
		% within Gender	15.2%	11.4%	.0%	12.6%
		% of Total	4.8%	7.7%	.0%	12.6%
	3-5 times	Count	22	50	0	72
		Expected Count	22.9	48.9	.2	72.0
		% within Had a memory loss	30.6%	69.4%	.0%	100.0%
		% within Gender	9.6%	10.2%	.0%	9.9%
		% of Total	3.0%	6.9%	.0%	9.9%
	6-9 times	Count	7	20	0	27
		Expected Count	8.6	18.3	.1	27.0
		% within Had a memory loss	25.9%	74.1%	.0%	100.0%
		% within Gender	3.0%	4.1%	.0%	3.7%
		% of Total	1.0%	2.8%	.0%	3.7%
	10+ times	Count	18	25	1	44
		Expected Count	14.0	29.9	.1	44.0
		% within Had a memory loss	40.9%	56.8%	2.3%	100.0%
		% within Gender	7.8%	5.1%	50.0%	6.1%
		% of Total	2.5%	3.5%	.1%	6.1%
Total	Count	230	492	2	724	
	Expected Count	230.0	492.0	2.0	724.0	
	% within Had a memory loss	31.8%	68.0%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	31.8%	68.0%	.3%	100.0%	

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	12.533(a)	10	.251

Likelihood Ratio	9.311	10	.503
Linear-by-Linear Association	1.295	1	.255
N of Valid Cases	724		

a 6 cells (33.3%) have expected count less than 5. The minimum expected count is .07.

## Later regretted action \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
Later regretted action	Never	Count	108	262	1	371
		Expected Count	117.7	252.3	1.0	371.0
		% within Later regretted action	29.1%	70.6%	.3%	100.0%
		% within Gender	46.8%	52.9%	50.0%	51.0%
		% of Total	14.8%	36.0%	.1%	51.0%
	Once	Count	33	80	0	113
		Expected Count	35.9	76.8	.3	113.0
		% within Later regretted action	29.2%	70.8%	.0%	100.0%
		% within Gender	14.3%	16.2%	.0%	15.5%
		% of Total	4.5%	11.0%	.0%	15.5%
	Twice	Count	33	69	0	102
		Expected Count	32.4	69.4	.3	102.0
		% within Later regretted action	32.4%	67.6%	.0%	100.0%
		% within Gender	14.3%	13.9%	.0%	14.0%
		% of Total	4.5%	9.5%	.0%	14.0%
	3-5 times	Count	35	54	1	90
		Expected Count	28.6	61.2	.2	90.0
		% within Later regretted action	38.9%	60.0%	1.1%	100.0%
		% within Gender	15.2%	10.9%	50.0%	12.4%
		% of Total	4.8%	7.4%	.1%	12.4%
	6-9 times	Count	11	20	0	31
		Expected Count	9.8	21.1	.1	31.0
		% within Later regretted action	35.5%	64.5%	.0%	100.0%
		% within Gender	4.8%	4.0%	.0%	4.3%
		% of Total	1.5%	2.7%	.0%	4.3%
	10+ times	Count	11	10	0	21
		Expected Count	6.7	14.3	.1	21.0
		% within Later regretted action	52.4%	47.6%	.0%	100.0%
		% within Gender	4.8%	2.0%	.0%	2.9%

		% of Total	1.5%	1.4%	.0%	2.9%
Total		Count	231	495	2	728
		Expected Count	231.0	495.0	2.0	728.0
		% within Later regretted action	31.7%	68.0%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.7%	68.0%	.3%	100.0%

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	11.173(a)	10	.344
Likelihood Ratio	10.543	10	.394
Linear-by-Linear Association	5.896	1	.015
N of Valid Cases	728		
a 6 cells (33.3%) have expected count less than 5. The minimum expected count is .06.			

## Arrested for DWI/DUI \* Gender

### Crosstab

			Gender			Total
			Male	Female	Unknown	
Arrested for DWI/DUI	Never	Count	228	490	2	720
		Expected Count	230.1	487.9	2.0	720.0
		% within Arrested for DWI/DUI	31.7%	68.1%	.3%	100.0%
		% within Gender	98.3%	99.6%	100.0%	99.2%
		% of Total	31.4%	67.5%	.3%	99.2%
	Once	Count	4	1	0	5
		Expected Count	1.6	3.4	.0	5.0
		% within Arrested for DWI/DUI	80.0%	20.0%	.0%	100.0%
		% within Gender	1.7%	.2%	.0%	.7%
		% of Total	.6%	.1%	.0%	.7%
	Twice	Count	0	1	0	1
		Expected Count	.3	.7	.0	1.0
		% within Arrested for DWI/DUI	.0%	100.0%	.0%	100.0%
		% within Gender	.0%	.2%	.0%	.1%
		% of Total	.0%	.1%	.0%	.1%
Total		Count	232	492	2	726
		Expected Count	232.0	492.0	2.0	726.0
		% within Arrested for DWI/DUI	32.0%	67.8%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%

	% of Total	32.0%	67.8%	.3%	100.0%
--	------------	-------	-------	-----	--------

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	5.812(a)	4	.214
Likelihood Ratio	5.707	4	.222
Linear-by-Linear Association	1.600	1	.206
N of Valid Cases	726		

a 7 cells (77.8%) have expected count less than 5. The minimum expected count is .00.

# HAVE BEEN TAKEN ADVANTAGE SEXUALLY \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
HAVE BEEN TAKEN ADVANTAGE SEXUALLY	Never	Count	212	423	2	637
		Expected Count	203.2	432.0	1.8	637.0
		% within HAVE BEEN TAKEN ADVANTAGE SEXUALLY	33.3%	66.4%	.3%	100.0%
		% within Gender	91.8%	86.2%	100.0%	88.0%
		% of Total	29.3%	58.4%	.3%	88.0%
	Once	Count	6	39	0	45
		Expected Count	14.4	30.5	.1	45.0
		% within HAVE BEEN TAKEN ADVANTAGE SEXUALLY	13.3%	86.7%	.0%	100.0%
		% within Gender	2.6%	7.9%	.0%	6.2%
		% of Total	.8%	5.4%	.0%	6.2%
	Twice	Count	6	23	0	29
		Expected Count	9.3	19.7	.1	29.0
		% within HAVE BEEN TAKEN ADVANTAGE SEXUALLY	20.7%	79.3%	.0%	100.0%
		% within Gender	2.6%	4.7%	.0%	4.0%
		% of Total	.8%	3.2%	.0%	4.0%
	3-5 times	Count	4	4	0	8
		Expected Count	2.6	5.4	.0	8.0
		% within HAVE BEEN TAKEN ADVANTAGE SEXUALLY	50.0%	50.0%	.0%	100.0%
		% within Gender	1.7%	.8%	.0%	1.1%
		% of Total	.6%	.6%	.0%	1.1%

		Count	1	1	0	2
		Expected Count	.6	1.4	.0	2.0
	6-9 times	% within HAVE BEEN TAKEN ADVANTAGE SEXUALLY	50.0%	50.0%	.0%	100.0%
		% within Gender	.4%	.2%	.0%	.3%
		% of Total	.1%	.1%	.0%	.3%
		Count	2	1	0	3
		Expected Count	1.0	2.0	.0	3.0
	10+ times	% within HAVE BEEN TAKEN ADVANTAGE SEXUALLY	66.7%	33.3%	.0%	100.0%
		% within Gender	.9%	.2%	.0%	.4%
		% of Total	.3%	.1%	.0%	.4%
Total		Count	231	491	2	724
		Expected Count	231.0	491.0	2.0	724.0
		% within HAVE BEEN TAKEN ADVANTAGE SEXUALLY	31.9%	67.8%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.9%	67.8%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	12.925(a)	10	.228
Likelihood Ratio	14.226	10	.163
Linear-by-Linear Association	.173	1	.677
N of Valid Cases	724		

a 11 cells (61.1%) have expected count less than 5. The minimum expected count is .01.

## HAVE TAKEN ADVANTAGE OF SOMEONE SEXUALLY \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
Never		Count	219	479	2	700
		Expected Count	222.1	476.0	1.9	700.0
		% within HAVE TAKEN ADVANTAGE OF SOMEONE SEXUALLY	31.3%	68.4%	.3%	100.0%
		% within Gender	95.2%	97.2%	100.0%	96.6%
		% of Total	30.2%	66.1%	.3%	96.6%

HAVE TAKEN ADVANTAGE OF SOMEONE SEXUALLY	Once	Count	7	8	0	15
		Expected Count	4.8	10.2	.0	15.0
		% within HAVE TAKEN ADVANTAGE OF SOMEONE SEXUALLY	46.7%	53.3%	.0%	100.0%
		% within Gender	3.0%	1.6%	.0%	2.1%
		% of Total	1.0%	1.1%	.0%	2.1%
	Twice	Count	1	4	0	5
		Expected Count	1.6	3.4	.0	5.0
		% within HAVE TAKEN ADVANTAGE OF SOMEONE SEXUALLY	20.0%	80.0%	.0%	100.0%
		% within Gender	.4%	.8%	.0%	.7%
		% of Total	.1%	.6%	.0%	.7%
	3-5 times	Count	3	1	0	4
		Expected Count	1.3	2.7	.0	4.0
		% within HAVE TAKEN ADVANTAGE OF SOMEONE SEXUALLY	75.0%	25.0%	.0%	100.0%
		% within Gender	1.3%	.2%	.0%	.6%
		% of Total	.4%	.1%	.0%	.6%
	10+ times	Count	0	1	0	1
		Expected Count	.3	.7	.0	1.0
		% within HAVE TAKEN ADVANTAGE OF SOMEONE SEXUALLY	.0%	100.0%	.0%	100.0%
		% within Gender	.0%	.2%	.0%	.1%
		% of Total	.0%	.1%	.0%	.1%
	Total	Count	230	493	2	725
		Expected Count	230.0	493.0	2.0	725.0
		% within HAVE TAKEN ADVANTAGE OF SOMEONE SEXUALLY	31.7%	68.0%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.7%	68.0%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	5.902(a)	8	.658
Likelihood Ratio	5.890	8	.660
Linear-by-Linear Association	1.108	1	.292
N of Valid Cases	725		

a 12 cells (80.0%) have expected count less than 5. The minimum expected count is .00.

# TRIED/FAILED TO STOP \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
TRIED/FAILED TO STOP	Never	Count	211	470	2	683
		Expected Count	217.1	464.1	1.9	683.0
		% within TRIED/FAILED TO STOP	30.9%	68.8%	.3%	100.0%
		% within Gender	90.9%	94.8%	100.0%	93.6%
		% of Total	28.9%	64.4%	.3%	93.6%
	Once	Count	12	12	0	24
		Expected Count	7.6	16.3	.1	24.0
		% within TRIED/FAILED TO STOP	50.0%	50.0%	.0%	100.0%
		% within Gender	5.2%	2.4%	.0%	3.3%
		% of Total	1.6%	1.6%	.0%	3.3%
	Twice	Count	2	9	0	11
		Expected Count	3.5	7.5	.0	11.0
		% within TRIED/FAILED TO STOP	18.2%	81.8%	.0%	100.0%
		% within Gender	.9%	1.8%	.0%	1.5%
		% of Total	.3%	1.2%	.0%	1.5%
	3-5 times	Count	4	5	0	9
		Expected Count	2.9	6.1	.0	9.0
		% within TRIED/FAILED TO STOP	44.4%	55.6%	.0%	100.0%
		% within Gender	1.7%	1.0%	.0%	1.2%
		% of Total	.5%	.7%	.0%	1.2%
	6-9 times	Count	3	0	0	3
		Expected Count	1.0	2.0	.0	3.0
		% within TRIED/FAILED TO STOP	100.0%	.0%	.0%	100.0%
		% within Gender	1.3%	.0%	.0%	.4%
		% of Total	.4%	.0%	.0%	.4%
Total		Count	232	496	2	730
		Expected Count	232.0	496.0	2.0	730.0
		% within TRIED/FAILED TO STOP	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	12.068(a)	8	.148

Likelihood Ratio	12.429	8	.133
Linear-by-Linear Association	4.215	1	.040
N of Valid Cases	730		

a 9 cells (60.0%) have expected count less than 5. The minimum expected count is .01.

# THOUGHT ABOUT SUICIDE \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
THOUGHT ABOUT SUICIDE	Never	Count	214	472	2	688
		Expected Count	219.6	466.6	1.9	688.0
		% within THOUGHT ABOUT SUICIDE	31.1%	68.6%	.3%	100.0%
		% within Gender	92.2%	95.7%	100.0%	94.6%
		% of Total	29.4%	64.9%	.3%	94.6%
	Once	Count	5	16	0	21
		Expected Count	6.7	14.2	.1	21.0
		% within THOUGHT ABOUT SUICIDE	23.8%	76.2%	.0%	100.0%
		% within Gender	2.2%	3.2%	.0%	2.9%
		% of Total	.7%	2.2%	.0%	2.9%
	Twice	Count	6	2	0	8
		Expected Count	2.6	5.4	.0	8.0
		% within THOUGHT ABOUT SUICIDE	75.0%	25.0%	.0%	100.0%
		% within Gender	2.6%	.4%	.0%	1.1%
		% of Total	.8%	.3%	.0%	1.1%
	3-5 times	Count	4	1	0	5
		Expected Count	1.6	3.4	.0	5.0
		% within THOUGHT ABOUT SUICIDE	80.0%	20.0%	.0%	100.0%
		% within Gender	1.7%	.2%	.0%	.7%
		% of Total	.6%	.1%	.0%	.7%
	6-9 times	Count	1	1	0	2
		Expected Count	.6	1.4	.0	2.0
		% within THOUGHT ABOUT SUICIDE	50.0%	50.0%	.0%	100.0%
		% within Gender	.4%	.2%	.0%	.3%
		% of Total	.1%	.1%	.0%	.3%
		Count	2	1	0	3

		10+ times	Expected Count	1.0	2.0	.0	3.0
			% within THOUGHT ABOUT SUICIDE	66.7%	33.3%	.0%	100.0%
			% within Gender	.9%	.2%	.0%	.4%
			% of Total	.3%	.1%	.0%	.4%
			Count	232	493	2	727
			Expected Count	232.0	493.0	2.0	727.0
	Total		% within THOUGHT ABOUT SUICIDE	31.9%	67.8%	.3%	100.0%
			% within Gender	100.0%	100.0%	100.0%	100.0%
			% of Total	31.9%	67.8%	.3%	100.0%

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	15.052(a)	10	.130
Likelihood Ratio	13.998	10	.173
Linear-by-Linear Association	8.478	1	.004
N of Valid Cases	727		

a 13 cells (72.2%) have expected count less than 5. The minimum expected count is .01.

## TRIED TO COMMIT SUICIDE \* Gender

### Crosstab

			Gender			Total
			Male	Female	Unknown	
TRIED TO COMMIT SUICIDE	Never	Count	227	489	2	718
		Expected Count	227.8	488.2	2.0	718.0
		% within TRIED TO COMMIT SUICIDE	31.6%	68.1%	.3%	100.0%
		% within Gender	98.3%	98.8%	100.0%	98.6%
		% of Total	31.2%	67.2%	.3%	98.6%
	Once	Count	2	5	0	7
		Expected Count	2.2	4.8	.0	7.0
		% within TRIED TO COMMIT SUICIDE	28.6%	71.4%	.0%	100.0%
		% within Gender	.9%	1.0%	.0%	1.0%
		% of Total	.3%	.7%	.0%	1.0%
	Twice	Count	1	1	0	2
		Expected Count	.6	1.4	.0	2.0
		% within TRIED TO COMMIT SUICIDE	50.0%	50.0%	.0%	100.0%

		% within Gender	.4%	.2%	.0%	.3%
		% of Total	.1%	.1%	.0%	.3%
3-5 times	Count	1	0	0	1	
	Expected Count	.3	.7	.0	1.0	
	% within TRIED TO COMMIT SUICIDE	100.0%	.0%	.0%	100.0%	
	% within Gender	.4%	.0%	.0%	.1%	
	% of Total	.1%	.0%	.0%	.1%	
Total	Count	231	495	2	728	
	Expected Count	231.0	495.0	2.0	728.0	
	% within TRIED TO COMMIT SUICIDE	31.7%	68.0%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	31.7%	68.0%	.3%	100.0%	

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	2.521(a)	6	.866
Likelihood Ratio	2.668	6	.849
Linear-by-Linear Association	1.283	1	.257
N of Valid Cases	728		

a 10 cells (83.3%) have expected count less than 5. The minimum expected count is .00.

## BEEN HURT/INJURED \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
Never	Count	162	370	1	533	
	Expected Count	169.4	362.1	1.5	533.0	
	% within BEEN HURT/INJURED	30.4%	69.4%	.2%	100.0%	
	% within Gender	69.8%	74.6%	50.0%	73.0%	
	% of Total	22.2%	50.7%	.1%	73.0%	
Once	Count	28	67	0	95	
	Expected Count	30.2	64.5	.3	95.0	
	% within BEEN HURT/INJURED	29.5%	70.5%	.0%	100.0%	
	% within Gender	12.1%	13.5%	.0%	13.0%	
	% of Total	3.8%	9.2%	.0%	13.0%	
	Count	24	24	0	48	
	Expected Count	15.3	32.6	.1	48.0	

BEEN HURT/INJURED	Twice	% within BEEN HURT/INJURED	50.0%	50.0%	.0%	100.0%
		% within Gender	10.3%	4.8%	.0%	6.6%
		% of Total	3.3%	3.3%	.0%	6.6%
	3-5 times	Count	11	26	1	38
		Expected Count	12.1	25.8	.1	38.0
		% within BEEN HURT/INJURED	28.9%	68.4%	2.6%	100.0%
		% within Gender	4.7%	5.2%	50.0%	5.2%
	6-9 times	% of Total	1.5%	3.6%	.1%	5.2%
		Count	2	6	0	8
		Expected Count	2.5	5.4	.0	8.0
		% within BEEN HURT/INJURED	25.0%	75.0%	.0%	100.0%
	10+ times	% within Gender	.9%	1.2%	.0%	1.1%
		% of Total	.3%	.8%	.0%	1.1%
		Count	5	3	0	8
		Expected Count	2.5	5.4	.0	8.0
	Total	% within BEEN HURT/INJURED	62.5%	37.5%	.0%	100.0%
		% within Gender	2.2%	.6%	.0%	1.1%
		% of Total	.7%	.4%	.0%	1.1%
		Count	232	496	2	730
		Expected Count	232.0	496.0	2.0	730.0
		% within BEEN HURT/INJURED	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	20.062(a)	10	.029
Likelihood Ratio	14.722	10	.143
Linear-by-Linear Association	2.420	1	.120
N of Valid Cases	730		

a 8 cells (44.4%) have expected count less than 5. The minimum expected count is .02.

## A/D problems: mother \* Gender

### Crosstab

		Gender			Total	
		Male	Female	Unknown		
		Count	216	442	2	660
		Expected Count	209.9	448.3	1.8	660.0

A/D problems: mother	No	% within A/D problems: mother	32.7%	67.0%	.3%	100.0%
		% within Gender	94.3%	90.4%	100.0%	91.7%
		% of Total	30.0%	61.4%	.3%	91.7%
	Yes	Count	13	47	0	60
		Expected Count	19.1	40.8	.2	60.0
		% within A/D problems: mother	21.7%	78.3%	.0%	100.0%
		% within Gender	5.7%	9.6%	.0%	8.3%
		% of Total	1.8%	6.5%	.0%	8.3%
		Count	229	489	2	720
		Expected Count	229.0	489.0	2.0	720.0
		% within A/D problems: mother	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%
Total						

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	3.343(a)	2	.188
Likelihood Ratio	3.709	2	.157
Linear-by-Linear Association	2.870	1	.090
N of Valid Cases	720		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .17.

## A/D problems: father \* Gender

## Crosstab

			Gender			Total	
			Male	Female	Unknown		
A/D problems: father	No	Count	184	389	2	575	
		Expected Count	182.9	390.5	1.6	575.0	
		% within A/D problems: father	32.0%	67.7%	.3%	100.0%	
		% within Gender	80.3%	79.6%	100.0%	79.9%	
		% of Total	25.6%	54.0%	.3%	79.9%	
	Yes	Count	45	100	0	145	
		Expected Count	46.1	98.5	.4	145.0	
		% within A/D problems: father	31.0%	69.0%	.0%	100.0%	
		% within Gender	19.7%	20.4%	.0%	20.1%	
		% of Total	6.3%	13.9%	.0%	20.1%	
		Count	229	489	2	720	
		Expected Count	229.0	489.0	2.0	720.0	

Total	% within A/D problems: father	31.8%	67.9%	.3%	100.0%
	% within Gender	100.0%	100.0%	100.0%	100.0%
	% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.568(a)	2	.753
Likelihood Ratio	.963	2	.618
Linear-by-Linear Association	.020	1	.888
N of Valid Cases	720		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .40.

**A/D problems: stepmother \* Gender**

## Crosstab

		Gender			Total
		Male	Female	Unknown	
A/D problems: stepmother	No	Count	227	483	2
		Expected Count	226.5	483.6	2.0
		% within A/D problems: stepmother	31.9%	67.8%	.3% 100.0%
		% within Gender	99.1%	98.8%	100.0% 98.9%
		% of Total	31.5%	67.1%	.3% 98.9%
	Yes	Count	2	6	0
		Expected Count	2.5	5.4	.0
		% within A/D problems: stepmother	25.0%	75.0%	.0% 100.0%
		% within Gender	.9%	1.2%	.0% 1.1%
		% of Total	.3%	.8%	.0% 1.1%
Total		Count	229	489	2
		Expected Count	229.0	489.0	2.0
		% within A/D problems: stepmother	31.8%	67.9%	.3% 100.0%
		% within Gender	100.0%	100.0%	100.0% 100.0%
		% of Total	31.8%	67.9%	.3% 100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.200(a)	2	.905
Likelihood Ratio	.230	2	.892
Linear-by-Linear Association	.155	1	.693
N of Valid Cases	720		

a 3 cells (50.0%) have expected count less than 5. The minimum expected count is .02.

## A/D problems: stepfather \* Gender

Crosstab

		Gender			Total	
		Male	Female	Unknown		
A/D problems: stepfather	No	Count	222	474	2	698
		Expected Count	222.0	474.1	1.9	698.0
		% within A/D problems: stepfather	31.8%	67.9%	.3%	100.0%
		% within Gender	96.9%	96.9%	100.0%	96.9%
		% of Total	30.8%	65.8%	.3%	96.9%
	Yes	Count	7	15	0	22
		Expected Count	7.0	14.9	.1	22.0
		% within A/D problems: stepfather	31.8%	68.2%	.0%	100.0%
		% within Gender	3.1%	3.1%	.0%	3.1%
		% of Total	1.0%	2.1%	.0%	3.1%
Total		Count	229	489	2	720
		Expected Count	229.0	489.0	2.0	720.0
		% within A/D problems: stepfather	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.063(a)	2	.969
Likelihood Ratio	.124	2	.940
Linear-by-Linear Association	.001	1	.977
N of Valid Cases	720		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .06.

## A/D problems: br/sister \* Gender

Crosstab

		Gender			Total	
		Male	Female	Unknown		
		Count	216	436	2	654
		Expected Count	208.0	444.2	1.8	654.0

A/D problems: br/sister	No	% within A/D problems: br/sister	33.0%	66.7%	.3%	100.0%
		% within Gender	94.3%	89.2%	100.0%	90.8%
		% of Total	30.0%	60.6%	.3%	90.8%
	Yes	Count	13	53	0	66
		Expected Count	21.0	44.8	.2	66.0
		% within A/D problems: br/sister	19.7%	80.3%	.0%	100.0%
		% within Gender	5.7%	10.8%	.0%	9.2%
		% of Total	1.8%	7.4%	.0%	9.2%
		Count	229	489	2	720
		Expected Count	229.0	489.0	2.0	720.0
		% within A/D problems: br/sister	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%
Total						

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	5.193(a)	2	.075
Likelihood Ratio	5.772	2	.056
Linear-by-Linear Association	4.586	1	.032
N of Valid Cases	720		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .18.

## A/D problems: mother's parents \* Gender

## Crosstab

A/D problems: mother's parents			Gender			Total
			Male	Female	Unknown	
			Count	Expected Count	%	
	No	Count	195	397	2	594
		Expected Count	188.9	403.4	1.7	594.0
		% within A/D problems: mother's parents	32.8%	66.8%	.3%	100.0%
	Yes	% within Gender	85.2%	81.2%	100.0%	82.5%
		% of Total	27.1%	55.1%	.3%	82.5%
		Count	34	92	0	126
		Expected Count	40.1	85.6	.4	126.0
		% within A/D problems: mother's parents	27.0%	73.0%	.0%	100.0%
		% within Gender	14.8%	18.8%	.0%	17.5%
		% of Total	4.7%	12.8%	.0%	17.5%
		Count	229	489	2	720

Total	Expected Count	229.0	489.0	2.0	720.0
	% within A/D problems: mother's parents	31.8%	67.9%	.3%	100.0%
	% within Gender	100.0%	100.0%	100.0%	100.0%
	% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	2.125(a)	2	.346
Likelihood Ratio	2.508	2	.285
Linear-by-Linear Association	1.422	1	.233
N of Valid Cases	720		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .35.

## A/D problems: father's parents \* Gender

## Crosstab

		No	Gender			Total
			Male	Female	Unknown	
A/D problems: father's parents	No	Count	200	421	2	623
		Expected Count	198.1	423.1	1.7	623.0
		% within A/D problems: father's parents	32.1%	67.6%	.3%	100.0%
		% within Gender	87.3%	86.1%	100.0%	86.5%
		% of Total	27.8%	58.5%	.3%	86.5%
	Yes	Count	29	68	0	97
		Expected Count	30.9	65.9	.3	97.0
		% within A/D problems: father's parents	29.9%	70.1%	.0%	100.0%
		% within Gender	12.7%	13.9%	.0%	13.5%
		% of Total	4.0%	9.4%	.0%	13.5%
Total		Count	229	489	2	720
		Expected Count	229.0	489.0	2.0	720.0
		% within A/D problems: father's parents	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.519(a)	2	.772

Likelihood Ratio	.788	2	.674
Linear-by-Linear Association	.134	1	.714
N of Valid Cases	720		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .27.

## A/D problems: aunts/uncles \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
A/D problems: aunts/uncles	No	Count	163	331	1	495
		Expected Count	157.4	336.2	1.4	495.0
		% within A/D problems: aunts/uncles	32.9%	66.9%	.2%	100.0%
		% within Gender	71.2%	67.7%	50.0%	68.8%
		% of Total	22.6%	46.0%	.1%	68.8%
	Yes	Count	66	158	1	225
		Expected Count	71.6	152.8	.6	225.0
		% within A/D problems: aunts/uncles	29.3%	70.2%	.4%	100.0%
		% within Gender	28.8%	32.3%	50.0%	31.3%
		% of Total	9.2%	21.9%	.1%	31.3%
Total		Count	229	489	2	720
		Expected Count	229.0	489.0	2.0	720.0
		% within A/D problems: aunts/uncles	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1.212(a)	2	.545
Likelihood Ratio	1.196	2	.550
Linear-by-Linear Association	1.028	1	.311
N of Valid Cases	720		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .63.

## A/D problems: spouse \* Gender

Crosstab

	Gender		Total
	Male	Female	

			Male	Female	Unknown	
A/D problems: spouse	No	Count	229	484	2	715
		Expected Count	227.4	485.6	2.0	715.0
		% within A/D problems: spouse	32.0%	67.7%	.3%	100.0%
		% within Gender	100.0%	99.0%	100.0%	99.3%
		% of Total	31.8%	67.2%	.3%	99.3%
	Yes	Count	0	5	0	5
		Expected Count	1.6	3.4	.0	5.0
		% within A/D problems: spouse	.0%	100.0%	.0%	100.0%
		% within Gender	.0%	1.0%	.0%	.7%
		% of Total	.0%	.7%	.0%	.7%
Total	Count		229	489	2	720
	Expected Count		229.0	489.0	2.0	720.0
	% within A/D problems: spouse		31.8%	67.9%	.3%	100.0%
	% within Gender		100.0%	100.0%	100.0%	100.0%
	% of Total		31.8%	67.9%	.3%	100.0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	2.378(a)	2	.304
Likelihood Ratio	3.885	2	.143
Linear-by-Linear Association	2.257	1	.133
N of Valid Cases	720		

a 4 cells (66.7%) have expected count less than 5. The minimum expected count is .01.

## A/D problems: children \* Gender

#### Crosstab

			Gender			Total
			Male	Female	Unknown	
A/D problems: children	No	Count	229	489	2	720
		Expected Count	229.0	489.0	2.0	720.0
		% within A/D problems: children	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%
	Total	Count	229	489	2	720
		Expected Count	229.0	489.0	2.0	720.0
		% within A/D problems: children	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

	% of Total	31.8%	67.9%	.3%	100.0%
--	------------	-------	-------	-----	--------

### Chi-Square Tests

	Value
Pearson Chi-Square	.(a)
N of Valid Cases	720

a No statistics are computed because A/D problems: children is a constant.

## A/D problems: none \* Gender

### Crosstab

		Count	Gender			Total
			Male	Female	Unknown	
A/D problems: none	No	Count	112	291	1	404
		Expected Count	128.5	274.4	1.1	404.0
		% within A/D problems: none	27.7%	72.0%	.2%	100.0%
		% within Gender	48.9%	59.5%	50.0%	56.1%
		% of Total	15.6%	40.4%	.1%	56.1%
	Yes	Count	117	198	1	316
		Expected Count	100.5	214.6	.9	316.0
		% within A/D problems: none	37.0%	62.7%	.3%	100.0%
		% within Gender	51.1%	40.5%	50.0%	43.9%
		% of Total	16.3%	27.5%	.1%	43.9%
Total		Count	229	489	2	720
		Expected Count	229.0	489.0	2.0	720.0
		% within A/D problems: none	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	7.148(a)	2	.028
Likelihood Ratio	7.122	2	.028
Linear-by-Linear Association	6.818	1	.009
N of Valid Cases	720		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .88.

## VOLUNTEERING \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
VOLUNTEERING	DON'T OR < 1 HOUR	Count	111	200	0	311
		Expected Count	98.6	211.4	.9	311.0
		% within VOLUNTEERING	35.7%	64.3%	.0%	100.0%
		% within Gender	53.4%	44.8%	.0%	47.4%
		% of Total	16.9%	30.5%	.0%	47.4%
	1-4 HOURS	Count	49	134	1	184
		Expected Count	58.3	125.1	.6	184.0
		% within VOLUNTEERING	26.6%	72.8%	.5%	100.0%
		% within Gender	23.6%	30.0%	50.0%	28.0%
		% of Total	7.5%	20.4%	.2%	28.0%
	5-9 HOURS	Count	23	62	1	86
		Expected Count	27.3	58.5	.3	86.0
		% within VOLUNTEERING	26.7%	72.1%	1.2%	100.0%
		% within Gender	11.1%	13.9%	50.0%	13.1%
		% of Total	3.5%	9.5%	.2%	13.1%
	10-15 HOURS	Count	9	27	0	36
		Expected Count	11.4	24.5	.1	36.0
		% within VOLUNTEERING	25.0%	75.0%	.0%	100.0%
		% within Gender	4.3%	6.1%	.0%	5.5%
		% of Total	1.4%	4.1%	.0%	5.5%
	16 OR MORE HOURS	Count	16	23	0	39
		Expected Count	12.4	26.5	.1	39.0
		% within VOLUNTEERING	41.0%	59.0%	.0%	100.0%
		% within Gender	7.7%	5.2%	.0%	5.9%
		% of Total	2.4%	3.5%	.0%	5.9%
Total	Count	208	446	2	656	
	Expected Count	208.0	446.0	2.0	656.0	
	% within VOLUNTEERING	31.7%	68.0%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	31.7%	68.0%	.3%	100.0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	11.088(a)	8	.197
Likelihood Ratio	11.344	8	.183
Linear-by-Linear Association	.693	1	.405
N of Valid Cases	656		

a 5 cells (33.3%) have expected count less than 5. The minimum expected count is .11.

# PARTICIPATED-INTERCOLLEGIATE ATHLETICS \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
PARTICIPATED-INTERCOLLEGIATE ATHLETICS	NOT INVOLVED	Count	201	461	2	664
		Expected Count	211.3	450.9	1.8	664.0
		% within PARTICIPATED-INTERCOLLEGIATE ATHLETICS	30.3%	69.4%	.3%	100.0%
		% within Gender	87.0%	93.5%	100.0%	91.5%
		% of Total	27.7%	63.5%	.3%	91.5%
	ACTIVE, NONLEADER	Count	23	28	0	51
		Expected Count	16.2	34.6	.1	51.0
		% within PARTICIPATED-INTERCOLLEGIATE ATHLETICS	45.1%	54.9%	.0%	100.0%
		% within Gender	10.0%	5.7%	.0%	7.0%
		% of Total	3.2%	3.9%	.0%	7.0%
	LEADERSHIP POSITION	Count	7	4	0	11
		Expected Count	3.5	7.5	.0	11.0
		% within PARTICIPATED-INTERCOLLEGIATE ATHLETICS	63.6%	36.4%	.0%	100.0%
		% within Gender	3.0%	.8%	.0%	1.5%
		% of Total	1.0%	.6%	.0%	1.5%
Total		Count	231	493	2	726
		Expected Count	231.0	493.0	2.0	726.0
		% within PARTICIPATED-INTERCOLLEGIATE ATHLETICS	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	10.121(a)	4	.038
Likelihood Ratio	9.596	4	.048
Linear-by-Linear Association	10.045	1	.002

N of Valid Cases	726	
a 4 cells (44.4%) have expected count less than 5. The minimum expected count is .03.		

# PARTICIPATED-INTRAMURAL OR CLUB SPORTS \* Gender

Crosstab

		Gender			Total
		Male	Female	Unknown	
PARTICIPATED-INTRAMURAL OR CLUB SPORTS	NOT INVOLVED	Count	149	421	1 571
		Expected Count	182.5	387.0	1.6 571.0
		% within PARTICIPATED-INTRAMURAL OR CLUB SPORTS	26.1%	73.7%	.2% 100.0%
		% within Gender	64.2%	85.6%	50.0% 78.7%
		% of Total	20.5%	58.0%	.1% 78.7%
	ACTIVE, NONLEADER	Count	77	64	1 142
		Expected Count	45.4	96.2	.4 142.0
		% within PARTICIPATED-INTRAMURAL OR CLUB SPORTS	54.2%	45.1%	.7% 100.0%
		% within Gender	33.2%	13.0%	50.0% 19.6%
		% of Total	10.6%	8.8%	.1% 19.6%
	LEADERSHIP POSITION	Count	6	7	0 13
		Expected Count	4.2	8.8	.0 13.0
		% within PARTICIPATED-INTRAMURAL OR CLUB SPORTS	46.2%	53.8%	.0% 100.0%
		% within Gender	2.6%	1.4%	.0% 1.8%
		% of Total	.8%	1.0%	.0% 1.8%
Total		Count	232	492	2 726
		Expected Count	232.0	492.0	2.0 726.0
		% within PARTICIPATED-INTRAMURAL OR CLUB SPORTS	32.0%	67.8%	.3% 100.0%
		% within Gender	100.0%	100.0%	100.0% 100.0%
		% of Total	32.0%	67.8%	.3% 100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	44.350(a)	4	.000
Likelihood Ratio	41.999	4	.000

Linear-by-Linear Association	35.079	1	.000
N of Valid Cases	726		
a 4 cells (44.4%) have expected count less than 5. The minimum expected count is .04.			

## PARTICIPATED-SOCIAL FRATERNITIES OR SOR \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
PARTICIPATED-SOCIAL FRATERNITIES OR SOR	NOT INVOLVED	Count	178	420	1	599
		Expected Count	190.3	407.1	1.7	599.0
		% within PARTICIPATED-SOCIAL FRATERNITIES OR SOR	29.7%	70.1%	.2%	100.0%
		% within Gender	77.4%	85.4%	50.0%	82.7%
		% of Total	24.6%	58.0%	.1%	82.7%
	ATTENDED	Count	18	38	0	56
		Expected Count	17.8	38.1	.2	56.0
		% within PARTICIPATED-SOCIAL FRATERNITIES OR SOR	32.1%	67.9%	.0%	100.0%
		% within Gender	7.8%	7.7%	.0%	7.7%
		% of Total	2.5%	5.2%	.0%	7.7%
	ACTIVE, NONLEADER	Count	22	24	0	46
		Expected Count	14.6	31.3	.1	46.0
		% within PARTICIPATED-SOCIAL FRATERNITIES OR SOR	47.8%	52.2%	.0%	100.0%
		% within Gender	9.6%	4.9%	.0%	6.4%
		% of Total	3.0%	3.3%	.0%	6.4%
	LEADERSHIP POSITION	Count	12	10	1	23
		Expected Count	7.3	15.6	.1	23.0
		% within PARTICIPATED-SOCIAL FRATERNITIES OR SOR	52.2%	43.5%	4.3%	100.0%
		% within Gender	5.2%	2.0%	50.0%	3.2%
		% of Total	1.7%	1.4%	.1%	3.2%
Total		Count	230	492	2	724
		Expected Count	230.0	492.0	2.0	724.0
		% within PARTICIPATED-SOCIAL FRATERNITIES OR SOR	31.8%	68.0%	.3%	100.0%

SOR				
% within Gender	100.0%	100.0%	100.0%	100.0%
% of Total	31.8%	68.0%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	26.014(a)	6	.000
Likelihood Ratio	15.636	6	.016
Linear-by-Linear Association	8.400	1	.004
N of Valid Cases	724		

a 4 cells (33.3%) have expected count less than 5. The minimum expected count is .06.

## PARTICIPATED-RELIGIOUS GROUPS \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
PARTICIPATED-RELIGIOUS GROUPS	NOT INVOLVED	Count	159	316	1	476
		Expected Count	152.1	322.6	1.3	476.0
		% within PARTICIPATED-RELIGIOUS GROUPS	33.4%	66.4%	.2%	100.0%
		% within Gender	68.5%	64.2%	50.0%	65.6%
		% of Total	21.9%	43.5%	.1%	65.6%
	ATTENDED	Count	47	110	0	157
		Expected Count	50.2	106.4	.4	157.0
		% within PARTICIPATED-RELIGIOUS GROUPS	29.9%	70.1%	.0%	100.0%
		% within Gender	20.3%	22.4%	.0%	21.6%
		% of Total	6.5%	15.2%	.0%	21.6%
	ACTIVE, NONLEADER	Count	17	48	1	66
		Expected Count	21.1	44.7	.2	66.0
		% within PARTICIPATED-RELIGIOUS GROUPS	25.8%	72.7%	1.5%	100.0%
		% within Gender	7.3%	9.8%	50.0%	9.1%
		% of Total	2.3%	6.6%	.1%	9.1%
	LEADERSHIP POSITION	Count	9	18	0	27
		Expected Count	8.6	18.3	.1	27.0
		% within PARTICIPATED-RELIGIOUS GROUPS	33.3%	66.7%	.0%	100.0%

		% within Gender	3.9%	3.7%	.0%	3.7%
		% of Total	1.2%	2.5%	.0%	3.7%
Total		Count	232	492	2	726
		Expected Count	232.0	492.0	2.0	726.0
		% within PARTICIPATED-RELIGIOUS GROUPS	32.0%	67.8%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	32.0%	67.8%	.3%	100.0%

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	6.085(a)	6	.414
Likelihood Ratio	4.740	6	.578
Linear-by-Linear Association	1.191	1	.275
N of Valid Cases	726		

a 4 cells (33.3%) have expected count less than 5. The minimum expected count is .07.

## PARTICIPATED-INTERNATIONAL AND LANGUAGE GPS \* Gender

**Crosstab**

			Gender			Total
			Male	Female	Unknown	
PARTICIPATED-INTERNATIONAL AND LANGUAGE GPS	NOT INVOLVED	Count	219	457	2	678
		Expected Count	215.7	460.4	1.9	678.0
		% within PARTICIPATED-INTERNATIONAL AND LANGUAGE GPS	32.3%	67.4%	.3%	100.0%
		% within Gender	94.8%	92.7%	100.0%	93.4%
		% of Total	30.2%	62.9%	.3%	93.4%
	ATTENDED	Count	10	29	0	39
		Expected Count	12.4	26.5	.1	39.0
		% within PARTICIPATED-INTERNATIONAL AND LANGUAGE GPS	25.6%	74.4%	.0%	100.0%
		% within Gender	4.3%	5.9%	.0%	5.4%
		% of Total	1.4%	4.0%	.0%	5.4%
	ACTIVE, NONLEADER	Count	1	5	0	6
		Expected Count	1.9	4.1	.0	6.0
		% within PARTICIPATED-INTERNATIONAL AND	16.7%	83.3%	.0%	100.0%

		LANGUAGE GPS			
		% within Gender	.4%	1.0%	.0%
		% of Total	.1%	.7%	.0%
Total	LEADERSHIP POSITION	Count	1	2	0
		Expected Count	1.0	2.0	.0
		% within PARTICIPATED-INTERNATIONAL AND LANGUAGE GPS	33.3%	66.7%	.0% 100.0%
		% within Gender	.4%	.4%	.0% .4%
		% of Total	.1%	.3%	.0% .4%
		Count	231	493	2
		Expected Count	231.0	493.0	2.0
		% within PARTICIPATED-INTERNATIONAL AND LANGUAGE GPS	31.8%	67.9%	.3% 100.0%
		% within Gender	100.0%	100.0%	100.0% 100.0%
		% of Total	31.8%	67.9%	.3% 100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1.569(a)	6	.955
Likelihood Ratio	1.805	6	.937
Linear-by-Linear Association	.817	1	.366
N of Valid Cases	726		

a 8 cells (66.7%) have expected count less than 5. The minimum expected count is .01.

## PARTICIPATED-MINORITY AND ETHNIC GROUPS \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
NOT INVOLVED		Count	208	457	2	667
		Expected Count	213.1	452.0	1.8	667.0
		% within PARTICIPATED-MINORITY AND ETHNIC GROUPS	31.2%	68.5%	.3%	100.0%
		% within Gender	89.7%	92.9%	100.0%	91.9%
		% of Total	28.7%	62.9%	.3%	91.9%
		Count	19	21	0	40
		Expected Count	12.8	27.1	.1	40.0

PARTICIPATED-MINORITY AND ETHNIC GROUPS	ATTENDED	% within PARTICIPATED-MINORITY AND ETHNIC GROUPS	47.5%	52.5%	.0%	100.0%
		% within Gender	8.2%	4.3%	.0%	5.5%
		% of Total	2.6%	2.9%	.0%	5.5%
	ACTIVE, NONLEADER	Count	2	13	0	15
		Expected Count	4.8	10.2	.0	15.0
		% within PARTICIPATED-MINORITY AND ETHNIC GROUPS	13.3%	86.7%	.0%	100.0%
		% within Gender	.9%	2.6%	.0%	2.1%
		% of Total	.3%	1.8%	.0%	2.1%
	LEADERSHIP POSITION	Count	3	1	0	4
		Expected Count	1.3	2.7	.0	4.0
		% within PARTICIPATED-MINORITY AND ETHNIC GROUPS	75.0%	25.0%	.0%	100.0%
		% within Gender	1.3%	.2%	.0%	.6%
		% of Total	.4%	.1%	.0%	.6%
Total		Count	232	492	2	726
		Expected Count	232.0	492.0	2.0	726.0
		% within PARTICIPATED-MINORITY AND ETHNIC GROUPS	32.0%	67.8%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	32.0%	67.8%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	10.574(a)	6	.102
Likelihood Ratio	10.557	6	.103
Linear-by-Linear Association	1.287	1	.257
N of Valid Cases	726		

a 7 cells (58.3%) have expected count less than 5. The minimum expected count is .01.

## PARTICIPATED-POLITICAL AND SOCIAL ACTION GPS \* Gender

## Crosstab

	Count	Gender			Total
		Male	Female	Unknown	
		190	423	2	615

PARTICIPATED-POLITICAL AND SOCIAL ACTION GPS	NOT INVOLVED	Expected Count	196.8	416.5	1.7	615.0
		% within PARTICIPATED-POLITICAL AND SOCIAL ACTION GPS	30.9%	68.8%	.3%	100.0%
		% within Gender	81.9%	86.2%	100.0%	84.8%
		% of Total	26.2%	58.3%	.3%	84.8%
	ATTENDED	Count	24	44	0	68
		Expected Count	21.8	46.1	.2	68.0
		% within PARTICIPATED-POLITICAL AND SOCIAL ACTION GPS	35.3%	64.7%	.0%	100.0%
		% within Gender	10.3%	9.0%	.0%	9.4%
		% of Total	3.3%	6.1%	.0%	9.4%
	ACTIVE, NONLEADER	Count	16	17	0	33
		Expected Count	10.6	22.3	.1	33.0
		% within PARTICIPATED-POLITICAL AND SOCIAL ACTION GPS	48.5%	51.5%	.0%	100.0%
		% within Gender	6.9%	3.5%	.0%	4.6%
		% of Total	2.2%	2.3%	.0%	4.6%
	LEADERSHIP POSITION	Count	2	7	0	9
		Expected Count	2.9	6.1	.0	9.0
		% within PARTICIPATED-POLITICAL AND SOCIAL ACTION GPS	22.2%	77.8%	.0%	100.0%
		% within Gender	.9%	1.4%	.0%	1.2%
		% of Total	.3%	1.0%	.0%	1.2%
Total	Count	232	491	2	725	
	Expected Count	232.0	491.0	2.0	725.0	
	% within PARTICIPATED-POLITICAL AND SOCIAL ACTION GPS	32.0%	67.7%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	32.0%	67.7%	.3%	100.0%	

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	5.502(a)	6	.481
Likelihood Ratio	5.555	6	.475
Linear-by-Linear Association	2.188	1	.139
N of Valid Cases	725		

a 5 cells (41.7%) have expected count less than 5. The minimum expected count is .02.

## PARTICIPATED-MUSIC,PERFORMING ARTS

# GROUPS \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
PARTICIPATED-MUSIC,PERFORMING ARTS GROUPS	NOT INVOLVED	Count	162	331	2	495
		Expected Count	158.2	335.5	1.4	495.0
		% within PARTICIPATED-MUSIC,PERFORMING ARTS GROUPS	32.7%	66.9%	.4%	100.0%
		% within Gender	69.8%	67.3%	100.0%	68.2%
		% of Total	22.3%	45.6%	.3%	68.2%
	ATTENDED	Count	45	109	0	154
		Expected Count	49.2	104.4	.4	154.0
		% within PARTICIPATED-MUSIC,PERFORMING ARTS GROUPS	29.2%	70.8%	.0%	100.0%
		% within Gender	19.4%	22.2%	.0%	21.2%
		% of Total	6.2%	15.0%	.0%	21.2%
	ACTIVE, NONLEADER	Count	18	43	0	61
		Expected Count	19.5	41.3	.2	61.0
		% within PARTICIPATED-MUSIC,PERFORMING ARTS GROUPS	29.5%	70.5%	.0%	100.0%
		% within Gender	7.8%	8.7%	.0%	8.4%
		% of Total	2.5%	5.9%	.0%	8.4%
	LEADERSHIP POSITION	Count	7	9	0	16
		Expected Count	5.1	10.8	.0	16.0
		% within PARTICIPATED-MUSIC,PERFORMING ARTS GROUPS	43.8%	56.3%	.0%	100.0%
		% within Gender	3.0%	1.8%	.0%	2.2%
		% of Total	1.0%	1.2%	.0%	2.2%
Total		Count	232	492	2	726
		Expected Count	232.0	492.0	2.0	726.0
		% within PARTICIPATED-MUSIC,PERFORMING ARTS GROUPS	32.0%	67.8%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	32.0%	67.8%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)

Pearson Chi-Square	2.842(a)	6	.828
Likelihood Ratio	3.397	6	.758
Linear-by-Linear Association	.005	1	.945
N of Valid Cases	726		

a 4 cells (33.3%) have expected count less than 5. The minimum expected count is .04.

## PARTICIPATED-NEWSPAPER, RADIO, TV \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
PARTICIPATED- NEWSPAPER, RADIO, TV	NOT INVOLVED	Count	204	450	2	656
		Expected Count	209.6	444.6	1.8	656.0
		% within PARTICIPATED- NEWSPAPER, RADIO, TV	31.1%	68.6%	.3%	100.0%
		% within Gender	88.3%	91.8%	100.0%	90.7%
		% of Total	28.2%	62.2%	.3%	90.7%
	ATTENDED	Count	16	21	0	37
		Expected Count	11.8	25.1	.1	37.0
		% within PARTICIPATED- NEWSPAPER, RADIO, TV	43.2%	56.8%	.0%	100.0%
		% within Gender	6.9%	4.3%	.0%	5.1%
		% of Total	2.2%	2.9%	.0%	5.1%
	ACTIVE, NONLEADER	Count	11	16	0	27
		Expected Count	8.6	18.3	.1	27.0
		% within PARTICIPATED- NEWSPAPER, RADIO, TV	40.7%	59.3%	.0%	100.0%
		% within Gender	4.8%	3.3%	.0%	3.7%
		% of Total	1.5%	2.2%	.0%	3.7%
	LEADERSHIP POSITION	Count	0	3	0	3
		Expected Count	1.0	2.0	.0	3.0
		% within PARTICIPATED- NEWSPAPER, RADIO, TV	.0%	100.0%	.0%	100.0%
		% within Gender	.0%	.6%	.0%	.4%
		% of Total	.0%	.4%	.0%	.4%
Total	Count	231	490	2	723	
	Expected Count	231.0	490.0	2.0	723.0	
	% within PARTICIPATED- NEWSPAPER, RADIO, TV	32.0%	67.8%	.3%	100.0%	

	% within Gender	100.0%	100.0%	100.0%	100.0%
	% of Total	32.0%	67.8%	.3%	100.0%

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	4.919(a)	6	.554
Likelihood Ratio	5.854	6	.440
Linear-by-Linear Association	1.139	1	.286
N of Valid Cases	723		

a 6 cells (50.0%) have expected count less than 5. The minimum expected count is .01.

# HAPPENED - ETHNIC HARASSMENT \*

## Gender

**Crosstab**

			Gender			Total
			Male	Female	Unknown	
HAPPENED - ETHNIC HARASSMENT	YES	Count	18	24	0	42
		Expected Count	13.4	28.4	.1	42.0
		% within HAPPENED - ETHNIC HARASSMENT	42.9%	57.1%	.0%	100.0%
		% within Gender	7.7%	4.9%	.0%	5.8%
		% of Total	2.5%	3.3%	.0%	5.8%
	2	Count	215	469	2	686
		Expected Count	219.6	464.6	1.9	686.0
		% within HAPPENED - ETHNIC HARASSMENT	31.3%	68.4%	.3%	100.0%
		% within Gender	92.3%	95.1%	100.0%	94.2%
		% of Total	29.5%	64.4%	.3%	94.2%
Total		Count	233	493	2	728
		Expected Count	233.0	493.0	2.0	728.0
		% within HAPPENED - ETHNIC HARASSMENT	32.0%	67.7%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	32.0%	67.7%	.3%	100.0%

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	2.499(a)	2	.287
Likelihood Ratio	2.504	2	.286

Linear-by-Linear Association	2.481	1	.115
N of Valid Cases	728		
a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .12.			

## HAPPENED - THREATS OF PHYSICAL VIOLENCE \* Gender

Crosstab

		Gender			Total
		Male	Female	Unknown	
HAPPENED - THREATS OF PHYSICAL VIOLENCE	YES	Count	57	37	0 94
		Expected Count	29.8	63.9	.3 94.0
		% within HAPPENED - THREATS OF PHYSICAL VIOLENCE	60.6%	39.4%	.0% 100.0%
		% within Gender	24.8%	7.5%	.0% 13.0%
		% of Total	7.9%	5.1%	.0% 13.0%
	2	Count	173	456	2 631
		Expected Count	200.2	429.1	1.7 631.0
		% within HAPPENED - THREATS OF PHYSICAL VIOLENCE	27.4%	72.3%	.3% 100.0%
		% within Gender	75.2%	92.5%	100.0% 87.0%
		% of Total	23.9%	62.9%	.3% 87.0%
Total		Count	230	493	2 725
		Expected Count	230.0	493.0	2.0 725.0
		% within HAPPENED - THREATS OF PHYSICAL VIOLENCE	31.7%	68.0%	.3% 100.0%
		% within Gender	100.0%	100.0%	100.0% 100.0%
		% of Total	31.7%	68.0%	.3% 100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	41.786(a)	2	.000
Likelihood Ratio	38.959	2	.000
Linear-by-Linear Association	41.564	1	.000
N of Valid Cases	725		
a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .26.			

## HAPPENED - ACTUAL PHYSICAL VIOLENCE \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
HAPPENED - ACTUAL PHYSICAL VIOLENCE	YES	Count	22	12	0	34
		Expected Count	10.9	23.0	.1	34.0
		% within HAPPENED - ACTUAL PHYSICAL VIOLENCE	64.7%	35.3%	.0%	100.0%
		% within Gender	9.4%	2.4%	.0%	4.7%
		% of Total	3.0%	1.7%	.0%	4.7%
	2	Count	211	480	2	693
		Expected Count	222.1	469.0	1.9	693.0
		% within HAPPENED - ACTUAL PHYSICAL VIOLENCE	30.4%	69.3%	.3%	100.0%
		% within Gender	90.6%	97.6%	100.0%	95.3%
		% of Total	29.0%	66.0%	.3%	95.3%
Total		Count	233	492	2	727
		Expected Count	233.0	492.0	2.0	727.0
		% within HAPPENED - ACTUAL PHYSICAL VIOLENCE	32.0%	67.7%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	32.0%	67.7%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	17.493(a)	2	.000
Likelihood Ratio	16.114	2	.000
Linear-by-Linear Association	17.378	1	.000
N of Valid Cases	727		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .09.

## HAPPENED -THEFT INVOLVING FORCE \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
YES		Count	3	12	0	15
		Expected Count	4.8	10.2	.0	15.0
		% within HAPPENED -THEFT	20.0%	80.0%	.0%	100.0%

HAPPENED -THEFT INVOLVING FORCE	INVOLVING FORCE				
		% within Gender		1.3%	2.4% .0% 2.1%
		% of Total		.4%	1.7% .0% 2.1%
	2	Count	229	481	2 712
		Expected Count	227.2	482.8	2.0 712.0
		% within HAPPENED -THEFT INVOLVING FORCE		32.2%	67.6% .3% 100.0%
		% within Gender		98.7%	97.6% 100.0% 97.9%
		% of Total		31.5%	66.2% .3% 97.9%
Total	Count	232	493	2 727	
	Expected Count	232.0	493.0	2.0 727.0	
	% within HAPPENED -THEFT INVOLVING FORCE		31.9%	67.8% .3% 100.0%	
	% within Gender		100.0%	100.0% 100.0% 100.0%	
	% of Total		31.9%	67.8% .3% 100.0%	

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1.059(a)	2	.589
Likelihood Ratio	1.185	2	.553
Linear-by-Linear Association	.934	1	.334
N of Valid Cases	727		

a 3 cells (50.0%) have expected count less than 5. The minimum expected count is .04.

## HAPPENED -FORCED SEXUAL TOUCHING \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
HAPPENED -FORCED SEXUAL TOUCHING	YES	Count	9	45	0	54
		Expected Count	17.3	36.6	.1	54.0
		% within HAPPENED -FORCED SEXUAL TOUCHING		16.7%	83.3% .0%	100.0%
		% within Gender		3.9%	9.1% .0%	7.4%
		% of Total		1.2%	6.2% .0%	7.4%
	2	Count	224	449	2	675
		Expected Count	215.7	457.4	1.9	675.0
		% within HAPPENED -FORCED SEXUAL TOUCHING		33.2%	66.5% .3%	100.0%
		% within Gender		7.4%	14.8% .3%	10.0%
		% of Total		40.7%	66.2% .5%	67.5%

	% within Gender	96.1%	90.9%	100.0%	92.6%
	% of Total	30.7%	61.6%	.3%	92.6%
Total	Count	233	494	2	729
	Expected Count	233.0	494.0	2.0	729.0
	% within HAPPENED -FORCED SEXUAL TOUCHING	32.0%	67.8%	.3%	100.0%
	% within Gender	100.0%	100.0%	100.0%	100.0%
	% of Total	32.0%	67.8%	.3%	100.0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	6.515(a)	2	.038
Likelihood Ratio	7.373	2	.025
Linear-by-Linear Association	5.920	1	.015
N of Valid Cases	729		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .15.

## HAPPENED - UNWANTED SEXUAL INTERCOURSE \* Gender

#### Crosstab

			Gender			Total
			Male	Female	Unknown	
HAPPENED - UNWANTED SEXUAL INTERCOURSE	YES	Count	6	21	0	27
		Expected Count	8.6	18.3	.1	27.0
		% within HAPPENED - UNWANTED SEXUAL INTERCOURSE	22.2%	77.8%	.0%	100.0%
		% within Gender	2.6%	4.3%	.0%	3.7%
		% of Total	.8%	2.9%	.0%	3.7%
	2	Count	226	472	2	700
		Expected Count	223.4	474.7	1.9	700.0
		% within HAPPENED - UNWANTED SEXUAL INTERCOURSE	32.3%	67.4%	.3%	100.0%
		% within Gender	97.4%	95.7%	100.0%	96.3%
		% of Total	31.1%	64.9%	.3%	96.3%
Total	Count	232	493	2	727	
	Expected Count	232.0	493.0	2.0	727.0	
	% within HAPPENED - UNWANTED SEXUAL	31.9%	67.8%	.3%	100.0%	

	INTERCOURSE				
	% within Gender	100.0%	100.0%	100.0%	100.0%
	% of Total	31.9%	67.8%	.3%	100.0%

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	1.313(a)	2	.519
<b>Likelihood Ratio</b>	1.464	2	.481
<b>Linear-by-Linear Association</b>	1.119	1	.290
<b>N of Valid Cases</b>	727		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .07.

## Crosstabs

### Notes

Output Created		16-JUN-2004 09:47:00
Comments		
Input	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	747
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
Syntax	CROSSTABS /TABLES= Q25B1 q25b2 q25b3 q25b4 q25b5 q25b6 q26a q26b q26c q26d q26e q26f q26g q26h q26i q26j q26k q26l q26m q27a q27b q27c q27d q27e q27f q27g q27h q27i q27j q27k q27l q27m q27n q28a q28b q28c q28d q28e q28f q28g q29a q29b q29c q30 q31a q31b q32a q32b q32c q32d q32e q32f q32g q32h q33 q34 q35a q35b q35c q35d q35e q35f q35g q35h q35i q35j q35k q35l q35m q35n q35o q35p q36a q36b q36c BY q05 /STATISTIC=CHISQ /CELLS= COUNT EXPECTED ROW COLUMN TOTAL .	
Resources	Dimensions Requested	2
	Cells Available	116508
	Elapsed Time	0:00:01.75

### Case Processing Summary

	Cases

	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
CONSUMED -ETHNIC HARASSMENT * Gender	268	35.9%	479	64.1%	747	100.0%
CONSUMED - THREATS OF PHYSICAL VIOLENCE * Gender	285	38.2%	462	61.8%	747	100.0%
CONSUMED-ACTUAL PHYSICAL VIOLENCE * Gender	270	36.1%	477	63.9%	747	100.0%
CONSUMED - THEFT INVOLVING FORCE * Gender	258	34.5%	489	65.5%	747	100.0%
CONSUMED - FORCED SEXUAL TOUCHING * Gender	267	35.7%	480	64.3%	747	100.0%
CONSUMED - UNWANTED SEXUAL INTERCOURSE * Gender	263	35.2%	484	64.8%	747	100.0%
CLOSE FRIENDS -TRYING MARIJUANA ONCE OR TWICE * Gender	728	97.5%	19	2.5%	747	100.0%
CLOSE FRIENDS -SMOKING MARIJUANA OCCASIONALLY * Gender	727	97.3%	20	2.7%	747	100.0%
CLOSE FRIENDS -SMOKING MARIJUANA REGULARLY * Gender	728	97.5%	19	2.5%	747	100.0%
CLOSE FRIENDS -TRYING COCAINE ONCE OR TWICE * Gender	729	97.6%	18	2.4%	747	100.0%
CLOSE FRIENDS -TAKING COCAINE REGULARLY * Gender	729	97.6%	18	2.4%	747	100.0%
CLOSE FRIENDS -TRYING LSD ONCE OR TWICE * Gender	727	97.3%	20	2.7%	747	100.0%
CLOSE FRIENDS -TAKING LSD REGULARLY * Gender	729	97.6%	18	2.4%	747	100.0%
CLOSE FRIENDS -TRYING AMPHETAMINES ONCE OR TWICE * Gender	726	97.2%	21	2.8%	747	100.0%
CLOSE FRIENDS -TAKING AMPHETAMINES REGULARLY * Gender	726	97.2%	21	2.8%	747	100.0%
CLOSE FRIENDS -TAKING ONE OR TWO DRINKS NEARLY EVERY DAY * Gender	728	97.5%	19	2.5%	747	100.0%
CLOSE FRIENDS - TAKING FOUR OR FIVE DRINKS NEARLY EVERY DAY * Gender	727	97.3%	20	2.7%	747	100.0%
CLOSE FRIENDS -HAVING 5 OR MORE DRINKS IN ONE SITTING * Gender	728	97.5%	19	2.5%	747	100.0%
CLOSE FRIENDS -TAKING STEROIDS FOR BODY BUILDING * Gender	728	97.5%	19	2.5%	747	100.0%
EFFECTS -BREAKS THE ICE * Gender	730	97.7%	17	2.3%	747	100.0%
EFFECTS - ENHANCES SOCIAL ACTIVITY * Gender	728	97.5%	19	2.5%	747	100.0%
EFFECTS - EASY TO DEAL WITH STRESS * Gender	730	97.7%	17	2.3%	747	100.0%
EFFECTS - FACILITATES A CONNECTION WITH PEERS * Gender	729	97.6%	18	2.4%	747	100.0%
EFFECTS - GIVES PEOPLE SOMETHING TO TALK ABOUT * Gender	731	97.9%	16	2.1%	747	100.0%
EFFECTS - FACILITATES MALE BONDING * Gender	728	97.5%	19	2.5%	747	100.0%
EFFECTS - FACILITATES FEMALE BONDING * Gender	731	97.9%	16	2.1%	747	100.0%
EFFECTS - ALLOWS PEOPLE TO HAVE MORE FUN * Gender	728	97.5%	19	2.5%	747	100.0%
EFFECTS - GIVES PEOPLE SOMETHING TO DO * Gender	728	97.5%	19	2.5%	747	100.0%
EFFECTS - MAKES FOOD TASTE BETTER * Gender	729	97.6%	18	2.4%	747	100.0%
EFFECTS - MAKES WOMEN SEXIER * Gender	729	97.6%	18	2.4%	747	100.0%
EFFECTS - MAKES MEN SEXIER * Gender	728	97.5%	19	2.5%	747	100.0%
EFFECTS - MAKES ME SEXIER * Gender	727	97.3%	20	2.7%	747	100.0%
EFFECTS - FACILITATES SEXUAL OPPORTUNITIES * Gender	731	97.9%	16	2.1%	747	100.0%
DRINKING IS CENTRAL - MALE STUDENTS * Gender	725	97.1%	22	2.9%	747	100.0%

DRINKING IS CENTRAL - FEMALE STUDENTS * Gender	723	96.8%	24	3.2%	747	100.0%
DRINKING IS CENTRAL - FACULTY * Gender	719	96.3%	28	3.7%	747	100.0%
DRINKING IS CENTRAL - ALUMNI * Gender	719	96.3%	28	3.7%	747	100.0%
DRINKING IS CENTRAL - ATHLETES * Gender	718	96.1%	29	3.9%	747	100.0%
DRINKING IS CENTRAL - FRATERNITIES * Gender	724	96.9%	23	3.1%	747	100.0%
DRINKING IS CENTRAL - SORORITIES * Gender	724	96.9%	23	3.1%	747	100.0%
CAMPUS ENVIRONMENT: PROMOTE ALCOHOL USE? * Gender	725	97.1%	22	2.9%	747	100.0%
CAMPUS ENVIRONMENT: PROMOTE OTHER DRUG USE? * Gender	726	97.2%	21	2.8%	747	100.0%
CAMPUS ENVIRONEMNT: DO YOU FEEL SAFE ON CAMPUS? * Gender	727	97.3%	20	2.7%	747	100.0%
COMPARED TO OTHER CAMPUSES: ALCOHOL USE IS * Gender	726	97.2%	21	2.8%	747	100.0%
DO YOU LIVE IN ALCOHOL FREE RESIDENCE HALL * Gender	514	68.8%	233	31.2%	747	100.0%
WOULD YOU LIVE IN ALCOHOL FREE RESIDENCE HALL * Gender	407	54.5%	340	45.5%	747	100.0%
STUDENTS CARE : ALCOHOL - DRUG USE * Gender	722	96.7%	25	3.3%	747	100.0%
STUDENTS CARE : CAMPUS VANDALISM * Gender	719	96.3%	28	3.7%	747	100.0%
STUDENTS CARE : SEXUAL ASSAULT * Gender	718	96.1%	29	3.9%	747	100.0%
STUDENTS CARE : ASSUALTS-NON SEXUAL * Gender	721	96.5%	26	3.5%	747	100.0%
STUDENTS CARE : HARASSMENT - GENDER * Gender	719	96.3%	28	3.7%	747	100.0%
STUDENTS CARE : HARASSMENT - SEXUAL ORIENTATION * Gender	719	96.3%	28	3.7%	747	100.0%
STUDENTS CARE : HARASSMENT - RACE OR ETHNICITY * Gender	719	96.3%	28	3.7%	747	100.0%
STUDENTS CARE : HARASSMENT - RELIGION * Gender	720	96.4%	27	3.6%	747	100.0%
TO WHAT EXTENT : ALCOHOL USE CHANGED-LAST 12 MONTHS * Gender	720	96.4%	27	3.6%	747	100.0%
TO WHAT EXTENT : DRUG USE CHANGED-LAST 12 MONTHS * Gender	714	95.6%	33	4.4%	747	100.0%
RISK - TRYING MARIJUANA ONCE OR TWICE * Gender	720	96.4%	27	3.6%	747	100.0%
RISK - SMOKING MARIJUANA OCCASIONALLY * Gender	716	95.9%	31	4.1%	747	100.0%
RISK - SMOKING MARIJUANA REGULARLY * Gender	721	96.5%	26	3.5%	747	100.0%
RISK - TRYING COCAINE ONCE OR TWICE * Gender	719	96.3%	28	3.7%	747	100.0%
RISK - TAKING COCAINE REGULARLY * Gender	721	96.5%	26	3.5%	747	100.0%
RISK - TRYING LSD ONCE OR TWICE * Gender	715	95.7%	32	4.3%	747	100.0%
RISK - TAKING LSD REGULARLY * Gender	713	95.4%	34	4.6%	747	100.0%
RISK - TRYING AMPHETAMINES ONCE OR TWICE * Gender	715	95.7%	32	4.3%	747	100.0%
RISK - TAKING AMPHETAMINES REGULARLY * Gender	717	96.0%	30	4.0%	747	100.0%
RISK - TAKING ONE OR TWO DRINKS NEARLY EVERY DAY * Gender	721	96.5%	26	3.5%	747	100.0%
RISK - TAKING FOUR OR FIVE DRINKS NEARLY EVERY DAY * Gender	717	96.0%	30	4.0%	747	100.0%
RISK - HAVING 5 OR MORE DRINKS IN ONE SITTING * Gender	719	96.3%	28	3.7%	747	100.0%
RISK - TAKING STEROIDS FOR BODY BUILDING * Gender	716	95.9%	31	4.1%	747	100.0%
RISK - CONSUME ALCOHOL PRIOR TO SEX * Gender	716	95.9%	31	4.1%	747	100.0%

RISK -REGULARLY ENGAGE IN UNPROTECTED SEX * Gender	719	96.3%	28	3.7%	747	100.0%
RISK -REGULARLY ENGAGE IN MULTILE PARTNER,UNPROTECTED SEX * Gender	721	96.5%	26	3.5%	747	100.0%
DID YOU HAVE SEX WITHIN THE LAST YEAR * Gender	721	96.5%	26	3.5%	747	100.0%
DID YOU DRINK ALCOHOL? * Gender	547	73.2%	200	26.8%	747	100.0%
DID YOU USE OTHER DRUGS? * Gender	546	73.1%	201	26.9%	747	100.0%

## CONSUMED -ETHNIC HARASSMENT \* Gender

Crosstab

			Gender		Total
			Male	Female	
CONSUMED -ETHNIC HARASSMENT	YES	Count	3	2	5
		Expected Count	1.8	3.2	5.0
		% within CONSUMED -ETHNIC HARASSMENT	60.0%	40.0%	100.0%
		% within Gender	3.2%	1.1%	1.9%
		% of Total	1.1%	.7%	1.9%
	2	Count	91	172	263
		Expected Count	92.2	170.8	263.0
		% within CONSUMED -ETHNIC HARASSMENT	34.6%	65.4%	100.0%
		% within Gender	96.8%	98.9%	98.1%
		% of Total	34.0%	64.2%	98.1%
Total		Count	94	174	268
		Expected Count	94.0	174.0	268.0
		% within CONSUMED -ETHNIC HARASSMENT	35.1%	64.9%	100.0%
		% within Gender	100.0%	100.0%	100.0%
		% of Total	35.1%	64.9%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	1.390(b)	1	.238		
Continuity Correction(a)	.498	1	.480		
Likelihood Ratio	1.310	1	.252		
Fisher's Exact Test				.348	.235
Linear-by-Linear Association	1.385	1	.239		
N of Valid Cases	268				

a Computed only for a 2x2 table

b 2 cells (50.0%) have expected count less than 5. The minimum expected count is 1.75.

# CONSUMED - THREATS OF PHYSICAL VIOLENCE \* Gender

Crosstab

		Gender		Total	
		Male	Female		
CONSUMED - THREATS OF PHYSICAL VIOLENCE	YES	Count	44	16	60
		Expected Count	22.5	37.5	60.0
		% within CONSUMED - THREATS OF PHYSICAL VIOLENCE	73.3%	26.7%	100.0%
		% within Gender	41.1%	9.0%	21.1%
		% of Total	15.4%	5.6%	21.1%
	2	Count	63	162	225
		Expected Count	84.5	140.5	225.0
		% within CONSUMED - THREATS OF PHYSICAL VIOLENCE	28.0%	72.0%	100.0%
		% within Gender	58.9%	91.0%	78.9%
		% of Total	22.1%	56.8%	78.9%
Total		Count	107	178	285
		Expected Count	107.0	178.0	285.0
		% within CONSUMED - THREATS OF PHYSICAL VIOLENCE	37.5%	62.5%	100.0%
		% within Gender	100.0%	100.0%	100.0%
		% of Total	37.5%	62.5%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	41.515(b)	1	.000		
Continuity Correction(a)	39.605	1	.000		
Likelihood Ratio	40.800	1	.000		
Fisher's Exact Test				.000	.000
Linear-by-Linear Association	41.370	1	.000		
N of Valid Cases	285				

a Computed only for a 2x2 table

b 0 cells (.0%) have expected count less than 5. The minimum expected count is 22.53.

# CONSUMED-ACTUAL PHYSICAL VIOLENCE

## \* Gender

Crosstab

			Gender		Total
			Male	Female	
CONSUMED-ACTUAL PHYSICAL VIOLENCE	YES	Count	21	3	24
		Expected Count	8.7	15.3	24.0
		% within CONSUMED-ACTUAL PHYSICAL VIOLENCE	87.5%	12.5%	100.0%
		% within Gender	21.4%	1.7%	8.9%
		% of Total	7.8%	1.1%	8.9%
	2	Count	77	169	246
		Expected Count	89.3	156.7	246.0
		% within CONSUMED-ACTUAL PHYSICAL VIOLENCE	31.3%	68.7%	100.0%
		% within Gender	78.6%	98.3%	91.1%
		% of Total	28.5%	62.6%	91.1%
Total		Count	98	172	270
		Expected Count	98.0	172.0	270.0
		% within CONSUMED-ACTUAL PHYSICAL VIOLENCE	36.3%	63.7%	100.0%
		% within Gender	100.0%	100.0%	100.0%
		% of Total	36.3%	63.7%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	29.869(b)	1	.000		
Continuity Correction(a)	27.488	1	.000		
Likelihood Ratio	29.900	1	.000		
Fisher's Exact Test				.000	.000
Linear-by-Linear Association	29.758	1	.000		
N of Valid Cases	270				
a Computed only for a 2x2 table					
b 0 cells (.0%) have expected count less than 5. The minimum expected count is 8.71.					

# CONSUMED - THEFT INVOLVING FORCE \* Gender

Crosstab

			Gender		Total
			Male	Female	
CONSUMED - THEFT INVOLVING FORCE	YES	Count	4	1	5
		Expected Count	1.7	3.3	5.0
		% within CONSUMED - THEFT INVOLVING FORCE	80.0%	20.0%	100.0%
		% within Gender	4.5%	.6%	1.9%
		% of Total	1.6%	.4%	1.9%
	2	Count	85	168	253
	2	Expected Count	87.3	165.7	253.0
		% within CONSUMED - THEFT INVOLVING FORCE	33.6%	66.4%	100.0%
		% within Gender	95.5%	99.4%	98.1%
		% of Total	32.9%	65.1%	98.1%
		Count	89	169	258
Total	Total	Expected Count	89.0	169.0	258.0
		% within CONSUMED - THEFT INVOLVING FORCE	34.5%	65.5%	100.0%
		% within Gender	100.0%	100.0%	100.0%
		% of Total	34.5%	65.5%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	4.672(b)	1	.031		
Continuity Correction(a)	2.844	1	.092		
Likelihood Ratio	4.448	1	.035		
Fisher's Exact Test				.050	.050
Linear-by-Linear Association	4.654	1	.031		
N of Valid Cases	258				
a Computed only for a 2x2 table					
b 2 cells (50.0%) have expected count less than 5. The minimum expected count is 1.72.					

## CONSUMED - FORCED SEXUAL TOUCHING \* Gender

Crosstab

			Gender		Total
			Male	Female	

CONSUMED - FORCED SEXUAL TOUCHING	YES	Count	4	34	38
		Expected Count	13.0	25.0	38.0
		% within CONSUMED - FORCED SEXUAL TOUCHING	10.5%	89.5%	100.0%
		% within Gender	4.4%	19.3%	14.2%
		% of Total	1.5%	12.7%	14.2%
	2	Count	87	142	229
		Expected Count	78.0	151.0	229.0
		% within CONSUMED - FORCED SEXUAL TOUCHING	38.0%	62.0%	100.0%
		% within Gender	95.6%	80.7%	85.8%
		% of Total	32.6%	53.2%	85.8%
Total		Count	91	176	267
		Expected Count	91.0	176.0	267.0
		% within CONSUMED - FORCED SEXUAL TOUCHING	34.1%	65.9%	100.0%
		% within Gender	100.0%	100.0%	100.0%
		% of Total	34.1%	65.9%	100.0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	10.943(b)	1	.001		
Continuity Correction(a)	9.755	1	.002		
Likelihood Ratio	12.909	1	.000		
Fisher's Exact Test				.001	.000
Linear-by-Linear Association	10.902	1	.001		
N of Valid Cases	267				

a Computed only for a 2x2 table

b 0 cells (.0%) have expected count less than 5. The minimum expected count is 12.95.

## CONSUMED - UNWANTED SEXUAL INTERCOURSE \* Gender

#### Crosstab

	YES	Count	Gender		Total
			Male	Female	
	YES	Expected Count	6.9	13.1	20.0
		% within CONSUMED - UNWANTED SEXUAL INTERCOURSE	30.0%	70.0%	100.0%

CONSUMED - UNWANTED SEXUAL INTERCOURSE	2	% within Gender	6.6%	8.1%	7.6%	
		% of Total	2.3%	5.3%	7.6%	
		Count	85	158	243	
		Expected Count	84.1	158.9	243.0	
		% within CONSUMED - UNWANTED SEXUAL INTERCOURSE	35.0%	65.0%	100.0%	
		% within Gender	93.4%	91.9%	92.4%	
Total		% of Total	32.3%	60.1%	92.4%	
		Count	91	172	263	
		Expected Count	91.0	172.0	263.0	
		% within CONSUMED - UNWANTED SEXUAL INTERCOURSE	34.6%	65.4%	100.0%	
		% within Gender	100.0%	100.0%	100.0%	
		% of Total	34.6%	65.4%	100.0%	

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.202(b)	1		.653	
Continuity Correction(a)	.042	1		.837	
Likelihood Ratio	.207	1		.649	
Fisher's Exact Test					.808
Linear-by-Linear Association	.202	1		.653	
N of Valid Cases	263				
a Computed only for a 2x2 table					
b 0 cells (.0%) have expected count less than 5. The minimum expected count is 6.92.					

## CLOSE FRIENDS -TRYING MARIJUANA ONCE OR TWICE \* Gender

#### Crosstab

			Gender			Total
			Male	Female	Unknown	
DONT DISAPPROVE		Count	144	265	1	410
		Expected Count	131.2	277.7	1.1	410.0
		% within CLOSE FRIENDS - TRYING MARIJUANA ONCE OR TWICE	35.1%	64.6%	.2%	100.0%
		% within Gender	61.8%	53.8%	50.0%	56.3%
		% of Total	19.8%	36.4%	.1%	56.3%
		Count	52	110	1	163

CLOSE FRIENDS - TRYING MARIJUANA ONCE OR TWICE	DISAPPROVE	Expected Count	52.2	110.4	.4	163.0
		% within CLOSE FRIENDS - TRYING MARIJUANA ONCE OR TWICE	31.9%	67.5%	.6%	100.0%
		% within Gender	22.3%	22.3%	50.0%	22.4%
		% of Total	7.1%	15.1%	.1%	22.4%
	STRONGLY DISAPPROVE	Count	37	118	0	155
		Expected Count	49.6	105.0	.4	155.0
		% within CLOSE FRIENDS - TRYING MARIJUANA ONCE OR TWICE	23.9%	76.1%	.0%	100.0%
		% within Gender	15.9%	23.9%	.0%	21.3%
		% of Total	5.1%	16.2%	.0%	21.3%
Total	Count	233	493	2	728	
	Expected Count	233.0	493.0	2.0	728.0	
	% within CLOSE FRIENDS - TRYING MARIJUANA ONCE OR TWICE	32.0%	67.7%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	32.0%	67.7%	.3%	100.0%	

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	7.767(a)	4	.101
Likelihood Ratio	8.234	4	.083
Linear-by-Linear Association	5.948	1	.015
N of Valid Cases	728		

a 3 cells (33.3%) have expected count less than 5. The minimum expected count is .43.

## CLOSE FRIENDS -SMOKING MARIJUANA OCCASIONALLY \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
DONT DISAPPROVE	DONT DISAPPROVE	Count	109	199	0	308
		Expected Count	98.7	208.4	.8	308.0
		% within CLOSE FRIENDS - SMOKING MARIJUANA OCCASIONALLY	35.4%	64.6%	.0%	100.0%
		% within Gender	46.8%	40.4%	.0%	42.4%
		% of Total	15.0%	27.4%	.0%	42.4%

CLOSE FRIENDS - SMOKING MARIJUANA OCCASIONALLY	DISAPPROVE	Count	59	107	1	167
		Expected Count	53.5	113.0	.5	167.0
		% within CLOSE FRIENDS - SMOKING MARIJUANA OCCASIONALLY	35.3%	64.1%	.6%	100.0%
		% within Gender	25.3%	21.7%	50.0%	23.0%
		% of Total	8.1%	14.7%	.1%	23.0%
	STRONGLY DISAPPROVE	Count	65	186	1	252
	Expected Count	80.8	170.5	.7	252.0	
	% within CLOSE FRIENDS - SMOKING MARIJUANA OCCASIONALLY	25.8%	73.8%	.4%	100.0%	
	% within Gender	27.9%	37.8%	50.0%	34.7%	
	% of Total	8.9%	25.6%	.1%	34.7%	
Total		Count	233	492	2	727
		Expected Count	233.0	492.0	2.0	727.0
		% within CLOSE FRIENDS - SMOKING MARIJUANA OCCASIONALLY	32.0%	67.7%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	32.0%	67.7%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	8.478(a)	4	.076
Likelihood Ratio	9.288	4	.054
Linear-by-Linear Association	5.984	1	.014
N of Valid Cases	727		

a 3 cells (33.3%) have expected count less than 5. The minimum expected count is .46.

## CLOSE FRIENDS -SMOKING MARIJUANA REGULARLY \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
DONT DISAPPROVE	DONT DISAPPROVE	Count	51	73	0	124
		Expected Count	39.5	84.1	.3	124.0
		% within CLOSE FRIENDS - SMOKING MARIJUANA REGULARLY	41.1%	58.9%	.0%	100.0%
		% within Gender	22.0%	14.8%	.0%	17.0%

CLOSE FRIENDS - SMOKING MARIJUANA REGULARLY	DISAPPROVE	% of Total	7.0%	10.0%	.0%	17.0%
		Count	73	135	0	208
		Expected Count	66.3	141.1	.6	208.0
		% within CLOSE FRIENDS - SMOKING MARIJUANA REGULARLY	35.1%	64.9%	.0%	100.0%
		% within Gender	31.5%	27.3%	.0%	28.6%
	STRONGLY DISAPPROVE	% of Total	10.0%	18.5%	.0%	28.6%
		Count	108	286	2	396
		Expected Count	126.2	268.7	1.1	396.0
		% within CLOSE FRIENDS - SMOKING MARIJUANA REGULARLY	27.3%	72.2%	.5%	100.0%
		% within Gender	46.6%	57.9%	100.0%	54.4%
		% of Total	14.8%	39.3%	.3%	54.4%
Total		Count	232	494	2	728
		Expected Count	232.0	494.0	2.0	728.0
		% within CLOSE FRIENDS - SMOKING MARIJUANA REGULARLY	31.9%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.9%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	11.173(a)	4	.025
Likelihood Ratio	11.814	4	.019
Linear-by-Linear Association	10.307	1	.001
N of Valid Cases	728		

a 3 cells (33.3%) have expected count less than 5. The minimum expected count is .34.

## CLOSE FRIENDS -TRYING COCAINE ONCE OR TWICE \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
DONT DISAPPROVE	DONT DISAPPROVE	Count	25	45	0	70
		Expected Count	22.4	47.4	.2	70.0
		% within CLOSE FRIENDS - TRYING COCAINE ONCE OR TWICE	35.7%	64.3%	.0%	100.0%

CLOSE FRIENDS - TRYING COCAINE ONCE OR TWICE	DISAPPROVE	% within Gender	10.7%	9.1%	.0%	9.6%
		% of Total	3.4%	6.2%	.0%	9.6%
		Count	47	71	0	118
		Expected Count	37.7	80.0	.3	118.0
		% within CLOSE FRIENDS - TRYING COCAINE ONCE OR TWICE	39.8%	60.2%	.0%	100.0%
		% within Gender	20.2%	14.4%	.0%	16.2%
	STRONGLY DISAPPROVE	% of Total	6.4%	9.7%	.0%	16.2%
		Count	161	378	2	541
		Expected Count	172.9	366.6	1.5	541.0
		% within CLOSE FRIENDS - TRYING COCAINE ONCE OR TWICE	29.8%	69.9%	.4%	100.0%
		% within Gender	69.1%	76.5%	100.0%	74.2%
Total		% of Total	22.1%	51.9%	.3%	74.2%
		Count	233	494	2	729
		Expected Count	233.0	494.0	2.0	729.0
		% within CLOSE FRIENDS - TRYING COCAINE ONCE OR TWICE	32.0%	67.8%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	32.0%	67.8%	.3%	100.0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	5.594(a)	4	.232
Likelihood Ratio	5.971	4	.201
Linear-by-Linear Association	3.410	1	.065
N of Valid Cases	729		

a 3 cells (33.3%) have expected count less than 5. The minimum expected count is .19.

## CLOSE FRIENDS -TAKING COCAINE REGULARLY \* Gender

#### Crosstab

			Gender			Total
			Male	Female	Unknown	
DONT		Count	0	5	0	5
		Expected Count	1.6	3.4	.0	5.0
		% within CLOSE FRIENDS -				

CLOSE FRIENDS - TAKING COCAINE REGULARLY	DISAPPROVE	TAKING COCAINE REGULARLY	.0%	100.0%	.0%	100.0%	
		% within Gender	.0%	1.0%	.0%	.7%	
		% of Total	.0%	.7%	.0%	.7%	
	DISAPPROVE	Count	33	46	0	79	
		Expected Count	25.2	53.5	.2	79.0	
		% within CLOSE FRIENDS - TAKING COCAINE REGULARLY	41.8%	58.2%	.0%	100.0%	
		% within Gender	14.2%	9.3%	.0%	10.8%	
		% of Total	4.5%	6.3%	.0%	10.8%	
	STRONGLY DISAPPROVE	Count	200	443	2	645	
		Expected Count	206.2	437.1	1.8	645.0	
		% within CLOSE FRIENDS - TAKING COCAINE REGULARLY	31.0%	68.7%	.3%	100.0%	
		% within Gender	85.8%	89.7%	100.0%	88.5%	
		% of Total	27.4%	60.8%	.3%	88.5%	
			Count	233	494	2	729
			Expected Count	233.0	494.0	2.0	729.0
			% within CLOSE FRIENDS - TAKING COCAINE REGULARLY	32.0%	67.8%	.3%	100.0%
			% within Gender	100.0%	100.0%	100.0%	100.0%
			% of Total	32.0%	67.8%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	6.328(a)	4	.176
Likelihood Ratio	7.900	4	.095
Linear-by-Linear Association	1.175	1	.278
N of Valid Cases	729		

a 5 cells (55.6%) have expected count less than 5. The minimum expected count is .01.

## CLOSE FRIENDS -TRYING LSD ONCE OR TWICE \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
		Count	29	51	0	80

CLOSE FRIENDS - TRYING LSD ONCE OR TWICE	DONT DISAPPROVE	Expected Count	25.5	54.3	.2	80.0
		% within CLOSE FRIENDS - TRYING LSD ONCE OR TWICE	36.3%	63.8%	.0%	100.0%
		% within Gender	12.5%	10.3%	.0%	11.0%
		% of Total	4.0%	7.0%	.0%	11.0%
	DISAPPROVE	Count	48	68	0	116
		Expected Count	37.0	78.7	.3	116.0
		% within CLOSE FRIENDS - TRYING LSD ONCE OR TWICE	41.4%	58.6%	.0%	100.0%
		% within Gender	20.7%	13.8%	.0%	16.0%
		% of Total	6.6%	9.4%	.0%	16.0%
	STRONGLY DISAPPROVE	Count	155	374	2	531
		Expected Count	169.5	360.1	1.5	531.0
		% within CLOSE FRIENDS - TRYING LSD ONCE OR TWICE	29.2%	70.4%	.4%	100.0%
		% within Gender	66.8%	75.9%	100.0%	73.0%
		% of Total	21.3%	51.4%	.3%	73.0%
Total	Count	232	493	2	727	
	Expected Count	232.0	493.0	2.0	727.0	
	% within CLOSE FRIENDS - TRYING LSD ONCE OR TWICE	31.9%	67.8%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	31.9%	67.8%	.3%	100.0%	

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	7.878(a)	4	.096
Likelihood Ratio	8.199	4	.085
Linear-by-Linear Association	4.745	1	.029
N of Valid Cases	727		

a 3 cells (33.3%) have expected count less than 5. The minimum expected count is .22.

# CLOSE FRIENDS -TAKING LSD REGULARLY \* Gender

## Crosstab

	Gender			Total
	Male	Female	Unknown	

CLOSE FRIENDS - TAKING LSD REGULARLY	DONT DISAPPROVE	Count	1	4	0	5
		Expected Count	1.6	3.4	.0	5.0
		% within CLOSE FRIENDS -TAKING LSD REGULARLY	20.0%	80.0%	.0%	100.0%
		% within Gender	.4%	.8%	.0%	.7%
		% of Total	.1%	.5%	.0%	.7%
	DISAPRROVE	Count	35	56	0	91
		Expected Count	29.1	61.7	.2	91.0
		% within CLOSE FRIENDS -TAKING LSD REGULARLY	38.5%	61.5%	.0%	100.0%
		% within Gender	15.0%	11.3%	.0%	12.5%
		% of Total	4.8%	7.7%	.0%	12.5%
	STRONGLY DISAPPROVE	Count	197	434	2	633
		Expected Count	202.3	428.9	1.7	633.0
		% within CLOSE FRIENDS -TAKING LSD REGULARLY	31.1%	68.6%	.3%	100.0%
		% within Gender	84.5%	87.9%	100.0%	86.8%
		% of Total	27.0%	59.5%	.3%	86.8%
	Total	Count	233	494	2	729
		Expected Count	233.0	494.0	2.0	729.0
		% within CLOSE FRIENDS -TAKING LSD REGULARLY	32.0%	67.8%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	32.0%	67.8%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	2.560(a)	4	.634
Likelihood Ratio	2.794	4	.593
Linear-by-Linear Association	1.158	1	.282
N of Valid Cases	729		

a 5 cells (55.6%) have expected count less than 5. The minimum expected count is .01.

## CLOSE FRIENDS -TRYING AMPHETAMINES ONCE OR TWICE \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
		Count	14	42	0	56

CLOSE FRIENDS - TRYING AMPHETAMINES ONCE OR TWICE	DONT DISAPPROVE	Expected Count	18.0	37.9	.2	56.0
		% within CLOSE FRIENDS - TRYING AMPHETAMINES ONCE OR TWICE	25.0%	75.0%	.0%	100.0%
		% within Gender	6.0%	8.6%	.0%	7.7%
		% of Total	1.9%	5.8%	.0%	7.7%
	DISAPPROVE	Count	55	88	0	143
		Expected Count	45.9	96.7	.4	143.0
		% within CLOSE FRIENDS - TRYING AMPHETAMINES ONCE OR TWICE	38.5%	61.5%	.0%	100.0%
		% within Gender	23.6%	17.9%	.0%	19.7%
		% of Total	7.6%	12.1%	.0%	19.7%
	STRONGLY DISAPPROVE	Count	164	361	2	527
		Expected Count	169.1	356.4	1.5	527.0
		% within CLOSE FRIENDS - TRYING AMPHETAMINES ONCE OR TWICE	31.1%	68.5%	.4%	100.0%
		% within Gender	70.4%	73.5%	100.0%	72.6%
		% of Total	22.6%	49.7%	.3%	72.6%
Total	Count	233	491	2	726	
	Expected Count	233.0	491.0	2.0	726.0	
	% within CLOSE FRIENDS - TRYING AMPHETAMINES ONCE OR TWICE	32.1%	67.6%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	32.1%	67.6%	.3%	100.0%	

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	4.889(a)	4	.299
Likelihood Ratio	5.390	4	.250
Linear-by-Linear Association	.056	1	.813
N of Valid Cases	726		

a 3 cells (33.3%) have expected count less than 5. The minimum expected count is .15.

# CLOSE FRIENDS -TAKING AMPHETAMINES REGULARLY \* Gender

## Crosstab

	Gender			Total
	Male	Female	Unknown	

CLOSE FRIENDS - TAKING AMPHETAMINES REGULARLY	DONT DISAPPROVE	Count	3	8	0	11	
		Expected Count	3.5	7.5	.0	11.0	
		% within CLOSE FRIENDS - TAKING AMPHETAMINES REGULARLY	27.3%	72.7%	.0%	100.0%	
		% within Gender	1.3%	1.6%	.0%	1.5%	
		% of Total	.4%	1.1%	.0%	1.5%	
	DISAPRROVE	Count	37	60	0	97	
		Expected Count	31.0	65.7	.3	97.0	
		% within CLOSE FRIENDS - TAKING AMPHETAMINES REGULARLY	38.1%	61.9%	.0%	100.0%	
		% within Gender	15.9%	12.2%	.0%	13.4%	
		% of Total	5.1%	8.3%	.0%	13.4%	
	STRONGLY DISAPPROVE	Count	192	424	2	618	
		Expected Count	197.5	418.8	1.7	618.0	
		% within CLOSE FRIENDS - TAKING AMPHETAMINES REGULARLY	31.1%	68.6%	.3%	100.0%	
		% within Gender	82.8%	86.2%	100.0%	85.1%	
		% of Total	26.4%	58.4%	.3%	85.1%	
	Total		Count	232	492	2	726
			Expected Count	232.0	492.0	2.0	726.0
			% within CLOSE FRIENDS - TAKING AMPHETAMINES REGULARLY	32.0%	67.8%	.3%	100.0%
			% within Gender	100.0%	100.0%	100.0%	100.0%
			% of Total	32.0%	67.8%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	2.345(a)	4	.673
Likelihood Ratio	2.590	4	.629
Linear-by-Linear Association	1.040	1	.308
N of Valid Cases	726		

a 4 cells (44.4%) have expected count less than 5. The minimum expected count is .03.

# CLOSE FRIENDS -TAKING ONE OR TWO DRINKS NEARLY EVERY DAY \* Gender

## Crosstab

	Gender
--	--------

			Male	Female	Unknown	Total
CLOSE FRIENDS - TAKING ONE OR TWO DRINKS NEARLY EVERY DAY	DONT DISAPPROVE	Count	135	179	1	315
		Expected Count	100.4	213.8	.9	315.0
		% within CLOSE FRIENDS - TAKING ONE OR TWO DRINKS NEARLY EVERY DAY	42.9%	56.8%	.3%	100.0%
		% within Gender	58.2%	36.2%	50.0%	43.3%
		% of Total	18.5%	24.6%	.1%	43.3%
	DISAPRROVE	Count	69	199	1	269
		Expected Count	85.7	182.5	.7	269.0
		% within CLOSE FRIENDS - TAKING ONE OR TWO DRINKS NEARLY EVERY DAY	25.7%	74.0%	.4%	100.0%
		% within Gender	29.7%	40.3%	50.0%	37.0%
		% of Total	9.5%	27.3%	.1%	37.0%
	STRONGLY DISAPPROVE	Count	28	116	0	144
		Expected Count	45.9	97.7	.4	144.0
		% within CLOSE FRIENDS - TAKING ONE OR TWO DRINKS NEARLY EVERY DAY	19.4%	80.6%	.0%	100.0%
		% within Gender	12.1%	23.5%	.0%	19.8%
		% of Total	3.8%	15.9%	.0%	19.8%
Total		Count	232	494	2	728
		Expected Count	232.0	494.0	2.0	728.0
		% within CLOSE FRIENDS - TAKING ONE OR TWO DRINKS NEARLY EVERY DAY	31.9%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.9%	67.9%	.3%	100.0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	33.239(a)	4	.000
Likelihood Ratio	33.919	4	.000
Linear-by-Linear Association	29.064	1	.000
N of Valid Cases	728		

a 3 cells (33.3%) have expected count less than 5. The minimum expected count is .40.

## CLOSE FRIENDS - TAKING FOUR OR FIVE DRINKS NEARLY EVERY DAY \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
CLOSE FRIENDS - TAKING FOUR OR FIVE DRINKS NEARLY EVERY DAY	DONT DISAPPROVE	Count	54	32	1	87
		Expected Count	27.9	59.0	.1	87.0
		% within CLOSE FRIENDS - TAKING FOUR OR FIVE DRINKS NEARLY EVERY DAY	62.1%	36.8%	1.1%	100.0%
		% within Gender	23.2%	6.5%	100.0%	12.0%
		% of Total	7.4%	4.4%	.1%	12.0%
	DISAPPROVE	Count	95	146	0	241
		Expected Count	77.2	163.4	.3	241.0
		% within CLOSE FRIENDS - TAKING FOUR OR FIVE DRINKS NEARLY EVERY DAY	39.4%	60.6%	.0%	100.0%
		% within Gender	40.8%	29.6%	.0%	33.1%
		% of Total	13.1%	20.1%	.0%	33.1%
	STRONGLY DISAPPROVE	Count	84	315	0	399
		Expected Count	127.9	270.6	.5	399.0
		% within CLOSE FRIENDS - TAKING FOUR OR FIVE DRINKS NEARLY EVERY DAY	21.1%	78.9%	.0%	100.0%
		% within Gender	36.1%	63.9%	.0%	54.9%
		% of Total	11.6%	43.3%	.0%	54.9%
Total		Count	233	493	1	727
		Expected Count	233.0	493.0	1.0	727.0
		% within CLOSE FRIENDS - TAKING FOUR OR FIVE DRINKS NEARLY EVERY DAY	32.0%	67.8%	.1%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	32.0%	67.8%	.1%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	72.466(a)	4	.000
Likelihood Ratio	68.047	4	.000
Linear-by-Linear Association	60.598	1	.000
N of Valid Cases	727		

a 3 cells (33.3%) have expected count less than 5. The minimum expected count is .12.

# CLOSE FRIENDS - HAVING 5 OR MORE DRINKS IN ONE SITTING \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
CLOSE FRIENDS - HAVING 5 OR MORE DRINKS IN ONE SITTING	DONT DISAPPROVE	Count	161	267	2	430
		Expected Count	137.0	291.8	1.2	430.0
		% within CLOSE FRIENDS - HAVING 5 OR MORE DRINKS IN ONE SITTING	37.4%	62.1%	.5%	100.0%
		% within Gender	69.4%	54.0%	100.0%	59.1%
		% of Total	22.1%	36.7%	.3%	59.1%
	DISAPPROVE	Count	40	109	0	149
		Expected Count	47.5	101.1	.4	149.0
		% within CLOSE FRIENDS - HAVING 5 OR MORE DRINKS IN ONE SITTING	26.8%	73.2%	.0%	100.0%
		% within Gender	17.2%	22.1%	.0%	20.5%
		% of Total	5.5%	15.0%	.0%	20.5%
	STRONGLY DISAPPROVE	Count	31	118	0	149
		Expected Count	47.5	101.1	.4	149.0
		% within CLOSE FRIENDS - HAVING 5 OR MORE DRINKS IN ONE SITTING	20.8%	79.2%	.0%	100.0%
		% within Gender	13.4%	23.9%	.0%	20.5%
		% of Total	4.3%	16.2%	.0%	20.5%
Total		Count	232	494	2	728
		Expected Count	232.0	494.0	2.0	728.0
		% within CLOSE FRIENDS - HAVING 5 OR MORE DRINKS IN ONE SITTING	31.9%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.9%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	18.023(a)	4	.001
Likelihood Ratio	19.296	4	.001
Linear-by-Linear Association	14.733	1	.000
N of Valid Cases	728		
a 3 cells (33.3%) have expected count less than 5. The minimum expected count is .41.			

# CLOSE FRIENDS -TAKING STEROIDS FOR BODY BUILDING \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
CLOSE FRIENDS - TAKING STEROIDS FOR BODY BUILDING	DONT DISAPPROVE	Count	21	10	0	31
		Expected Count	9.9	21.0	.1	31.0
		% within CLOSE FRIENDS - TAKING STEROIDS FOR BODY BUILDING	67.7%	32.3%	.0%	100.0%
		% within Gender	9.0%	2.0%	.0%	4.3%
		% of Total	2.9%	1.4%	.0%	4.3%
	DISAPPROVE	Count	63	123	1	187
		Expected Count	59.9	126.6	.5	187.0
		% within CLOSE FRIENDS - TAKING STEROIDS FOR BODY BUILDING	33.7%	65.8%	.5%	100.0%
		% within Gender	27.0%	24.9%	50.0%	25.7%
		% of Total	8.7%	16.9%	.1%	25.7%
	STRONGLY DISAPPROVE	Count	149	360	1	510
		Expected Count	163.2	345.4	1.4	510.0
		% within CLOSE FRIENDS - TAKING STEROIDS FOR BODY BUILDING	29.2%	70.6%	.2%	100.0%
		% within Gender	63.9%	73.0%	50.0%	70.1%
		% of Total	20.5%	49.5%	.1%	70.1%
Total		Count	233	493	2	728
		Expected Count	233.0	493.0	2.0	728.0
		% within CLOSE FRIENDS - TAKING STEROIDS FOR BODY BUILDING	32.0%	67.7%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	32.0%	67.7%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	20.917(a)	4	.000
Likelihood Ratio	19.304	4	.001
Linear-by-Linear Association	12.433	1	.000
N of Valid Cases	728		

a 3 cells (33.3%) have expected count less than 5. The minimum expected count is .09.

## EFFECTS -BREAKS THE ICE \* Gender

Crosstab

		Gender			Total	
		Male	Female	Unknown		
EFFECTS -BREAKS THE ICE	YES	Count	194	411	3	608
		Expected Count	193.2	412.3	2.5	608.0
		% within EFFECTS -BREAKS THE ICE	31.9%	67.6%	.5%	100.0%
		% within Gender	83.6%	83.0%	100.0%	83.3%
		% of Total	26.6%	56.3%	.4%	83.3%
	2	Count	38	84	0	122
		Expected Count	38.8	82.7	.5	122.0
		% within EFFECTS -BREAKS THE ICE	31.1%	68.9%	.0%	100.0%
		% within Gender	16.4%	17.0%	.0%	16.7%
		% of Total	5.2%	11.5%	.0%	16.7%
Total	Count	232	495	3	730	
	Expected Count	232.0	495.0	3.0	730.0	
	% within EFFECTS -BREAKS THE ICE	31.8%	67.8%	.4%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	31.8%	67.8%	.4%	100.0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.644(a)	2	.725
Likelihood Ratio	1.139	2	.566
Linear-by-Linear Association	.003	1	.955
N of Valid Cases	730		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .50.

## EFFECTS - ENHANCES SOCIAL ACTIVITY \* Gender

Crosstab

	Gender

			Male	Female	Unknown	Total
EFFECTS - ENHANCES SOCIAL ACTIVITY	YES	Count	190	397	3	590
		Expected Count	186.4	401.2	2.4	590.0
		% within EFFECTS - ENHANCES SOCIAL ACTIVITY	32.2%	67.3%	.5%	100.0%
		% within Gender	82.6%	80.2%	100.0%	81.0%
		% of Total	26.1%	54.5%	.4%	81.0%
	2	Count	40	98	0	138
		Expected Count	43.6	93.8	.6	138.0
		% within EFFECTS - ENHANCES SOCIAL ACTIVITY	29.0%	71.0%	.0%	100.0%
		% within Gender	17.4%	19.8%	.0%	19.0%
		% of Total	5.5%	13.5%	.0%	19.0%
Total	Count	230	495	3	728	
	Expected Count	230.0	495.0	3.0	728.0	
	% within EFFECTS - ENHANCES SOCIAL ACTIVITY	31.6%	68.0%	.4%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	31.6%	68.0%	.4%	100.0%	

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1.297(a)	2	.523
Likelihood Ratio	1.862	2	.394
Linear-by-Linear Association	.368	1	.544
N of Valid Cases	728		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .57.

## EFFECTS - EASY TO DEAL WITH STRESS \* Gender

#### Crosstab

			Gender			Total
			Male	Female	Unknown	
EFFECTS - EASY TO DEAL	YES	Count	109	217	2	328
		Expected Count	103.8	222.9	1.3	328.0
		% within EFFECTS - EASY TO DEAL WITH STRESS	33.2%	66.2%	.6%	100.0%
		% within Gender	47.2%	43.8%	66.7%	44.9%
		% of Total	14.9%	29.7%	.3%	44.9%
	2	Count	103	214	2	320
		Expected Count	103.8	222.9	1.3	320.0
		% within EFFECTS - EASY TO DEAL WITH STRESS	32.2%	67.8%	.6%	100.0%
		% within Gender	47.2%	43.8%	66.7%	44.9%
		% of Total	14.9%	29.7%	.3%	44.9%

WITH STRESS	2	Count	122	279	1	402
		Expected Count	127.2	273.1	1.7	402.0
		% within EFFECTS - EASY TO DEAL WITH STRESS	30.3%	69.4%	.2%	100.0%
		% within Gender	52.8%	56.3%	33.3%	55.1%
		% of Total	16.7%	38.2%	.1%	55.1%
Total		Count	231	496	3	730
		Expected Count	231.0	496.0	3.0	730.0
		% within EFFECTS - EASY TO DEAL WITH STRESS	31.6%	67.9%	.4%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.6%	67.9%	.4%	100.0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1.327(a)	2	.515
Likelihood Ratio	1.328	2	.515
Linear-by-Linear Association	.515	1	.473
N of Valid Cases	730		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is 1.35.

## EFFECTS - FACILITATES A CONNECTION WITH PEERS \* Gender

#### Crosstab

			Gender			Total	
			Male	Female	Unknown		
EFFECTS - FACILITATES A CONNECTION WITH PEERS	YES	Count	177	329	2	508	
		Expected Count	160.3	345.6	2.1	508.0	
		% within EFFECTS - FACILITATES A CONNECTION WITH PEERS	34.8%	64.8%	.4%	100.0%	
		% within Gender	77.0%	66.3%	66.7%	69.7%	
		% of Total	24.3%	45.1%	.3%	69.7%	
	2	Count	53	167	1	221	
		Expected Count	69.7	150.4	.9	221.0	
		% within EFFECTS - FACILITATES A CONNECTION WITH PEERS	24.0%	75.6%	.5%	100.0%	
		% within Gender	23.0%	33.7%	33.3%	30.3%	
		% of Total	7.3%	22.9%	.1%	30.3%	
			Count	230	496	3	
						729	

Total	Expected Count	230.0	496.0	3.0	729.0
	% within EFFECTS - FACILITATES A CONNECTION WITH PEERS	31.6%	68.0%	.4%	100.0%
	% within Gender	100.0%	100.0%	100.0%	100.0%
	% of Total	31.6%	68.0%	.4%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	8.411(a)	2	.015
Likelihood Ratio	8.668	2	.013
Linear-by-Linear Association	8.236	1	.004
N of Valid Cases	729		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .91.

## EFFECTS - GIVES PEOPLE SOMETHING TO TALK ABOUT \* Gender

## Crosstab

		Gender			Total
		Male	Female	Unknown	
EFFECTS - GIVES PEOPLE SOMETHING TO TALK ABOUT	YES	Count	185	371	3 559
		Expected Count	177.4	379.3	2.3 559.0
		% within EFFECTS - GIVES PEOPLE SOMETHING TO TALK ABOUT	33.1%	66.4%	.5% 100.0%
		% within Gender	79.7%	74.8%	100.0% 76.5%
		% of Total	25.3%	50.8%	.4% 76.5%
	2	Count	47	125	0 172
		Expected Count	54.6	116.7	.7 172.0
		% within EFFECTS - GIVES PEOPLE SOMETHING TO TALK ABOUT	27.3%	72.7%	.0% 100.0%
		% within Gender	20.3%	25.2%	.0% 23.5%
		% of Total	6.4%	17.1%	.0% 23.5%
Total		Count	232	496	3 731
		Expected Count	232.0	496.0	3.0 731.0
		% within EFFECTS - GIVES PEOPLE SOMETHING TO TALK ABOUT	31.7%	67.9%	.4% 100.0%
		% within Gender	100.0%	100.0%	100.0% 100.0%
		% of Total	31.7%	67.9%	.4% 100.0%

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	3.073(a)	2	.215
<b>Likelihood Ratio</b>	3.796	2	.150
<b>Linear-by-Linear Association</b>	1.610	1	.204
<b>N of Valid Cases</b>	731		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .71.

# **EFFECTS - FACILITATES MALE BONDING \***

## **Gender**

**Crosstab**

			Gender			Total
			Male	Female	Unknown	
EFFECTS - FACILITATES MALE BONDING	YES	Count	162	308	3	473
		Expected Count	150.1	321.0	1.9	473.0
		% within EFFECTS - FACILITATES MALE BONDING	34.2%	65.1%	.6%	100.0%
		% within Gender	70.1%	62.3%	100.0%	65.0%
		% of Total	22.3%	42.3%	.4%	65.0%
	2	Count	69	186	0	255
		Expected Count	80.9	173.0	1.1	255.0
		% within EFFECTS - FACILITATES MALE BONDING	27.1%	72.9%	.0%	100.0%
		% within Gender	29.9%	37.7%	.0%	35.0%
		% of Total	9.5%	25.5%	.0%	35.0%
Total	Count	231	494	3	728	
	Expected Count	231.0	494.0	3.0	728.0	
	% within EFFECTS - FACILITATES MALE BONDING	31.7%	67.9%	.4%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	31.7%	67.9%	.4%	100.0%	

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	5.812(a)	2	.055
<b>Likelihood Ratio</b>	6.835	2	.033
<b>Linear-by-Linear Association</b>	3.184	1	.074
<b>N of Valid Cases</b>	728		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is 1.05.

## EFFECTS - FACILITATES FEMALE BONDING \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
EFFECTS - FACILITATES FEMALE BONDING	YES	Count	140	263	3	406
		Expected Count	128.9	275.5	1.7	406.0
		% within EFFECTS - FACILITATES FEMALE BONDING	34.5%	64.8%	.7%	100.0%
		% within Gender	60.3%	53.0%	100.0%	55.5%
		% of Total	19.2%	36.0%	.4%	55.5%
	2	Count	92	233	0	325
		Expected Count	103.1	220.5	1.3	325.0
		% within EFFECTS - FACILITATES FEMALE BONDING	28.3%	71.7%	.0%	100.0%
		% within Gender	39.7%	47.0%	.0%	44.5%
		% of Total	12.6%	31.9%	.0%	44.5%
Total		Count	232	496	3	731
		Expected Count	232.0	496.0	3.0	731.0
		% within EFFECTS - FACILITATES FEMALE BONDING	31.7%	67.9%	.4%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.7%	67.9%	.4%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	5.842(a)	2	.054
Likelihood Ratio	6.984	2	.030
Linear-by-Linear Association	2.385	1	.122
N of Valid Cases	731		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is 1.33.

## EFFECTS - ALLOWS PEOPLE TO HAVE MORE FUN \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
EFFECTS - ALLOWS PEOPLE TO HAVE MORE FUN	YES	Count	170	315	2	487
		Expected Count	153.9	331.1	2.0	487.0
		% within EFFECTS - ALLOWS PEOPLE TO HAVE MORE FUN	34.9%	64.7%	.4%	100.0%
		% within Gender	73.9%	63.6%	66.7%	66.9%
		% of Total	23.4%	43.3%	.3%	66.9%
	2	Count	60	180	1	241
		Expected Count	76.1	163.9	1.0	241.0
		% within EFFECTS - ALLOWS PEOPLE TO HAVE MORE FUN	24.9%	74.7%	.4%	100.0%
		% within Gender	26.1%	36.4%	33.3%	33.1%
		% of Total	8.2%	24.7%	.1%	33.1%
Total		Count	230	495	3	728
		Expected Count	230.0	495.0	3.0	728.0
		% within EFFECTS - ALLOWS PEOPLE TO HAVE MORE FUN	31.6%	68.0%	.4%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.6%	68.0%	.4%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	7.489(a)	2	.024
Likelihood Ratio	7.667	2	.022
Linear-by-Linear Association	7.248	1	.007
N of Valid Cases	728		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .99.

## EFFECTS - GIVES PEOPLE SOMETHING TO DO \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
YES	YES	Count	201	394	2	597
		Expected Count	190.3	404.3	2.5	597.0
		% within EFFECTS - GIVES PEOPLE SOMETHING TO DO	33.7%	66.0%	.3%	100.0%

EFFECTS - GIVES PEOPLE SOMETHING TO DO	2	% within Gender	86.6%	79.9%	66.7%	82.0%	
		% of Total	27.6%	54.1%	.3%	82.0%	
		Count	31	99	1	131	
		Expected Count	41.7	88.7	.5	131.0	
		% within EFFECTS - GIVES PEOPLE SOMETHING TO DO	23.7%	75.6%	.8%	100.0%	
	Total	% within Gender	13.4%	20.1%	33.3%	18.0%	
		% of Total	4.3%	13.6%	.1%	18.0%	
		Count	232	493	3	728	
		Expected Count	232.0	493.0	3.0	728.0	
		% within EFFECTS - GIVES PEOPLE SOMETHING TO DO	31.9%	67.7%	.4%	100.0%	
		% within Gender	100.0%	100.0%	100.0%	100.0%	
		% of Total	31.9%	67.7%	.4%	100.0%	

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	5.307(a)	2	.070
Likelihood Ratio	5.454	2	.065
Linear-by-Linear Association	5.216	1	.022
N of Valid Cases	728		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .54.

## EFFECTS - MAKES FOOD TASTE BETTER \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
EFFECTS - MAKES FOOD TASTE BETTER	YES	Count	44	84	1	129
		Expected Count	40.9	87.6	.5	129.0
		% within EFFECTS - MAKES FOOD TASTE BETTER	34.1%	65.1%	.8%	100.0%
		% within Gender	19.0%	17.0%	33.3%	17.7%
		% of Total	6.0%	11.5%	.1%	17.7%
	2	Count	187	411	2	600
		Expected Count	190.1	407.4	2.5	600.0
		% within EFFECTS - MAKES FOOD TASTE BETTER	31.2%	68.5%	.3%	100.0%
		% within Gender	81.0%	83.0%	66.7%	82.3%

	% of Total	25.7%	56.4%	.3%	82.3%
Total	Count	231	495	3	729
	Expected Count	231.0	495.0	3.0	729.0
	% within EFFECTS - MAKES FOOD TASTE BETTER	31.7%	67.9%	.4%	100.0%
	% within Gender	100.0%	100.0%	100.0%	100.0%
	% of Total	31.7%	67.9%	.4%	100.0%

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.973(a)	2	.615
Likelihood Ratio	.889	2	.641
Linear-by-Linear Association	.297	1	.586
N of Valid Cases	729		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .53.

## EFFECTS - MAKES WOMEN SEXIER \* Gender

### Crosstab

			Gender			Total
			Male	Female	Unknown	
EFFECTS - MAKES WOMEN SEXIER	YES	Count	117	105	1	223
		Expected Count	71.0	151.1	.9	223.0
		% within EFFECTS - MAKES WOMEN SEXIER	52.5%	47.1%	.4%	100.0%
		% within Gender	50.4%	21.3%	33.3%	30.6%
		% of Total	16.0%	14.4%	.1%	30.6%
	2	Count	115	389	2	506
		Expected Count	161.0	342.9	2.1	506.0
		% within EFFECTS - MAKES WOMEN SEXIER	22.7%	76.9%	.4%	100.0%
		% within Gender	49.6%	78.7%	66.7%	69.4%
		% of Total	15.8%	53.4%	.3%	69.4%
Total	Count	232	494	3	729	
	Expected Count	232.0	494.0	3.0	729.0	
	% within EFFECTS - MAKES WOMEN SEXIER	31.8%	67.8%	.4%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	31.8%	67.8%	.4%	100.0%	

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	63.300(a)	2	.000
Likelihood Ratio	61.276	2	.000
Linear-by-Linear Association	60.898	1	.000
N of Valid Cases	729		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .92.

## EFFECTS - MAKES MEN SEXIER \* Gender

## Crosstab

		Gender			Total
		Male	Female	Unknown	
EFFECTS - MAKES MEN SEXIER	YES	Count	43	125	0 168
		Expected Count	53.1	114.2	.7 168.0
		% within EFFECTS - MAKES MEN SEXIER	25.6%	74.4%	.0% 100.0%
		% within Gender	18.7%	25.3%	.0% 23.1%
		% of Total	5.9%	17.2%	.0% 23.1%
	2	Count	187	370	3 560
		Expected Count	176.9	380.8	2.3 560.0
		% within EFFECTS - MAKES MEN SEXIER	33.4%	66.1%	.5% 100.0%
		% within Gender	81.3%	74.7%	100.0% 76.9%
		% of Total	25.7%	50.8%	.4% 76.9%
Total		Count	230	495	3 728
		Expected Count	230.0	495.0	3.0 728.0
		% within EFFECTS - MAKES MEN SEXIER	31.6%	68.0%	.4% 100.0%
		% within Gender	100.0%	100.0%	100.0% 100.0%
		% of Total	31.6%	68.0%	.4% 100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	4.707(a)	2	.095
Likelihood Ratio	5.477	2	.065
Linear-by-Linear Association	3.054	1	.081
N of Valid Cases	728		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .69.

# EFFECTS - MAKES ME SEXIER \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
EFFECTS - MAKES ME SEXIER	YES	Count	50	89	1	140
		Expected Count	44.5	94.9	.6	140.0
		% within EFFECTS - MAKES ME SEXIER	35.7%	63.6%	.7%	100.0%
		% within Gender	21.6%	18.1%	33.3%	19.3%
		% of Total	6.9%	12.2%	.1%	19.3%
	2	Count	181	404	2	587
		Expected Count	186.5	398.1	2.4	587.0
		% within EFFECTS - MAKES ME SEXIER	30.8%	68.8%	.3%	100.0%
		% within Gender	78.4%	81.9%	66.7%	80.7%
		% of Total	24.9%	55.6%	.3%	80.7%
Total	Count	231	493	3	727	
	Expected Count	231.0	493.0	3.0	727.0	
	% within EFFECTS - MAKES ME SEXIER	31.8%	67.8%	.4%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	31.8%	67.8%	.4%	100.0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1.689(a)	2	.430
Likelihood Ratio	1.621	2	.445
Linear-by-Linear Association	1.025	1	.311
N of Valid Cases	727		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .58.

# EFFECTS - FACILITATES SEXUAL OPPORTUNITIES \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
	Count	138	247	1	386	

EFFECTS - FACILITATES SEXUAL OPPORTUNITIES	YES	Expected Count	122.5	261.9	1.6	386.0
		% within EFFECTS - FACILITATES SEXUAL OPPORTUNITIES	35.8%	64.0%	.3%	100.0%
		% within Gender	59.5%	49.8%	33.3%	52.8%
		% of Total	18.9%	33.8%	.1%	52.8%
	2	Count	94	249	2	345
		Expected Count	109.5	234.1	1.4	345.0
		% within EFFECTS - FACILITATES SEXUAL OPPORTUNITIES	27.2%	72.2%	.6%	100.0%
		% within Gender	40.5%	50.2%	66.7%	47.2%
		% of Total	12.9%	34.1%	.3%	47.2%
	Total	Count	232	496	3	731
		Expected Count	232.0	496.0	3.0	731.0
		% within EFFECTS - FACILITATES SEXUAL OPPORTUNITIES	31.7%	67.9%	.4%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.7%	67.9%	.4%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	6.407(a)	2	.041
Likelihood Ratio	6.443	2	.040
Linear-by-Linear Association	6.345	1	.012
N of Valid Cases	731		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is 1.42.

# DRINKING IS CENTRAL - MALE STUDENTS

## \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
DRINKING IS CENTRAL - MALE STUDENTS	YES	Count	214	474	2	690
		Expected Count	218.9	469.2	1.9	690.0
		% within DRINKING IS CENTRAL - MALE STUDENTS	31.0%	68.7%	.3%	100.0%
		% within Gender	93.0%	96.1%	100.0%	95.2%
		% of Total	29.5%	65.4%	.3%	95.2%
		Count	16	19	0	35
		Expected Count	11.1	23.8	.1	35.0

		% within DRINKING IS CENTRAL - MALE STUDENTS	45.7%	54.3%	.0%	100.0%
	2	% within Gender	7.0%	3.9%	.0%	4.8%
		% of Total	2.2%	2.6%	.0%	4.8%
Total		Count	230	493	2	725
		Expected Count	230.0	493.0	2.0	725.0
		% within DRINKING IS CENTRAL - MALE STUDENTS	31.7%	68.0%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.7%	68.0%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	3.388(a)	2	.184
Likelihood Ratio	3.295	2	.192
Linear-by-Linear Association	3.380	1	.066
N of Valid Cases	725		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .10.

## DRINKING IS CENTRAL - FEMALE STUDENTS \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
DRINKING IS CENTRAL - FEMALE STUDENTS	YES	Count	196	440	1	637
		Expected Count	200.9	434.4	1.8	637.0
		% within DRINKING IS CENTRAL - FEMALE STUDENTS	30.8%	69.1%	.2%	100.0%
		% within Gender	86.0%	89.2%	50.0%	88.1%
		% of Total	27.1%	60.9%	.1%	88.1%
	2	Count	32	53	1	86
		Expected Count	27.1	58.6	.2	86.0
		% within DRINKING IS CENTRAL - FEMALE STUDENTS	37.2%	61.6%	1.2%	100.0%
		% within Gender	14.0%	10.8%	50.0%	11.9%
		% of Total	4.4%	7.3%	.1%	11.9%
Total	Count	228	493	2	723	
	Expected Count	228.0	493.0	2.0	723.0	
	% within DRINKING IS CENTRAL -	31.5%	68.2%	.3%	100.0%	

	FEMALE STUDENTS				
	% within Gender	100.0%	100.0%	100.0%	100.0%
	% of Total	31.5%	68.2%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	4.384(a)	2	.112
Likelihood Ratio	3.322	2	.190
Linear-by-Linear Association	1.014	1	.314
N of Valid Cases	723		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .24.

# DRINKING IS CENTRAL - FACULTY \*

## Gender

## Crosstab

		Gender			Total
		Male	Female	Unknown	
DRINKING IS CENTRAL - FACULTY	YES	Count	39	65	0 104
		Expected Count	32.7	71.0	.3 104.0
		% within DRINKING IS CENTRAL - FACULTY	37.5%	62.5%	.0% 100.0%
		% within Gender	17.3%	13.2%	.0% 14.5%
		% of Total	5.4%	9.0%	.0% 14.5%
	2	Count	187	426	2 615
		Expected Count	193.3	420.0	1.7 615.0
		% within DRINKING IS CENTRAL - FACULTY	30.4%	69.3%	.3% 100.0%
		% within Gender	82.7%	86.8%	100.0% 85.5%
		% of Total	26.0%	59.2%	.3% 85.5%
Total		Count	226	491	2 719
		Expected Count	226.0	491.0	2.0 719.0
		% within DRINKING IS CENTRAL - FACULTY	31.4%	68.3%	.3% 100.0%
		% within Gender	100.0%	100.0%	100.0% 100.0%
		% of Total	31.4%	68.3%	.3% 100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	2.359(a)	2	.307

Likelihood Ratio	2.591	2	.274
Linear-by-Linear Association	2.222	1	.136
N of Valid Cases	719		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .29.

## DRINKING IS CENTRAL - ALUMNI \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
DRINKING IS CENTRAL - ALUMNI	YES	Count	83	122	0	205
		Expected Count	64.7	139.7	.6	205.0
		% within DRINKING IS CENTRAL - ALUMNI	40.5%	59.5%	.0%	100.0%
		% within Gender	36.6%	24.9%	.0%	28.5%
		% of Total	11.5%	17.0%	.0%	28.5%
	2	Count	144	368	2	514
		Expected Count	162.3	350.3	1.4	514.0
		% within DRINKING IS CENTRAL - ALUMNI	28.0%	71.6%	.4%	100.0%
		% within Gender	63.4%	75.1%	100.0%	71.5%
		% of Total	20.0%	51.2%	.3%	71.5%
Total		Count	227	490	2	719
		Expected Count	227.0	490.0	2.0	719.0
		% within DRINKING IS CENTRAL - ALUMNI	31.6%	68.2%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.6%	68.2%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	11.158(a)	2	.004
Likelihood Ratio	11.444	2	.003
Linear-by-Linear Association	10.975	1	.001
N of Valid Cases	719		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .57.

## DRINKING IS CENTRAL -ATHLETES \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
DRINKING IS CENTRAL - ATHLETES	YES	Count	163	352	1	516
		Expected Count	164.6	350.0	1.4	516.0
		% within DRINKING IS CENTRAL - ATHLETES	31.6%	68.2%	.2%	100.0%
		% within Gender	71.2%	72.3%	50.0%	71.9%
		% of Total	22.7%	49.0%	.1%	71.9%
	2	Count	66	135	1	202
		Expected Count	64.4	137.0	.6	202.0
		% within DRINKING IS CENTRAL - ATHLETES	32.7%	66.8%	.5%	100.0%
		% within Gender	28.8%	27.7%	50.0%	28.1%
		% of Total	9.2%	18.8%	.1%	28.1%
Total	Count	229	487	2	718	
	Expected Count	229.0	487.0	2.0	718.0	
	% within DRINKING IS CENTRAL - ATHLETES	31.9%	67.8%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	31.9%	67.8%	.3%	100.0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.568(a)	2	.753
Likelihood Ratio	.519	2	.771
Linear-by-Linear Association	.040	1	.841
N of Valid Cases	718		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .56.

## DRINKING IS CENTRAL - FRATERNITIES \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
YES	Count	219	487	2	708	
	Expected Count	223.9	482.1	2.0	708.0	
	% within DRINKING IS CENTRAL -	30.9%	68.8%	.3%	100.0%	

		FRATERNITIES			
		% within Gender	95.6%	98.8%	100.0%
		% of Total	30.2%	67.3%	.3%
DRINKING IS CENTRAL - FRATERNITIES	2	Count	10	6	0
		Expected Count	5.1	10.9	.0
		% within DRINKING IS CENTRAL - FRATERNITIES	62.5%	37.5%	.0%
		% within Gender	4.4%	1.2%	.0%
		% of Total	1.4%	.8%	.0%
Total		Count	229	493	2
		Expected Count	229.0	493.0	2.0
		% within DRINKING IS CENTRAL - FRATERNITIES	31.6%	68.1%	.3%
		% within Gender	100.0%	100.0%	100.0%
		% of Total	31.6%	68.1%	.3%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	7.224(a)	2	.027
Likelihood Ratio	6.623	2	.036
Linear-by-Linear Association	7.180	1	.007
N of Valid Cases	724		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .04.

## DRINKING IS CENTRAL - SORORITIES \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
DRINKING IS CENTRAL - SORORITIES	YES	Count	209	453	2	664
		Expected Count	210.9	451.2	1.8	664.0
		% within DRINKING IS CENTRAL - SORORITIES	31.5%	68.2%	.3%	100.0%
		% within Gender	90.9%	92.1%	100.0%	91.7%
		% of Total	28.9%	62.6%	.3%	91.7%
	2	Count	21	39	0	60
		Expected Count	19.1	40.8	.2	60.0
		% within DRINKING IS CENTRAL - SORORITIES	35.0%	65.0%	.0%	100.0%

	% within Gender	9.1%	7.9%	.0%	8.3%
	% of Total	2.9%	5.4%	.0%	8.3%
Total	Count	230	492	2	724
	Expected Count	230.0	492.0	2.0	724.0
	% within DRINKING IS CENTRAL - SORORITIES	31.8%	68.0%	.3%	100.0%
	% within Gender	100.0%	100.0%	100.0%	100.0%
	% of Total	31.8%	68.0%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.480(a)	2	.787
Likelihood Ratio	.640	2	.726
Linear-by-Linear Association	.363	1	.547
N of Valid Cases	724		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .17.

## CAMPUS ENVIRONMENT:PROMOTE ALCOHOL USE? \* Gender

## Crosstab

		Gender			Total
		Male	Female	Unknown	
CAMPUS ENVIRONMENT:PROMOTE ALCOHOL USE?	YES	Count	179	339	2 520
		Expected Count	165.7	352.9	1.4 520.0
		% within CAMPUS ENVIRONMENT:PROMOTE ALCOHOL USE?	34.4%	65.2%	.4% 100.0%
		% within Gender	77.5%	68.9%	100.0% 71.7%
		% of Total	24.7%	46.8%	.3% 71.7%
	2	Count	52	153	0 205
		Expected Count	65.3	139.1	.6 205.0
		% within CAMPUS ENVIRONMENT:PROMOTE ALCOHOL USE?	25.4%	74.6%	.0% 100.0%
		% within Gender	22.5%	31.1%	.0% 28.3%
		% of Total	7.2%	21.1%	.0% 28.3%
Total	Count	231	492	2	725
	Expected Count	231.0	492.0	2.0	725.0
	% within CAMPUS ENVIRONMENT:PROMOTE	31.9%	67.9%	.3%	100.0%

	ALCOHOL USE?				
	% within Gender	100.0%	100.0%	100.0%	100.0%
	% of Total	31.9%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	6.506(a)	2	.039
Likelihood Ratio	7.188	2	.027
Linear-by-Linear Association	4.983	1	.026
N of Valid Cases	725		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .57.

# CAMPUS ENVIRONMENT:PROMOTE OTHER DRUG USE? \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
CAMPUS ENVIRONMENT:PROMOTE OTHER DRUG USE?	YES	Count	67	109	0	176
		Expected Count	56.0	119.5	.5	176.0
		% within CAMPUS ENVIRONMENT:PROMOTE OTHER DRUG USE?	38.1%	61.9%	.0%	100.0%
		% within Gender	29.0%	22.1%	.0%	24.2%
		% of Total	9.2%	15.0%	.0%	24.2%
	2	Count	164	384	2	550
		Expected Count	175.0	373.5	1.5	550.0
		% within CAMPUS ENVIRONMENT:PROMOTE OTHER DRUG USE?	29.8%	69.8%	.4%	100.0%
		% within Gender	71.0%	77.9%	100.0%	75.8%
		% of Total	22.6%	52.9%	.3%	75.8%
Total		Count	231	493	2	726
		Expected Count	231.0	493.0	2.0	726.0
		% within CAMPUS ENVIRONMENT:PROMOTE OTHER DRUG USE?	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	4.713(a)	2	.095
<b>Likelihood Ratio</b>	5.095	2	.078
<b>Linear-by-Linear Association</b>	4.461	1	.035
<b>N of Valid Cases</b>	726		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .48.

## CAMPUS ENVIRONEMNT:DO YOU FEEL SAFE ON CAMPUS? \* Gender

Crosstab

		Gender			Total
		Male	Female	Unknown	
CAMPUS ENVIRONEMNT:DO YOU FEEL SAFE ON CAMPUS?	YES	Count	225	464	2 691
		Expected Count	219.6	469.5	1.9 691.0
		% within CAMPUS ENVIRONEMNT:DO YOU FEEL SAFE ON CAMPUS?	32.6%	67.1%	.3% 100.0%
		% within Gender	97.4%	93.9%	100.0% 95.0%
		% of Total	30.9%	63.8%	.3% 95.0%
	2	Count	6	30	0 36
		Expected Count	11.4	24.5	.1 36.0
		% within CAMPUS ENVIRONEMNT:DO YOU FEEL SAFE ON CAMPUS?	16.7%	83.3%	.0% 100.0%
		% within Gender	2.6%	6.1%	.0% 5.0%
		% of Total	.8%	4.1%	.0% 5.0%
Total		Count	231	494	2 727
		Expected Count	231.0	494.0	2.0 727.0
		% within CAMPUS ENVIRONEMNT:DO YOU FEEL SAFE ON CAMPUS?	31.8%	68.0%	.3% 100.0%
		% within Gender	100.0%	100.0%	100.0% 100.0%
		% of Total	31.8%	68.0%	.3% 100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	4.144(a)	2	.126
<b>Likelihood Ratio</b>	4.705	2	.095
<b>Linear-by-Linear Association</b>	3.761	1	.052
<b>N of Valid Cases</b>	727		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .10.

# COMPARED TO OTHER CAMPUSES:ALCOHOL USE IS \* Gender

Crosstab

		Gender			Total
		Male	Female	Unknown	
COMPARED TO OTHER CAMPUSES:ALCOHOL USE IS	GREATER THAN OTHER CAMPUSES	Count	44	53	1 98
		Expected Count	31.2	66.5	.3 98.0
		% within COMPARED TO OTHER CAMPUSES:ALCOHOL USE IS	44.9%	54.1%	1.0% 100.0%
		% within Gender	19.0%	10.8%	50.0% 13.5%
		% of Total	6.1%	7.3%	.1% 13.5%
	LESS THAN OTHER CAMPUSES	Count	44	71	0 115
		Expected Count	36.6	78.1	.3 115.0
		% within COMPARED TO OTHER CAMPUSES:ALCOHOL USE IS	38.3%	61.7%	.0% 100.0%
		% within Gender	19.0%	14.4%	.0% 15.8%
		% of Total	6.1%	9.8%	.0% 15.8%
	ABOUT THE SAME AS OTHER CAMPUSES	Count	143	369	1 513
		Expected Count	163.2	348.4	1.4 513.0
		% within COMPARED TO OTHER CAMPUSES:ALCOHOL USE IS	27.9%	71.9%	.2% 100.0%
		% within Gender	61.9%	74.8%	50.0% 70.7%
		% of Total	19.7%	50.8%	.1% 70.7%
Total	Count	231	493	2	726
	Expected Count	231.0	493.0	2.0	726.0
	% within COMPARED TO OTHER CAMPUSES:ALCOHOL USE IS	31.8%	67.9%	.3%	100.0%
	% within Gender	100.0%	100.0%	100.0%	100.0%
	% of Total	31.8%	67.9%	.3%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)

Pearson Chi-Square	16.313(a)	4	.003
Likelihood Ratio	15.450	4	.004
Linear-by-Linear Association	12.279	1	.000
N of Valid Cases	726		

a 3 cells (33.3%) have expected count less than 5. The minimum expected count is .27.

## DO YOU LIVE IN ALCOHOL FREE RESIDENCE HALL \* Gender

Crosstab

		Gender			Total
		Male	Female	Unknown	
DO YOU LIVE IN ALCOHOL FREE RESIDENCE HALL	YES	Count	58	159	1 218
		Expected Count	70.8	146.3	.8 218.0
		% within DO YOU LIVE IN ALCOHOL FREE RESIDENCE HALL	26.6%	72.9%	.5% 100.0%
		% within Gender	34.7%	46.1%	50.0% 42.4%
		% of Total	11.3%	30.9%	.2% 42.4%
	2	Count	109	186	1 296
		Expected Count	96.2	198.7	1.2 296.0
		% within DO YOU LIVE IN ALCOHOL FREE RESIDENCE HALL	36.8%	62.8%	.3% 100.0%
		% within Gender	65.3%	53.9%	50.0% 57.6%
		% of Total	21.2%	36.2%	.2% 57.6%
Total		Count	167	345	2 514
		Expected Count	167.0	345.0	2.0 514.0
		% within DO YOU LIVE IN ALCOHOL FREE RESIDENCE HALL	32.5%	67.1%	.4% 100.0%
		% within Gender	100.0%	100.0%	100.0% 100.0%
		% of Total	32.5%	67.1%	.4% 100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	5.989(a)	2	.050
Likelihood Ratio	6.059	2	.048
Linear-by-Linear Association	5.934	1	.015
N of Valid Cases	514		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .85.

# WOULD YOU LIVE IN ALCOHOL FREE RESIDENCE HALL \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
WOULD YOU LIVE IN ALCOHOL FREE RESIDENCE HALL	YES	Count	24	57	0	81
		Expected Count	28.1	52.7	.2	81.0
		% within WOULD YOU LIVE IN ALCOHOL FREE RESIDENCE HALL	29.6%	70.4%	.0%	100.0%
		% within Gender	17.0%	21.5%	.0%	19.9%
		% of Total	5.9%	14.0%	.0%	19.9%
	2	Count	117	208	1	326
		Expected Count	112.9	212.3	.8	326.0
		% within WOULD YOU LIVE IN ALCOHOL FREE RESIDENCE HALL	35.9%	63.8%	.3%	100.0%
		% within Gender	83.0%	78.5%	100.0%	80.1%
		% of Total	28.7%	51.1%	.2%	80.1%
Total		Count	141	265	1	407
		Expected Count	141.0	265.0	1.0	407.0
		% within WOULD YOU LIVE IN ALCOHOL FREE RESIDENCE HALL	34.6%	65.1%	.2%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	34.6%	65.1%	.2%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1.412(a)	2	.494
Likelihood Ratio	1.628	2	.443
Linear-by-Linear Association	.995	1	.319
N of Valid Cases	407		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .20.

# STUDENTS CARE : ALCOHOL -DRUG USE \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
STUDENTS CARE : ALCOHOL -DRUG USE	NOT AT ALL	Count	40	71	0	111
		Expected Count	35.2	75.5	.3	111.0
		% within STUDENTS CARE : ALCOHOL -DRUG USE	36.0%	64.0%	.0%	100.0%
		% within Gender	17.5%	14.5%	.0%	15.4%
		% of Total	5.5%	9.8%	.0%	15.4%
	SLIGHTLY	Count	117	198	2	317
		Expected Count	100.5	215.6	.9	317.0
		% within STUDENTS CARE : ALCOHOL -DRUG USE	36.9%	62.5%	.6%	100.0%
		% within Gender	51.1%	40.3%	100.0%	43.9%
		% of Total	16.2%	27.4%	.3%	43.9%
	SOMEWHAT	Count	65	207	0	272
		Expected Count	86.3	185.0	.8	272.0
		% within STUDENTS CARE : ALCOHOL -DRUG USE	23.9%	76.1%	.0%	100.0%
		% within Gender	28.4%	42.2%	.0%	37.7%
		% of Total	9.0%	28.7%	.0%	37.7%
	VERY MUCH	Count	7	15	0	22
		Expected Count	7.0	15.0	.1	22.0
		% within STUDENTS CARE : ALCOHOL -DRUG USE	31.8%	68.2%	.0%	100.0%
		% within Gender	3.1%	3.1%	.0%	3.0%
		% of Total	1.0%	2.1%	.0%	3.0%
Total	Count	229	491	2	722	
	Expected Count	229.0	491.0	2.0	722.0	
	% within STUDENTS CARE : ALCOHOL -DRUG USE	31.7%	68.0%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	31.7%	68.0%	.3%	100.0%	

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	15.468(a)	6	.017
Likelihood Ratio	16.483	6	.011
Linear-by-Linear Association	7.089	1	.008
N of Valid Cases	722		

a 4 cells (33.3%) have expected count less than 5. The minimum expected count is .06.

# STUDENTS CARE : CAMPUS VANDALISM \*

## Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
STUDENTS CARE : CAMPUS VANDALISM	NOT AT ALL	Count	34	58	0	92
		Expected Count	29.0	62.7	.3	92.0
		% within STUDENTS CARE : CAMPUS VANDALISM	37.0%	63.0%	.0%	100.0%
		% within Gender	15.0%	11.8%	.0%	12.8%
		% of Total	4.7%	8.1%	.0%	12.8%
	SLIGHTLY	Count	75	148	0	223
		Expected Count	70.4	152.0	.6	223.0
		% within STUDENTS CARE : CAMPUS VANDALISM	33.6%	66.4%	.0%	100.0%
		% within Gender	33.0%	30.2%	.0%	31.0%
		% of Total	10.4%	20.6%	.0%	31.0%
	SOMEWHAT	Count	87	225	1	313
		Expected Count	98.8	213.3	.9	313.0
		% within STUDENTS CARE : CAMPUS VANDALISM	27.8%	71.9%	.3%	100.0%
		% within Gender	38.3%	45.9%	50.0%	43.5%
		% of Total	12.1%	31.3%	.1%	43.5%
	VERY MUCH	Count	31	59	1	91
		Expected Count	28.7	62.0	.3	91.0
		% within STUDENTS CARE : CAMPUS VANDALISM	34.1%	64.8%	1.1%	100.0%
		% within Gender	13.7%	12.0%	50.0%	12.7%
		% of Total	4.3%	8.2%	.1%	12.7%
Total	Count	227	490	2	719	
	Expected Count	227.0	490.0	2.0	719.0	
	% within STUDENTS CARE : CAMPUS VANDALISM	31.6%	68.2%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	31.6%	68.2%	.3%	100.0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	7.080(a)	6	.313

Likelihood Ratio	7.012	6	.320
Linear-by-Linear Association	1.660	1	.198
N of Valid Cases	719		

a 4 cells (33.3%) have expected count less than 5. The minimum expected count is .25.

# STUDENTS CARE : SEXUAL ASSAULT \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
STUDENTS CARE : SEXUAL ASSAULT	NOT AT ALL	Count	7	7	0	14
		Expected Count	4.4	9.5	.0	14.0
		% within STUDENTS CARE : SEXUAL ASSAULT	50.0%	50.0%	.0%	100.0%
		% within Gender	3.1%	1.4%	.0%	1.9%
		% of Total	1.0%	1.0%	.0%	1.9%
	SLIGHTLY	Count	20	54	0	74
		Expected Count	23.4	50.4	.2	74.0
		% within STUDENTS CARE : SEXUAL ASSAULT	27.0%	73.0%	.0%	100.0%
		% within Gender	8.8%	11.0%	.0%	10.3%
		% of Total	2.8%	7.5%	.0%	10.3%
	SOMEWHAT	Count	73	151	0	224
		Expected Count	70.8	152.6	.6	224.0
		% within STUDENTS CARE : SEXUAL ASSAULT	32.6%	67.4%	.0%	100.0%
		% within Gender	32.2%	30.9%	.0%	31.2%
		% of Total	10.2%	21.0%	.0%	31.2%
	VERY MUCH	Count	127	277	2	406
		Expected Count	128.4	276.5	1.1	406.0
		% within STUDENTS CARE : SEXUAL ASSAULT	31.3%	68.2%	.5%	100.0%
		% within Gender	55.9%	56.6%	100.0%	56.5%
		% of Total	17.7%	38.6%	.3%	56.5%
Total		Count	227	489	2	718
		Expected Count	227.0	489.0	2.0	718.0
		% within STUDENTS CARE : SEXUAL ASSAULT	31.6%	68.1%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%

	% of Total	31.6%	68.1%	.3%	100.0%
--	------------	-------	-------	-----	--------

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	4.556(a)	6	.602
Likelihood Ratio	5.161	6	.523
Linear-by-Linear Association	.202	1	.653
N of Valid Cases	718		

a 5 cells (41.7%) have expected count less than 5. The minimum expected count is .04.

# STUDENTS CARE : ASSUALTS-NON SEXUAL \* Gender

## Crosstab

		Gender			Total
		Male	Female	Unknown	
STUDENTS CARE : ASSUALTS-NON SEXUAL	NOT AT ALL	Count	12	20	0 32
		Expected Count	10.2	21.7	.1 32.0
		% within STUDENTS CARE : ASSUALTS-NON SEXUAL	37.5%	62.5%	.0% 100.0%
		% within Gender	5.2%	4.1%	.0% 4.4%
		% of Total	1.7%	2.8%	.0% 4.4%
	SLIGHTLY	Count	43	85	0 128
		Expected Count	40.7	87.0	.4 128.0
		% within STUDENTS CARE : ASSUALTS-NON SEXUAL	33.6%	66.4%	.0% 100.0%
		% within Gender	18.8%	17.3%	.0% 17.8%
		% of Total	6.0%	11.8%	.0% 17.8%
	SOMEWHAT	Count	97	215	1 313
		Expected Count	99.4	212.7	.9 313.0
		% within STUDENTS CARE : ASSUALTS-NON SEXUAL	31.0%	68.7%	.3% 100.0%
		% within Gender	42.4%	43.9%	50.0% 43.4%
		% of Total	13.5%	29.8%	.1% 43.4%
	VERY MUCH	Count	77	170	1 248
		Expected Count	78.8	168.5	.7 248.0
		% within STUDENTS CARE : ASSUALTS-NON SEXUAL	31.0%	68.5%	.4% 100.0%
		% within Gender	33.6%	34.7%	50.0% 34.4%
		% of Total	10.7%	23.6%	.1% 34.4%

Total	Count	229	490	2	721
	Expected Count	229.0	490.0	2.0	721.0
	% within STUDENTS CARE : ASSUALTS-NON SEXUAL	31.8%	68.0%	.3%	100.0%
	% within Gender	100.0%	100.0%	100.0%	100.0%
	% of Total	31.8%	68.0%	.3%	100.0%

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1.394(a)	6	.966
Likelihood Ratio	1.803	6	.937
Linear-by-Linear Association	.673	1	.412
N of Valid Cases	721		

a 4 cells (33.3%) have expected count less than 5. The minimum expected count is .09.

## STUDENTS CARE : HARASSMENT - GENDER \* Gender

### Crosstab

			Gender			Total
			Male	Female	Unknown	
STUDENTS CARE : HARASSMENT - GENDER	NOT AT ALL	Count	27	39	0	66
		Expected Count	21.0	44.8	.2	66.0
		% within STUDENTS CARE : HARASSMENT - GENDER	40.9%	59.1%	.0%	100.0%
		% within Gender	11.8%	8.0%	.0%	9.2%
		% of Total	3.8%	5.4%	.0%	9.2%
	SLIGHTLY	Count	45	120	0	165
		Expected Count	52.6	112.0	.5	165.0
		% within STUDENTS CARE : HARASSMENT - GENDER	27.3%	72.7%	.0%	100.0%
		% within Gender	19.7%	24.6%	.0%	22.9%
		% of Total	6.3%	16.7%	.0%	22.9%
	SOMEWHAT	Count	86	206	2	294
		Expected Count	93.6	199.5	.8	294.0
		% within STUDENTS CARE : HARASSMENT - GENDER	29.3%	70.1%	.7%	100.0%
		% within Gender	37.6%	42.2%	100.0%	40.9%
		% of Total	12.0%	28.7%	.3%	40.9%
		Count	71	123	0	194

	VERY MUCH	Expected Count	61.8	131.7	.5	194.0
		% within STUDENTS CARE : HARASSMENT - GENDER	36.6%	63.4%	.0%	100.0%
		% within Gender	31.0%	25.2%	.0%	27.0%
		% of Total	9.9%	17.1%	.0%	27.0%
Total	Count	229	488	2	719	
	Expected Count	229.0	488.0	2.0	719.0	
	% within STUDENTS CARE : HARASSMENT - GENDER	31.8%	67.9%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	31.8%	67.9%	.3%	100.0%	

### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	9.776(a)	6	.134
Likelihood Ratio	10.363	6	.110
Linear-by-Linear Association	.151	1	.697
N of Valid Cases	719		

a 4 cells (33.3%) have expected count less than 5. The minimum expected count is .18.

## STUDENTS CARE : HARASSMENT - SEXUAL ORIENTATION \* Gender

### Crosstab

			Gender			Total
			Male	Female	Unknown	
STUDENTS CARE : HARASSMENT - SEXUAL ORIENTATION	NOT AT ALL	Count	26	24	0	50
		Expected Count	15.9	33.9	.1	50.0
		% within STUDENTS CARE : HARASSMENT - SEXUAL ORIENTATION	52.0%	48.0%	.0%	100.0%
		% within Gender	11.4%	4.9%	.0%	7.0%
		% of Total	3.6%	3.3%	.0%	7.0%
	SLIGHTLY	Count	44	118	0	162
		Expected Count	51.6	110.0	.5	162.0
		% within STUDENTS CARE : HARASSMENT - SEXUAL ORIENTATION	27.2%	72.8%	.0%	100.0%
		% within Gender	19.2%	24.2%	.0%	22.5%
		% of Total	6.1%	16.4%	.0%	22.5%
		Count	84	216	1	301

		Expected Count	95.9	204.3	.8	301.0
	SOMEWHAT	% within STUDENTS CARE : HARASSMENT - SEXUAL ORIENTATION	27.9%	71.8%	.3%	100.0%
		% within Gender	36.7%	44.3%	50.0%	41.9%
		% of Total	11.7%	30.0%	.1%	41.9%
	VERY MUCH	Count	75	130	1	206
		Expected Count	65.6	139.8	.6	206.0
		% within STUDENTS CARE : HARASSMENT - SEXUAL ORIENTATION	36.4%	63.1%	.5%	100.0%
		% within Gender	32.8%	26.6%	50.0%	28.7%
		% of Total	10.4%	18.1%	.1%	28.7%
Total		Count	229	488	2	719
		Expected Count	229.0	488.0	2.0	719.0
		% within STUDENTS CARE : HARASSMENT - SEXUAL ORIENTATION	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	16.103(a)	6	.013
Likelihood Ratio	15.988	6	.014
Linear-by-Linear Association	.149	1	.699
N of Valid Cases	719		

a 4 cells (33.3%) have expected count less than 5. The minimum expected count is .14.

## STUDENTS CARE : HARASSMENT - RACE OR ETHNICITY \* Gender

## Crosstab

		Gender			Total	
		Male	Female	Unknown		
NOT AT ALL		Count	18	24	0	42
		Expected Count	13.3	28.6	.1	42.0
		% within STUDENTS CARE : HARASSMENT - RACE OR ETHNICITY	42.9%	57.1%	.0%	100.0%
		% within Gender	7.9%	4.9%	.0%	5.8%
		% of Total	2.5%	3.3%	.0%	5.8%

STUDENTS CARE : HARASSMENT - RACE OR ETHNICITY	SLIGHTLY	Count	31	86	1	118
		Expected Count	37.3	80.4	.3	118.0
		% within STUDENTS CARE : HARASSMENT - RACE OR ETHNICITY	26.3%	72.9%	.8%	100.0%
		% within Gender	13.7%	17.6%	50.0%	16.4%
		% of Total	4.3%	12.0%	.1%	16.4%
	SOMEWHAT	Count	82	196	1	279
		Expected Count	88.1	190.1	.8	279.0
		% within STUDENTS CARE : HARASSMENT - RACE OR ETHNICITY	29.4%	70.3%	.4%	100.0%
		% within Gender	36.1%	40.0%	50.0%	38.8%
		% of Total	11.4%	27.3%	.1%	38.8%
	VERY MUCH	Count	96	184	0	280
		Expected Count	88.4	190.8	.8	280.0
		% within STUDENTS CARE : HARASSMENT - RACE OR ETHNICITY	34.3%	65.7%	.0%	100.0%
		% within Gender	42.3%	37.6%	.0%	38.9%
		% of Total	13.4%	25.6%	.0%	38.9%
Total	Count	227	490	2	719	
	Expected Count	227.0	490.0	2.0	719.0	
	% within STUDENTS CARE : HARASSMENT - RACE OR ETHNICITY	31.6%	68.2%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	31.6%	68.2%	.3%	100.0%	

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	7.712(a)	6	.260
Likelihood Ratio	8.032	6	.236
Linear-by-Linear Association	.254	1	.614
N of Valid Cases	719		

a 4 cells (33.3%) have expected count less than 5. The minimum expected count is .12.

## STUDENTS CARE : HARASSMENT - RELIGION \* Gender

## Crosstab

	Gender
--	--------

			Male	Female	Unknown	Total	
STUDENTS CARE : HARASSMENT - RELIGION	NOT AT ALL	Count	40	64	0	104	
		Expected Count	32.9	70.8	.3	104.0	
		% within STUDENTS CARE : HARASSMENT - RELIGION	38.5%	61.5%	.0%	100.0%	
		% within Gender	17.5%	13.1%	.0%	14.4%	
		% of Total	5.6%	8.9%	.0%	14.4%	
	SLIGHTLY	Count	57	128	1	186	
		Expected Count	58.9	126.6	.5	186.0	
		% within STUDENTS CARE : HARASSMENT - RELIGION	30.6%	68.8%	.5%	100.0%	
		% within Gender	25.0%	26.1%	50.0%	25.8%	
		% of Total	7.9%	17.8%	.1%	25.8%	
	SOMEWHAT	Count	76	195	1	272	
		Expected Count	86.1	185.1	.8	272.0	
		% within STUDENTS CARE : HARASSMENT - RELIGION	27.9%	71.7%	.4%	100.0%	
		% within Gender	33.3%	39.8%	50.0%	37.8%	
		% of Total	10.6%	27.1%	.1%	37.8%	
	VERY MUCH	Count	55	103	0	158	
		Expected Count	50.0	107.5	.4	158.0	
		% within STUDENTS CARE : HARASSMENT - RELIGION	34.8%	65.2%	.0%	100.0%	
		% within Gender	24.1%	21.0%	.0%	21.9%	
		% of Total	7.6%	14.3%	.0%	21.9%	
Total			Count	228	490	2	720
			Expected Count	228.0	490.0	2.0	720.0
			% within STUDENTS CARE : HARASSMENT - RELIGION	31.7%	68.1%	.3%	100.0%
			% within Gender	100.0%	100.0%	100.0%	100.0%
			% of Total	31.7%	68.1%	.3%	100.0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	5.906(a)	6	.434
Likelihood Ratio	6.480	6	.372
Linear-by-Linear Association	.317	1	.573
N of Valid Cases	720		

a 4 cells (33.3%) have expected count less than 5. The minimum expected count is .29.

## TO WHAT EXTENT : ALCOHOL USE

# CHANGED-LAST 12 MONTHS \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
TO WHAT EXTENT : ALCOHOL USE CHANGED-LAST 12 MONTHS	INCREASED	Count	47	136	0	183
		Expected Count	57.7	124.8	.5	183.0
		% within TO WHAT EXTENT : ALCOHOL USE CHANGED-LAST 12 MONTHS	25.7%	74.3%	.0%	100.0%
		% within Gender	20.7%	27.7%	.0%	25.4%
		% of Total	6.5%	18.9%	.0%	25.4%
	ABOUT THE SAME	Count	99	182	2	283
		Expected Count	89.2	193.0	.8	283.0
		% within TO WHAT EXTENT : ALCOHOL USE CHANGED-LAST 12 MONTHS	35.0%	64.3%	.7%	100.0%
		% within Gender	43.6%	37.1%	100.0%	39.3%
		% of Total	13.8%	25.3%	.3%	39.3%
	DECREASED	Count	61	125	0	186
		Expected Count	58.6	126.8	.5	186.0
		% within TO WHAT EXTENT : ALCOHOL USE CHANGED-LAST 12 MONTHS	32.8%	67.2%	.0%	100.0%
		% within Gender	26.9%	25.5%	.0%	25.8%
		% of Total	8.5%	17.4%	.0%	25.8%
	NOT USED	Count	20	48	0	68
		Expected Count	21.4	46.4	.2	68.0
		% within TO WHAT EXTENT : ALCOHOL USE CHANGED-LAST 12 MONTHS	29.4%	70.6%	.0%	100.0%
		% within Gender	8.8%	9.8%	.0%	9.4%
		% of Total	2.8%	6.7%	.0%	9.4%
Total	Count	227	491	2	720	
	Expected Count	227.0	491.0	2.0	720.0	
	% within TO WHAT EXTENT : ALCOHOL USE CHANGED-LAST 12 MONTHS	31.5%	68.2%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	

	% of Total	31.5%	68.2%	.3%	100.0%
--	------------	-------	-------	-----	--------

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	8.049(a)	6	.235
Likelihood Ratio	8.778	6	.186
Linear-by-Linear Association	.823	1	.364
N of Valid Cases	720		

a 4 cells (33.3%) have expected count less than 5. The minimum expected count is .19.

# TO WHAT EXTENT :DRUG USE CHANGED- LAST 12 MONTHS \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
TO WHAT EXTENT :DRUG USE CHANGED-LAST 12 MONTHS	INCREASED	Count	13	33	0	46
		Expected Count	14.5	31.4	.1	46.0
		% within TO WHAT EXTENT :DRUG USE CHANGED-LAST 12 MONTHS	28.3%	71.7%	.0%	100.0%
		% within Gender	5.8%	6.8%	.0%	6.4%
		% of Total	1.8%	4.6%	.0%	6.4%
	ABOUT THE SAME	Count	26	50	0	76
		Expected Count	23.9	51.8	.2	76.0
		% within TO WHAT EXTENT :DRUG USE CHANGED-LAST 12 MONTHS	34.2%	65.8%	.0%	100.0%
		% within Gender	11.6%	10.3%	.0%	10.6%
		% of Total	3.6%	7.0%	.0%	10.6%
	DECREASED	Count	47	67	0	114
		Expected Count	35.9	77.8	.3	114.0
		% within TO WHAT EXTENT :DRUG USE CHANGED-LAST 12 MONTHS	41.2%	58.8%	.0%	100.0%
		% within Gender	20.9%	13.8%	.0%	16.0%
		% of Total	6.6%	9.4%	.0%	16.0%
		Count	139	337	2	478
		Expected Count	150.6	326.0	1.3	478.0

		% within TO WHAT EXTENT :DRUG USE CHANGED-LAST 12 MONTHS	29.1%	70.5%	.4%	100.0%
		% within Gender	61.8%	69.2%	100.0%	66.9%
		% of Total	19.5%	47.2%	.3%	66.9%
Total		Count	225	487	2	714
		Expected Count	225.0	487.0	2.0	714.0
		% within TO WHAT EXTENT :DRUG USE CHANGED-LAST 12 MONTHS	31.5%	68.2%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.5%	68.2%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	7.636(a)	6	.266
Likelihood Ratio	8.043	6	.235
Linear-by-Linear Association	1.049	1	.306
N of Valid Cases	714		

a 4 cells (33.3%) have expected count less than 5. The minimum expected count is .13.

## RISK -TRYING MARIJUANA ONCE OR TWICE \* Gender

## Crosstab

		Gender			Total	
		Male	Female	Unknown		
NO RISK		Count	120	203	1	324
		Expected Count	102.6	220.5	.9	324.0
		% within RISK -TRYING MARIJUANA ONCE OR TWICE	37.0%	62.7%	.3%	100.0%
		% within Gender	52.6%	41.4%	50.0%	45.0%
		% of Total	16.7%	28.2%	.1%	45.0%
	SLIGHT RISK	Count	72	174	1	247
		Expected Count	78.2	168.1	.7	247.0
		% within RISK -TRYING MARIJUANA ONCE OR TWICE	29.1%	70.4%	.4%	100.0%
		% within Gender	31.6%	35.5%	50.0%	34.3%

RISK -TRYING MARIJUANA ONCE OR TWICE	MODERATE RISK	% of Total	10.0%	24.2%	.1%	34.3%	
		Count	19	47	0	66	
		Expected Count	20.9	44.9	.2	66.0	
		% within RISK -TRYING MARIJUANA ONCE OR TWICE	28.8%	71.2%	.0%	100.0%	
		% within Gender	8.3%	9.6%	.0%	9.2%	
		% of Total	2.6%	6.5%	.0%	9.2%	
	GREAT RISK	Count	9	43	0	52	
		Expected Count	16.5	35.4	.1	52.0	
		% within RISK -TRYING MARIJUANA ONCE OR TWICE	17.3%	82.7%	.0%	100.0%	
		% within Gender	3.9%	8.8%	.0%	7.2%	
		% of Total	1.3%	6.0%	.0%	7.2%	
	CANNOT SAY	Count	8	23	0	31	
		Expected Count	9.8	21.1	.1	31.0	
		% within RISK -TRYING MARIJUANA ONCE OR TWICE	25.8%	74.2%	.0%	100.0%	
		% within Gender	3.5%	4.7%	.0%	4.3%	
		% of Total	1.1%	3.2%	.0%	4.3%	
	Total		Count	228	490	2	720
			Expected Count	228.0	490.0	2.0	720.0
			% within RISK -TRYING MARIJUANA ONCE OR TWICE	31.7%	68.1%	.3%	100.0%
			% within Gender	100.0%	100.0%	100.0%	100.0%
			% of Total	31.7%	68.1%	.3%	100.0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	11.410(a)	8	.180
Likelihood Ratio	12.290	8	.139
Linear-by-Linear Association	7.827	1	.005
N of Valid Cases	720		

a 5 cells (33.3%) have expected count less than 5. The minimum expected count is .09.

## RISK -SMOKING MARIJUANA OCCASIONALLY \* Gender

#### Crosstab

		Gender			Total	
		Male	Female	Unknown		
RISK -SMOKING MARIJUANA OCCASIONALLY	NO RISK	Count	62	75	1	138
		Expected Count	43.6	94.1	.4	138.0
		% within RISK -SMOKING MARIJUANA OCCASIONALLY	44.9%	54.3%	.7%	100.0%
		% within Gender	27.4%	15.4%	50.0%	19.3%
		% of Total	8.7%	10.5%	.1%	19.3%
	SLIGHT RISK	Count	74	150	0	224
		Expected Count	70.7	152.7	.6	224.0
		% within RISK -SMOKING MARIJUANA OCCASIONALLY	33.0%	67.0%	.0%	100.0%
		% within Gender	32.7%	30.7%	.0%	31.3%
		% of Total	10.3%	20.9%	.0%	31.3%
	MODERATE RISK	Count	60	174	1	235
		Expected Count	74.2	160.2	.7	235.0
		% within RISK -SMOKING MARIJUANA OCCASIONALLY	25.5%	74.0%	.4%	100.0%
		% within Gender	26.5%	35.7%	50.0%	32.8%
		% of Total	8.4%	24.3%	.1%	32.8%
	GREAT RISK	Count	22	65	0	87
		Expected Count	27.5	59.3	.2	87.0
		% within RISK -SMOKING MARIJUANA OCCASIONALLY	25.3%	74.7%	.0%	100.0%
		% within Gender	9.7%	13.3%	.0%	12.2%
		% of Total	3.1%	9.1%	.0%	12.2%
	CANNOT SAY	Count	8	24	0	32
		Expected Count	10.1	21.8	.1	32.0
		% within RISK -SMOKING MARIJUANA OCCASIONALLY	25.0%	75.0%	.0%	100.0%
		% within Gender	3.5%	4.9%	.0%	4.5%
		% of Total	1.1%	3.4%	.0%	4.5%
Total		Count	226	488	2	716
		Expected Count	226.0	488.0	2.0	716.0
		% within RISK -SMOKING MARIJUANA OCCASIONALLY	31.6%	68.2%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.6%	68.2%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	20.181(a)	8	.010
Likelihood Ratio	20.435	8	.009
Linear-by-Linear Association	13.143	1	.000
N of Valid Cases	716		

a 5 cells (33.3%) have expected count less than 5. The minimum expected count is .09.

# RISK -SMOKING MARIJUANA REGULARLY

## \* Gender

## Crosstab

		Gender			Total
		Male	Female	Unknown	
RISK -SMOKING MARIJUANA REGULARLY	NO RISK	Count	25	28	1 54
		Expected Count	17.1	36.8	.1 54.0
		% within RISK -SMOKING MARIJUANA REGULARLY	46.3%	51.9%	1.9% 100.0%
		% within Gender	11.0%	5.7%	50.0% 7.5%
		% of Total	3.5%	3.9%	.1% 7.5%
	SLIGHT RISK	Count	49	77	0 126
		Expected Count	39.8	85.8	.3 126.0
		% within RISK -SMOKING MARIJUANA REGULARLY	38.9%	61.1%	.0% 100.0%
		% within Gender	21.5%	15.7%	.0% 17.5%
		% of Total	6.8%	10.7%	.0% 17.5%
	MODERATE RISK	Count	58	115	0 173
		Expected Count	54.7	117.8	.5 173.0
		% within RISK -SMOKING MARIJUANA REGULARLY	33.5%	66.5%	.0% 100.0%
		% within Gender	25.4%	23.4%	.0% 24.0%
		% of Total	8.0%	16.0%	.0% 24.0%
	GREAT RISK	Count	92	254	1 347
		Expected Count	109.7	236.3	1.0 347.0
		% within RISK -SMOKING MARIJUANA REGULARLY	26.5%	73.2%	.3% 100.0%
		% within Gender	40.4%	51.7%	50.0% 48.1%
		% of Total	12.8%	35.2%	.1% 48.1%
		Count	4	17	0 21
		Expected Count	6.6	14.3	.1 21.0

	CANNOT SAY	% within RISK -SMOKING MARIJUANA REGULARLY	19.0%	81.0%	.0%	100.0%
		% within Gender	1.8%	3.5%	.0%	2.9%
		% of Total	.6%	2.4%	.0%	2.9%
Total	Count	228	491	2	721	
	Expected Count	228.0	491.0	2.0	721.0	
	% within RISK -SMOKING MARIJUANA REGULARLY	31.6%	68.1%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	31.6%	68.1%	.3%	100.0%	

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	20.507(a)	8	.009
Likelihood Ratio	18.558	8	.017
Linear-by-Linear Association	13.273	1	.000
N of Valid Cases	721		

a 5 cells (33.3%) have expected count less than 5. The minimum expected count is .06.

## RISK -TRYING COCAINE ONCE OR TWICE \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
	NO RISK	Count	21	23	0	44
		Expected Count	13.9	30.0	.1	44.0
		% within RISK -TRYING COCAINE ONCE OR TWICE	47.7%	52.3%	.0%	100.0%
		% within Gender	9.3%	4.7%	.0%	6.1%
		% of Total	2.9%	3.2%	.0%	6.1%
	SLIGHT RISK	Count	46	97	0	143
		Expected Count	45.1	97.5	.4	143.0
		% within RISK -TRYING COCAINE ONCE OR TWICE	32.2%	67.8%	.0%	100.0%
		% within Gender	20.3%	19.8%	.0%	19.9%
		% of Total	6.4%	13.5%	.0%	19.9%
RISK -TRYING COCAINE	MODERATE	Count	59	128	0	187
		Expected Count	59.0	127.4	.5	187.0
		% within RISK -TRYING	31.6%	68.4%	.0%	100.0%

ONCE OR TWICE	RISK	COCAINE ONCE OR TWICE				
		% within Gender		26.0%	26.1%	.0% 26.0%
		% of Total		8.2%	17.8%	.0% 26.0%
	GREAT RISK	Count	92	218	2	312
		Expected Count	98.5	212.6	.9	312.0
		% within RISK -TRYING COCAINE ONCE OR TWICE		29.5%	69.9%	.6% 100.0%
		% within Gender		40.5%	44.5%	100.0% 43.4%
		% of Total		12.8%	30.3%	.3% 43.4%
Total	CANNOT SAY	Count	9	24	0	33
		Expected Count	10.4	22.5	.1	33.0
		% within RISK -TRYING COCAINE ONCE OR TWICE		27.3%	72.7%	.0% 100.0%
		% within Gender		4.0%	4.9%	.0% 4.6%
		% of Total		1.3%	3.3%	.0% 4.6%
		Count	227	490	2	719
		Expected Count	227.0	490.0	2.0	719.0
		% within RISK -TRYING COCAINE ONCE OR TWICE		31.6%	68.2%	.3% 100.0%
		% within Gender		100.0%	100.0%	100.0% 100.0%
		% of Total		31.6%	68.2%	.3% 100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	8.754(a)	8	.363
Likelihood Ratio	9.146	8	.330
Linear-by-Linear Association	4.176	1	.041
N of Valid Cases	719		

a 5 cells (33.3%) have expected count less than 5. The minimum expected count is .09.

## RISK -TAKING COCAINE REGULARLY \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
NO RISK	NO RISK	Count	2	0	0	2
		Expected Count	.6	1.4	.0	2.0
		% within RISK -TAKING COCAINE REGULARLY	100.0%	.0%	.0%	100.0%

RISK-TAKING COCAINE REGULARLY	SLIGHT RISK	% within Gender	.9%	.0%	.0%	.3%
		% of Total	.3%	.0%	.0%	.3%
		Count	3	8	0	11
		Expected Count	3.5	7.5	.0	11.0
		% within RISK-TAKING COCAINE REGULARLY	27.3%	72.7%	.0%	100.0%
	MODERATE RISK	% within Gender	1.3%	1.6%	.0%	1.5%
		% of Total	.4%	1.1%	.0%	1.5%
		Count	20	28	0	48
		Expected Count	15.2	32.7	.1	48.0
		% within RISK-TAKING COCAINE REGULARLY	41.7%	58.3%	.0%	100.0%
	GREAT RISK	% within Gender	8.8%	5.7%	.0%	6.7%
		% of Total	2.8%	3.9%	.0%	6.7%
		Count	194	437	2	633
		Expected Count	200.2	431.1	1.8	633.0
		% within RISK-TAKING COCAINE REGULARLY	30.6%	69.0%	.3%	100.0%
	CANNOT SAY	% within Gender	85.1%	89.0%	100.0%	87.8%
		% of Total	26.9%	60.6%	.3%	87.8%
		Count	9	18	0	27
		Expected Count	8.5	18.4	.1	27.0
		% within RISK-TAKING COCAINE REGULARLY	33.3%	66.7%	.0%	100.0%
	Total	% within Gender	3.9%	3.7%	.0%	3.7%
		% of Total	1.2%	2.5%	.0%	3.7%
		Count	228	491	2	721
		Expected Count	228.0	491.0	2.0	721.0
		% within RISK-TAKING COCAINE REGULARLY	31.6%	68.1%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.6%	68.1%	.3%	100.0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	7.206(a)	8	.515
Likelihood Ratio	7.622	8	.471
Linear-by-Linear Association	1.976	1	.160
N of Valid Cases	721		

a 8 cells (53.3%) have expected count less than 5. The minimum expected count is .01.

## RISK-TRYING LSD ONCE OR TWICE \*

# Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
RISK -TRYING LSD ONCE OR TWICE	NO RISK	Count	17	22	0	39
		Expected Count	12.4	26.5	.1	39.0
		% within RISK -TRYING LSD ONCE OR TWICE	43.6%	56.4%	.0%	100.0%
		% within Gender	7.5%	4.5%	.0%	5.5%
		% of Total	2.4%	3.1%	.0%	5.5%
	SLIGHT RISK	Count	38	65	0	103
		Expected Count	32.7	70.0	.3	103.0
		% within RISK -TRYING LSD ONCE OR TWICE	36.9%	63.1%	.0%	100.0%
		% within Gender	16.7%	13.4%	.0%	14.4%
		% of Total	5.3%	9.1%	.0%	14.4%
	MODERATE RISK	Count	62	126	0	188
		Expected Count	59.7	127.8	.5	188.0
		% within RISK -TRYING LSD ONCE OR TWICE	33.0%	67.0%	.0%	100.0%
		% within Gender	27.3%	25.9%	.0%	26.3%
		% of Total	8.7%	17.6%	.0%	26.3%
	GREAT RISK	Count	99	241	2	342
		Expected Count	108.6	232.5	1.0	342.0
		% within RISK -TRYING LSD ONCE OR TWICE	28.9%	70.5%	.6%	100.0%
		% within Gender	43.6%	49.6%	100.0%	47.8%
		% of Total	13.8%	33.7%	.3%	47.8%
	CANNOT SAY	Count	11	32	0	43
		Expected Count	13.7	29.2	.1	43.0
		% within RISK -TRYING LSD ONCE OR TWICE	25.6%	74.4%	.0%	100.0%
		% within Gender	4.8%	6.6%	.0%	6.0%
		% of Total	1.5%	4.5%	.0%	6.0%
Total		Count	227	486	2	715
		Expected Count	227.0	486.0	2.0	715.0
		% within RISK -TRYING LSD ONCE OR TWICE	31.7%	68.0%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.7%	68.0%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	7.939(a)	8	.439
Likelihood Ratio	8.590	8	.378
Linear-by-Linear Association	6.197	1	.013
N of Valid Cases	715		

a 5 cells (33.3%) have expected count less than 5. The minimum expected count is .11.

## RISK -TAKING LSD REGULARLY \* Gender

## Crosstab

		Gender			Total
		Male	Female	Unknown	
RISK -TAKING LSD REGULARLY	NO RISK	Count	3	0	0 3
		Expected Count	1.0	2.0	.0 3.0
		% within RISK -TAKING LSD REGULARLY	100.0%	.0%	.0% 100.0%
		% within Gender	1.3%	.0%	.0% .4%
		% of Total	.4%	.0%	.0% .4%
	SLIGHT RISK	Count	5	7	0 12
		Expected Count	3.8	8.1	.0 12.0
		% within RISK -TAKING LSD REGULARLY	41.7%	58.3%	.0% 100.0%
		% within Gender	2.2%	1.4%	.0% 1.7%
		% of Total	.7%	1.0%	.0% 1.7%
	MODERATE RISK	Count	20	22	0 42
		Expected Count	13.4	28.5	.1 42.0
		% within RISK -TAKING LSD REGULARLY	47.6%	52.4%	.0% 100.0%
		% within Gender	8.8%	4.6%	.0% 5.9%
		% of Total	2.8%	3.1%	.0% 5.9%
	GREAT RISK	Count	190	425	2 617
		Expected Count	197.3	418.0	1.7 617.0
		% within RISK -TAKING LSD REGULARLY	30.8%	68.9%	.3% 100.0%
		% within Gender	83.3%	88.0%	100.0% 86.5%
		% of Total	26.6%	59.6%	.3% 86.5%
	CANNOT SAY	Count	10	29	0 39
		Expected Count	12.5	26.4	.1 39.0
		% within RISK -TAKING LSD	25.6%	74.4%	.0% 100.0%

		REGULARLY			
		% within Gender	4.4%	6.0%	.0% 5.5%
		% of Total	1.4%	4.1%	.0% 5.5%
Total	Count	228	483	2	713
	Expected Count	228.0	483.0	2.0	713.0
	% within RISK -TAKING LSD REGULARLY	32.0%	67.7%	.3%	100.0%
	% within Gender	100.0%	100.0%	100.0%	100.0%
	% of Total	32.0%	67.7%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	13.000(a)	8	.112
Likelihood Ratio	13.427	8	.098
Linear-by-Linear Association	9.021	1	.003
N of Valid Cases	713		

a 8 cells (53.3%) have expected count less than 5. The minimum expected count is .01.

## RISK -TRYING AMPHETAMINES ONCE OR TWICE \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
NO RISK	NO RISK	Count	9	23	0	32
		Expected Count	10.2	21.8	.1	32.0
		% within RISK -TRYING AMPHETAMINES ONCE OR TWICE	28.1%	71.9%	.0%	100.0%
		% within Gender	4.0%	4.7%	.0%	4.5%
		% of Total	1.3%	3.2%	.0%	4.5%
	SLIGHT RISK	Count	44	88	0	132
		Expected Count	41.9	89.7	.4	132.0
		% within RISK -TRYING AMPHETAMINES ONCE OR TWICE	33.3%	66.7%	.0%	100.0%
		% within Gender	19.4%	18.1%	.0%	18.5%
		% of Total	6.2%	12.3%	.0%	18.5%
		Count	58	125	0	183
		Expected Count	58.1	124.4	.5	183.0

RISK -TRYING AMPHETAMINES ONCE OR TWICE	MODERATE RISK	% within RISK -TRYING AMPHETAMINES ONCE OR TWICE	31.7%	68.3%	.0%	100.0%	
		% within Gender	25.6%	25.7%	.0%	25.6%	
		% of Total	8.1%	17.5%	.0%	25.6%	
	GREAT RISK	Count	99	204	2	305	
		Expected Count	96.8	207.3	.9	305.0	
		% within RISK -TRYING AMPHETAMINES ONCE OR TWICE	32.5%	66.9%	.7%	100.0%	
		% within Gender	43.6%	42.0%	100.0%	42.7%	
		% of Total	13.8%	28.5%	.3%	42.7%	
	CANNOT SAY	Count	17	46	0	63	
		Expected Count	20.0	42.8	.2	63.0	
		% within RISK -TRYING AMPHETAMINES ONCE OR TWICE	27.0%	73.0%	.0%	100.0%	
		% within Gender	7.5%	9.5%	.0%	8.8%	
		% of Total	2.4%	6.4%	.0%	8.8%	
	Total		Count	227	486	2	715
			Expected Count	227.0	486.0	2.0	715.0
			% within RISK -TRYING AMPHETAMINES ONCE OR TWICE	31.7%	68.0%	.3%	100.0%
			% within Gender	100.0%	100.0%	100.0%	100.0%
			% of Total	31.7%	68.0%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	3.821(a)	8	.873
Likelihood Ratio	4.562	8	.803
Linear-by-Linear Association	.150	1	.699
N of Valid Cases	715		

a 5 cells (33.3%) have expected count less than 5. The minimum expected count is .09.

# RISK -TAKING AMPHETAMINES REGULARLY \* Gender

## Crosstab

	Count	Gender			Total
		Male	Female	Unknown	
		2	1	0	3

		Expected Count	1.0	2.0	.0	3.0
RISK -TAKING AMPHETAMINES REGULARLY	NO RISK	% within RISK -TAKING AMPHETAMINES REGULARLY	66.7%	33.3%	.0%	100.0%
		% within Gender	.9%	.2%	.0%	.4%
		% of Total	.3%	.1%	.0%	.4%
	SLIGHT RISK	Count	5	13	0	18
		Expected Count	5.7	12.2	.1	18.0
	MODERATE RISK	% within RISK -TAKING AMPHETAMINES REGULARLY	27.8%	72.2%	.0%	100.0%
		% within Gender	2.2%	2.7%	.0%	2.5%
		% of Total	.7%	1.8%	.0%	2.5%
	GREAT RISK	Count	25	41	0	66
		Expected Count	21.0	44.8	.2	66.0
	CANNOT SAY	% within RISK -TAKING AMPHETAMINES REGULARLY	37.9%	62.1%	.0%	100.0%
		% within Gender	11.0%	8.4%	.0%	9.2%
		% of Total	3.5%	5.7%	.0%	9.2%
	Total	Count	181	393	2	576
		Expected Count	183.2	391.2	1.6	576.0
		% within RISK -TAKING AMPHETAMINES REGULARLY	31.4%	68.2%	.3%	100.0%
		% within Gender	79.4%	80.7%	100.0%	80.3%
		% of Total	25.2%	54.8%	.3%	80.3%
		Count	15	39	0	54
		Expected Count	17.2	36.7	.2	54.0
		% within RISK -TAKING AMPHETAMINES REGULARLY	27.8%	72.2%	.0%	100.0%
		% within Gender	6.6%	8.0%	.0%	7.5%
		% of Total	2.1%	5.4%	.0%	7.5%
		Count	228	487	2	717
		Expected Count	228.0	487.0	2.0	717.0
		% within RISK -TAKING AMPHETAMINES REGULARLY	31.8%	67.9%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.9%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	3.855(a)	8	.870

Likelihood Ratio	4.064	8	.851
Linear-by-Linear Association	1.357	1	.244
N of Valid Cases	717		

a 7 cells (46.7%) have expected count less than 5. The minimum expected count is .01.

## RISK -TAKING ONE OR TWO DRINKS NEARLY EVERY DAY \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
RISK -TAKING ONE OR TWO DRINKS NEARLY EVERY DAY	NO RISK	Count	76	59	0	135
		Expected Count	42.7	91.9	.4	135.0
		% within RISK -TAKING ONE OR TWO DRINKS NEARLY EVERY DAY	56.3%	43.7%	.0%	100.0%
		% within Gender	33.3%	12.0%	.0%	18.7%
		% of Total	10.5%	8.2%	.0%	18.7%
	SLIGHT RISK	Count	87	156	1	244
		Expected Count	77.2	166.2	.7	244.0
		% within RISK -TAKING ONE OR TWO DRINKS NEARLY EVERY DAY	35.7%	63.9%	.4%	100.0%
		% within Gender	38.2%	31.8%	50.0%	33.8%
		% of Total	12.1%	21.6%	.1%	33.8%
	MODERATE RISK	Count	46	185	1	232
		Expected Count	73.4	158.0	.6	232.0
		% within RISK -TAKING ONE OR TWO DRINKS NEARLY EVERY DAY	19.8%	79.7%	.4%	100.0%
		% within Gender	20.2%	37.7%	50.0%	32.2%
		% of Total	6.4%	25.7%	.1%	32.2%
	GREAT RISK	Count	15	80	0	95
		Expected Count	30.0	64.7	.3	95.0
		% within RISK -TAKING ONE OR TWO DRINKS NEARLY EVERY DAY	15.8%	84.2%	.0%	100.0%
		% within Gender	6.6%	16.3%	.0%	13.2%
		% of Total	2.1%	11.1%	.0%	13.2%
		Count	4	11	0	15
		Expected Count	4.7	10.2	.0	15.0

	CANNOT SAY	% within RISK -TAKING ONE OR TWO DRINKS NEARLY EVERY DAY	26.7%	73.3%	.0%	100.0%
		% within Gender	1.8%	2.2%	.0%	2.1%
		% of Total	.6%	1.5%	.0%	2.1%
Total		Count	228	491	2	721
		Expected Count	228.0	491.0	2.0	721.0
		% within RISK -TAKING ONE OR TWO DRINKS NEARLY EVERY DAY	31.6%	68.1%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.6%	68.1%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	66.849(a)	8	.000
Likelihood Ratio	67.033	8	.000
Linear-by-Linear Association	53.739	1	.000
N of Valid Cases	721		

a 6 cells (40.0%) have expected count less than 5. The minimum expected count is .04.

## RISK -TAKING FOUR OR FIVE DRINKS NEARLY EVERY DAY \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
	NO RISK	Count	13	7	0	20
		Expected Count	6.4	13.6	.1	20.0
		% within RISK -TAKING FOUR OR FIVE DRINKS NEARLY EVERY DAY	65.0%	35.0%	.0%	100.0%
		% within Gender	5.7%	1.4%	.0%	2.8%
		% of Total	1.8%	1.0%	.0%	2.8%
	SLIGHT RISK	Count	33	28	0	61
		Expected Count	19.4	41.4	.2	61.0
		% within RISK -TAKING FOUR OR FIVE DRINKS NEARLY EVERY DAY	54.1%	45.9%	.0%	100.0%
		% within Gender	14.5%	5.7%	.0%	8.5%
		% of Total	4.6%	3.9%	.0%	8.5%
	Count		75	114	1	190

RISK-TAKING FOUR OR FIVE DRINKS NEARLY EVERY DAY	MODERATE RISK	Expected Count	60.4	129.1	.5	190.0
		% within RISK-TAKING FOUR OR FIVE DRINKS NEARLY EVERY DAY	39.5%	60.0%	.5%	100.0%
		% within Gender	32.9%	23.4%	50.0%	26.5%
		% of Total	10.5%	15.9%	.1%	26.5%
	GREAT RISK	Count	104	328	1	433
		Expected Count	137.7	294.1	1.2	433.0
		% within RISK-TAKING FOUR OR FIVE DRINKS NEARLY EVERY DAY	24.0%	75.8%	.2%	100.0%
		% within Gender	45.6%	67.4%	50.0%	60.4%
		% of Total	14.5%	45.7%	.1%	60.4%
	CANNOT SAY	Count	3	10	0	13
		Expected Count	4.1	8.8	.0	13.0
		% within RISK-TAKING FOUR OR FIVE DRINKS NEARLY EVERY DAY	23.1%	76.9%	.0%	100.0%
		% within Gender	1.3%	2.1%	.0%	1.8%
		% of Total	.4%	1.4%	.0%	1.8%
Total	Count	228	487	2	717	
	Expected Count	228.0	487.0	2.0	717.0	
	% within RISK-TAKING FOUR OR FIVE DRINKS NEARLY EVERY DAY	31.8%	67.9%	.3%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	31.8%	67.9%	.3%	100.0%	

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	42.624(a)	8	.000
Likelihood Ratio	41.237	8	.000
Linear-by-Linear Association	39.821	1	.000
N of Valid Cases	717		

a 6 cells (40.0%) have expected count less than 5. The minimum expected count is .04.

## RISK - HAVING 5 OR MORE DRINKS IN ONE SITTING \* Gender

## Crosstab

	Gender			Total
	Male	Female	Unknown	

RISK - HAVING 5 OR MORE DRINKS IN ONE SITTING	NO RISK	Count	30	21	0	51
		Expected Count	16.2	34.7	.1	51.0
		% within RISK - HAVING 5 OR MORE DRINKS IN ONE SITTING	58.8%	41.2%	.0%	100.0%
		% within Gender	13.2%	4.3%	.0%	7.1%
		% of Total	4.2%	2.9%	.0%	7.1%
	SLIGHT RISK	Count	58	123	1	182
		Expected Count	57.7	123.8	.5	182.0
		% within RISK - HAVING 5 OR MORE DRINKS IN ONE SITTING	31.9%	67.6%	.5%	100.0%
		% within Gender	25.4%	25.2%	50.0%	25.3%
		% of Total	8.1%	17.1%	.1%	25.3%
	MODERATE RISK	Count	76	135	1	212
		Expected Count	67.2	144.2	.6	212.0
		% within RISK - HAVING 5 OR MORE DRINKS IN ONE SITTING	35.8%	63.7%	.5%	100.0%
		% within Gender	33.3%	27.6%	50.0%	29.5%
		% of Total	10.6%	18.8%	.1%	29.5%
	GREAT RISK	Count	60	194	0	254
		Expected Count	80.5	172.7	.7	254.0
		% within RISK - HAVING 5 OR MORE DRINKS IN ONE SITTING	23.6%	76.4%	.0%	100.0%
		% within Gender	26.3%	39.7%	.0%	35.3%
		% of Total	8.3%	27.0%	.0%	35.3%
	CANNOT SAY	Count	4	16	0	20
		Expected Count	6.3	13.6	.1	20.0
		% within RISK - HAVING 5 OR MORE DRINKS IN ONE SITTING	20.0%	80.0%	.0%	100.0%
		% within Gender	1.8%	3.3%	.0%	2.8%
		% of Total	.6%	2.2%	.0%	2.8%
Total		Count	228	489	2	719
		Expected Count	228.0	489.0	2.0	719.0
		% within RISK - HAVING 5 OR MORE DRINKS IN ONE SITTING	31.7%	68.0%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.7%	68.0%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)

Pearson Chi-Square	29.772(a)	8	.000
Likelihood Ratio	29.497	8	.000
Linear-by-Linear Association	17.060	1	.000
N of Valid Cases	719		

a 5 cells (33.3%) have expected count less than 5. The minimum expected count is .06.

## RISK -TAKING STEROIDS FOR BODY BUILDING \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
RISK -TAKING STEROIDS FOR BODY BUILDING	NO RISK	Count	3	2	0	5
		Expected Count	1.6	3.4	.0	5.0
		% within RISK -TAKING STEROIDS FOR BODY BUILDING	60.0%	40.0%	.0%	100.0%
		% within Gender	1.3%	.4%	.0%	.7%
		% of Total	.4%	.3%	.0%	.7%
	SLIGHT RISK	Count	27	29	0	56
		Expected Count	17.8	38.1	.2	56.0
		% within RISK -TAKING STEROIDS FOR BODY BUILDING	48.2%	51.8%	.0%	100.0%
		% within Gender	11.9%	6.0%	.0%	7.8%
		% of Total	3.8%	4.1%	.0%	7.8%
	MODERATE RISK	Count	50	118	1	169
		Expected Count	53.6	114.9	.5	169.0
		% within RISK -TAKING STEROIDS FOR BODY BUILDING	29.6%	69.8%	.6%	100.0%
		% within Gender	22.0%	24.2%	50.0%	23.6%
		% of Total	7.0%	16.5%	.1%	23.6%
	GREAT RISK	Count	132	299	1	432
		Expected Count	137.0	293.8	1.2	432.0
		% within RISK -TAKING STEROIDS FOR BODY BUILDING	30.6%	69.2%	.2%	100.0%
		% within Gender	58.1%	61.4%	50.0%	60.3%
		% of Total	18.4%	41.8%	.1%	60.3%
		Count	15	39	0	54

		Expected Count	17.1	36.7	.2	54.0
	CANNOT SAY	% within RISK -TAKING STEROIDS FOR BODY BUILDING	27.8%	72.2%	.0%	100.0%
		% within Gender	6.6%	8.0%	.0%	7.5%
		% of Total	2.1%	5.4%	.0%	7.5%
Total		Count	227	487	2	716
		Expected Count	227.0	487.0	2.0	716.0
		% within RISK -TAKING STEROIDS FOR BODY BUILDING	31.7%	68.0%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.7%	68.0%	.3%	100.0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	10.764(a)	8	.215
Likelihood Ratio	10.322	8	.243
Linear-by-Linear Association	4.834	1	.028
N of Valid Cases	716		

a 7 cells (46.7%) have expected count less than 5. The minimum expected count is .01.

## RISK -CONSUME ALCOHOL PRIOR TO SEX \* Gender

#### Crosstab

			Gender			Total
			Male	Female	Unknown	
NO RISK	NO RISK	Count	25	23	0	48
		Expected Count	15.2	32.7	.1	48.0
		% within RISK -CONSUME ALCOHOL PRIOR TO SEX	52.1%	47.9%	.0%	100.0%
		% within Gender	11.1%	4.7%	.0%	6.7%
		% of Total	3.5%	3.2%	.0%	6.7%
SLIGHT RISK	SLIGHT RISK	Count	62	77	0	139
		Expected Count	43.9	94.7	.4	139.0
		% within RISK -CONSUME ALCOHOL PRIOR TO SEX	44.6%	55.4%	.0%	100.0%
		% within Gender	27.4%	15.8%	.0%	19.4%
		% of Total	8.7%	10.8%	.0%	19.4%
		Count	64	134	1	199

RISK -CONSUME ALCOHOL PRIOR TO SEX	MODERATE RISK	Expected Count	62.8	135.6	.6	199.0
		% within RISK -CONSUME ALCOHOL PRIOR TO SEX	32.2%	67.3%	.5%	100.0%
		% within Gender	28.3%	27.5%	50.0%	27.8%
		% of Total	8.9%	18.7%	.1%	27.8%
	GREAT RISK	Count	60	224	1	285
		Expected Count	90.0	194.2	.8	285.0
		% within RISK -CONSUME ALCOHOL PRIOR TO SEX	21.1%	78.6%	.4%	100.0%
		% within Gender	26.5%	45.9%	50.0%	39.8%
		% of Total	8.4%	31.3%	.1%	39.8%
	CANNOT SAY	Count	15	30	0	45
		Expected Count	14.2	30.7	.1	45.0
		% within RISK -CONSUME ALCOHOL PRIOR TO SEX	33.3%	66.7%	.0%	100.0%
		% within Gender	6.6%	6.1%	.0%	6.3%
		% of Total	2.1%	4.2%	.0%	6.3%
Total		Count	226	488	2	716
		Expected Count	226.0	488.0	2.0	716.0
		% within RISK -CONSUME ALCOHOL PRIOR TO SEX	31.6%	68.2%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.6%	68.2%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	35.788(a)	8	.000
Likelihood Ratio	36.132	8	.000
Linear-by-Linear Association	26.284	1	.000
N of Valid Cases	716		

a 5 cells (33.3%) have expected count less than 5. The minimum expected count is .13.

## RISK -REGULARLY ENGAGE IN UNPROTECTED SEX \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
		Count	14	5	0	19
		Expected Count	6.0	12.9	.1	19.0

RISK -REGULARLY ENGAGE IN UNPROTECTED SEX	NO RISK	% within RISK -REGULARLY ENGAGE IN UNPROTECTED SEX	73.7%	26.3%	.0%	100.0%
		% within Gender	6.1%	1.0%	.0%	2.6%
		% of Total	1.9%	.7%	.0%	2.6%
	SLIGHT RISK	Count	51	86	0	137
		Expected Count	43.4	93.2	.4	137.0
		% within RISK -REGULARLY ENGAGE IN UNPROTECTED SEX	37.2%	62.8%	.0%	100.0%
		% within Gender	22.4%	17.6%	.0%	19.1%
		% of Total	7.1%	12.0%	.0%	19.1%
	MODERATE RISK	Count	69	135	2	206
		Expected Count	65.3	140.1	.6	206.0
		% within RISK -REGULARLY ENGAGE IN UNPROTECTED SEX	33.5%	65.5%	1.0%	100.0%
		% within Gender	30.3%	27.6%	100.0%	28.7%
		% of Total	9.6%	18.8%	.3%	28.7%
	GREAT RISK	Count	88	250	0	338
		Expected Count	107.2	229.9	.9	338.0
		% within RISK -REGULARLY ENGAGE IN UNPROTECTED SEX	26.0%	74.0%	.0%	100.0%
		% within Gender	38.6%	51.1%	.0%	47.0%
		% of Total	12.2%	34.8%	.0%	47.0%
	CANNOT SAY	Count	6	13	0	19
		Expected Count	6.0	12.9	.1	19.0
		% within RISK -REGULARLY ENGAGE IN UNPROTECTED SEX	31.6%	68.4%	.0%	100.0%
		% within Gender	2.6%	2.7%	.0%	2.6%
		% of Total	.8%	1.8%	.0%	2.6%
Total		Count	228	489	2	719
		Expected Count	228.0	489.0	2.0	719.0
		% within RISK -REGULARLY ENGAGE IN UNPROTECTED SEX	31.7%	68.0%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.7%	68.0%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	27.848(a)	8	.001
Likelihood Ratio	26.581	8	.001

Linear-by-Linear Association	14.078	1	.000
N of Valid Cases	719		
a 5 cells (33.3%) have expected count less than 5. The minimum expected count is .05.			

# RISK -REGULARLY ENGAGE IN MULTILE PARTNER,UNPROTECTED SEX \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
RISK -REGULARLY ENGAGE IN MULTILE PARTNER,UNPROTECTED SEX	NO RISK	Count	2	0	0	2
		Expected Count	.6	1.4	.0	2.0
		% within RISK -REGULARLY ENGAGE IN MULTILE PARTNER,UNPROTECTED SEX	100.0%	.0%	.0%	100.0%
		% within Gender	.9%	.0%	.0%	.3%
		% of Total	.3%	.0%	.0%	.3%
	SLIGHT RISK	Count	4	5	0	9
		Expected Count	2.8	6.1	.0	9.0
		% within RISK -REGULARLY ENGAGE IN MULTILE PARTNER,UNPROTECTED SEX	44.4%	55.6%	.0%	100.0%
		% within Gender	1.8%	1.0%	.0%	1.2%
		% of Total	.6%	.7%	.0%	1.2%
	MODERATE RISK	Count	13	7	0	20
		Expected Count	6.3	13.6	.1	20.0
		% within RISK -REGULARLY ENGAGE IN MULTILE PARTNER,UNPROTECTED SEX	65.0%	35.0%	.0%	100.0%
		% within Gender	5.7%	1.4%	.0%	2.8%
		% of Total	1.8%	1.0%	.0%	2.8%
	GREAT RISK	Count	205	471	2	678
		Expected Count	214.4	461.7	1.9	678.0
		% within RISK -REGULARLY ENGAGE IN MULTILE PARTNER,UNPROTECTED SEX	30.2%	69.5%	.3%	100.0%
		% within Gender	89.9%	95.9%	100.0%	94.0%
		% of Total	28.4%	65.3%	.3%	94.0%
		Count	4	8	0	12

		Expected Count	3.8	8.2	.0	12.0
	CANNOT SAY	% within RISK -REGULARLY ENGAGE IN MULTILE PARTNER,UNPROTECTED SEX	33.3%	66.7%	.0%	100.0%
		% within Gender	1.8%	1.6%	.0%	1.7%
		% of Total	.6%	1.1%	.0%	1.7%
Total		Count	228	491	2	721
		Expected Count	228.0	491.0	2.0	721.0
		% within RISK -REGULARLY ENGAGE IN MULTILE PARTNER,UNPROTECTED SEX	31.6%	68.1%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.6%	68.1%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	15.999(a)	8	.042
Likelihood Ratio	15.399	8	.052
Linear-by-Linear Association	8.997	1	.003
N of Valid Cases	721		

a 9 cells (60.0%) have expected count less than 5. The minimum expected count is .01.

## DID YOU HAVE SEX WITHIN THE LAST YEAR \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
DID YOU HAVE SEX WITHIN THE LAST YEAR	YES	Count	159	367	2	528
		Expected Count	165.5	361.0	1.5	528.0
		% within DID YOU HAVE SEX WITHIN THE LAST YEAR	30.1%	69.5%	.4%	100.0%
		% within Gender	70.4%	74.4%	100.0%	73.2%
		% of Total	22.1%	50.9%	.3%	73.2%
	2	Count	67	126	0	193
		Expected Count	60.5	132.0	.5	193.0
		% within DID YOU HAVE SEX WITHIN THE LAST YEAR	34.7%	65.3%	.0%	100.0%
		% within Gender	29.6%	25.6%	.0%	26.8%
		% of Total	9.3%	17.5%	.0%	26.8%

Total	Count	226	493	2	721
	Expected Count	226.0	493.0	2.0	721.0
	% within DID YOU HAVE SEX WITHIN THE LAST YEAR	31.3%	68.4%	.3%	100.0%
	% within Gender	100.0%	100.0%	100.0%	100.0%
	% of Total	31.3%	68.4%	.3%	100.0%

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	2.054(a)	2	.358
Likelihood Ratio	2.553	2	.279
Linear-by-Linear Association	1.593	1	.207
N of Valid Cases	721		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .54.

## DID YOU DRINK ALCOHOL? \* Gender

#### Crosstab

			Gender			Total
			Male	Female	Unknown	
DID YOU DRINK ALCOHOL?	YES	Count	50	78	0	128
		Expected Count	38.6	88.9	.5	128.0
		% within DID YOU DRINK ALCOHOL?	39.1%	60.9%	.0%	100.0%
		% within Gender	30.3%	20.5%	.0%	23.4%
		% of Total	9.1%	14.3%	.0%	23.4%
	2	Count	115	302	2	419
		Expected Count	126.4	291.1	1.5	419.0
		% within DID YOU DRINK ALCOHOL?	27.4%	72.1%	.5%	100.0%
		% within Gender	69.7%	79.5%	100.0%	76.6%
		% of Total	21.0%	55.2%	.4%	76.6%
Total	Count	165	380	2	547	
	Expected Count	165.0	380.0	2.0	547.0	
	% within DID YOU DRINK ALCOHOL?	30.2%	69.5%	.4%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	30.2%	69.5%	.4%	100.0%	

#### Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	6.748(a)	2	.034
Likelihood Ratio	7.000	2	.030

Linear-by-Linear Association	6.611	1	.010
N of Valid Cases	547		
a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .47.			

## DID YOU USE OTHER DRUGS? \* Gender

Crosstab

		Gender			Total
		Male	Female	Unknown	
DID YOU USE OTHER DRUGS?	YES	Count	5	18	0 23
		Expected Count	7.0	16.0	.1 23.0
		% within DID YOU USE OTHER DRUGS?	21.7%	78.3%	.0% 100.0%
		% within Gender	3.0%	4.7%	.0% 4.2%
		% of Total	.9%	3.3%	.0% 4.2%
	2	Count	160	361	2 523
		Expected Count	158.0	363.0	1.9 523.0
		% within DID YOU USE OTHER DRUGS?	30.6%	69.0%	.4% 100.0%
		% within Gender	97.0%	95.3%	100.0% 95.8%
		% of Total	29.3%	66.1%	.4% 95.8%
Total		Count	165	379	2 546
		Expected Count	165.0	379.0	2.0 546.0
		% within DID YOU USE OTHER DRUGS?	30.2%	69.4%	.4% 100.0%
		% within Gender	100.0%	100.0%	100.0% 100.0%
		% of Total	30.2%	69.4%	.4% 100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.930(a)	2	.628
Likelihood Ratio	1.064	2	.587
Linear-by-Linear Association	.728	1	.393
N of Valid Cases	546		
a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .08.			

## Crosstabs

Notes

Output Created	16-JUN-2004 09:47:02
----------------	----------------------

Comments		
Input	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	747
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each table are based on all the cases with valid data in the specified range(s) for all variables in each table.
Syntax	CROSSTABS /TABLES=q37a q37b q37c q37d q37e q37f q37g q37h q38a q38b q38c q38d q38e q39a q39b q39c q39d q39e q39f q39g BY q05 /STATISTIC=CHISQ /CELLS= COUNT EXPECTED ROW COLUMN TOTAL.	
Resources	Dimensions Requested	2
	Cells Available	116508
	Elapsed Time	0:00:00.69

## Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
PAST 30 DAYS:REFUSED AN OFFER OF ALCOHOL * Gender	717	96.0%	30	4.0%	747	100.0%
PAST 30 DAYS:BRAGGED ABOUT ALCOHOL USE * Gender	717	96.0%	30	4.0%	747	100.0%
PAST 30 DAYS:HEARD SOMEONE BRAG ABOUT ALCOHOL USE * Gender	718	96.1%	29	3.9%	747	100.0%
PAST 30 DAYS:CARRIED A WEAPON * Gender	716	95.9%	31	4.1%	747	100.0%
PAST 30 DAYS:EXPERIENCED PEER PRESSURE,ALCOHOL * Gender	718	96.1%	29	3.9%	747	100.0%
PAST 30 DAYS:HELD A DRINK-DECOY * Gender	719	96.3%	28	3.7%	747	100.0%
PAST 30 DAYS:THOUGHT A PARTNER NOT ATTRACTIVE,DRUNK * Gender	718	96.1%	29	3.9%	747	100.0%
PAST 30 DAYS:TOLD A PARTNER NOT ATTRACTIVE,DRUNK * Gender	717	96.0%	30	4.0%	747	100.0%
I FEEL VALUED AS A PERSON * Gender	717	96.0%	30	4.0%	747	100.0%
I FEEL THAT FACULTY CARE ABOUT ME * Gender	717	96.0%	30	4.0%	747	100.0%
I HAVE A RESPONSIBILITY TO CONTRIBUTE * Gender	716	95.9%	31	4.1%	747	100.0%
MY CAMPUS ENCOURAGES ME TO HELP OTHERS * Gender	717	96.0%	30	4.0%	747	100.0%
I ABIDE BY THE UNIVERSITY POLICY CONCERNING ALCOHOL * Gender	716	95.9%	31	4.1%	747	100.0%
OTHERS : INTERRPUTS STUDYING * Gender	717	96.0%	30	4.0%	747	100.0%
OTHERS : MAKES YOU FEEL UNSAFE * Gender	717	96.0%	30	4.0%	747	100.0%
OTHERS : MESSES UP YOUR SPACE * Gender	715	95.7%	32	4.3%	747	100.0%

OTHERS : AFFECTS YOUR INVOLVEMENTS IN ORGANIZED GROUPS * Gender	710	95.0%	37	5.0%	747	100.0%
OTHERS : PREVENTS YOU FROM ENJOYING EVENTS * Gender	710	95.0%	37	5.0%	747	100.0%
OTHERS : INTERFERES IN OTHER WAYS * Gender	716	95.9%	31	4.1%	747	100.0%
OTHERS : DOESNT INTERFERE WITH MY LIFE * Gender	710	95.0%	37	5.0%	747	100.0%

## PAST 30 DAYS:REFUSED AN OFFER OF ALCOHOL \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
PAST 30 DAYS:REFUSED AN OFFER OF ALCOHOL	0 TIMES	Count	59	112	0	171
		Expected Count	54.1	116.4	.5	171.0
		% within PAST 30 DAYS:REFUSED AN OFFER OF ALCOHOL	34.5%	65.5%	.0%	100.0%
		% within Gender	26.0%	23.0%	.0%	23.8%
		% of Total	8.2%	15.6%	.0%	23.8%
	ONE TIME	Count	41	91	0	132
		Expected Count	41.8	89.8	.4	132.0
		% within PAST 30 DAYS:REFUSED AN OFFER OF ALCOHOL	31.1%	68.9%	.0%	100.0%
		% within Gender	18.1%	18.6%	.0%	18.4%
		% of Total	5.7%	12.7%	.0%	18.4%
	TWO TIMES	Count	48	116	0	164
		Expected Count	51.9	111.6	.5	164.0
		% within PAST 30 DAYS:REFUSED AN OFFER OF ALCOHOL	29.3%	70.7%	.0%	100.0%
		% within Gender	21.1%	23.8%	.0%	22.9%
		% of Total	6.7%	16.2%	.0%	22.9%
	3-5 TIMES	Count	56	115	2	173
		Expected Count	54.8	117.7	.5	173.0
		% within PAST 30 DAYS:REFUSED AN OFFER OF ALCOHOL	32.4%	66.5%	1.2%	100.0%
		% within Gender	24.7%	23.6%	100.0%	24.1%
		% of Total	7.8%	16.0%	.3%	24.1%
		Count	12	28	0	40
		Expected Count	12.7	27.2	.1	40.0

6-9 TIMES	% within PAST 30 DAYS:REFUSED AN OFFER OF ALCOHOL	30.0%	70.0%	.0%	100.0%
	% within Gender	5.3%	5.7%	.0%	5.6%
	% of Total	1.7%	3.9%	.0%	5.6%
10 OR MORE TIMES	Count	11	26	0	37
	Expected Count	11.7	25.2	.1	37.0
	% within PAST 30 DAYS:REFUSED AN OFFER OF ALCOHOL	29.7%	70.3%	.0%	100.0%
	% within Gender	4.8%	5.3%	.0%	5.2%
	% of Total	1.5%	3.6%	.0%	5.2%
Total	Count	227	488	2	717
	Expected Count	227.0	488.0	2.0	717.0
	% within PAST 30 DAYS:REFUSED AN OFFER OF ALCOHOL	31.7%	68.1%	.3%	100.0%
	% within Gender	100.0%	100.0%	100.0%	100.0%
	% of Total	31.7%	68.1%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	7.607(a)	10	.667
Likelihood Ratio	7.002	10	.725
Linear-by-Linear Association	.561	1	.454
N of Valid Cases	717		

a 6 cells (33.3%) have expected count less than 5. The minimum expected count is .10.

## PAST 30 DAYS:BRAGGED ABOUT ALCOHOL USE \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
0 TIMES	0 TIMES	Count	161	379	1	541
		Expected Count	170.5	369.0	1.5	541.0
		% within PAST 30 DAYS:BRAGGED ABOUT ALCOHOL USE	29.8%	70.1%	.2%	100.0%
		% within Gender	71.2%	77.5%	50.0%	75.5%
		% of Total	22.5%	52.9%	.1%	75.5%
	Count	25	42	0	67	

PAST 30 DAYS:BRAGGED ABOUT ALCOHOL USE	ONE TIME	Expected Count	21.1	45.7	.2	67.0
		% within PAST 30 DAYS:BRAGGED ABOUT ALCOHOL USE	37.3%	62.7%	.0%	100.0%
		% within Gender	11.1%	8.6%	.0%	9.3%
		% of Total	3.5%	5.9%	.0%	9.3%
	TWO TIMES	Count	16	36	0	52
		Expected Count	16.4	35.5	.1	52.0
		% within PAST 30 DAYS:BRAGGED ABOUT ALCOHOL USE	30.8%	69.2%	.0%	100.0%
		% within Gender	7.1%	7.4%	.0%	7.3%
		% of Total	2.2%	5.0%	.0%	7.3%
	3-5 TIMES	Count	18	22	1	41
		Expected Count	12.9	28.0	.1	41.0
		% within PAST 30 DAYS:BRAGGED ABOUT ALCOHOL USE	43.9%	53.7%	2.4%	100.0%
		% within Gender	8.0%	4.5%	50.0%	5.7%
		% of Total	2.5%	3.1%	.1%	5.7%
	6-9 TIMES	Count	3	7	0	10
		Expected Count	3.2	6.8	.0	10.0
		% within PAST 30 DAYS:BRAGGED ABOUT ALCOHOL USE	30.0%	70.0%	.0%	100.0%
		% within Gender	1.3%	1.4%	.0%	1.4%
		% of Total	.4%	1.0%	.0%	1.4%
	10 OR MORE TIMES	Count	3	3	0	6
		Expected Count	1.9	4.1	.0	6.0
		% within PAST 30 DAYS:BRAGGED ABOUT ALCOHOL USE	50.0%	50.0%	.0%	100.0%
		% within Gender	1.3%	.6%	.0%	.8%
		% of Total	.4%	.4%	.0%	.8%
	Total	Count	226	489	2	717
		Expected Count	226.0	489.0	2.0	717.0
		% within PAST 30 DAYS:BRAGGED ABOUT ALCOHOL USE	31.5%	68.2%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.5%	68.2%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	13.460(a)	10	.199

Likelihood Ratio	9.356	10	.499
Linear-by-Linear Association	2.668	1	.102
N of Valid Cases	717		
a 9 cells (50.0%) have expected count less than 5. The minimum expected count is .02.			

## PAST 30 DAYS:HEARD SOMEONE BRAG ABOUT ALCOHOL USE \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
PAST 30 DAYS:HEARD SOMEONE BRAG ABOUT ALCOHOL USE	0 TIMES	Count	26	78	0	104
		Expected Count	32.7	71.0	.3	104.0
		% within PAST 30 DAYS:HEARD SOMEONE BRAG ABOUT ALCOHOL USE	25.0%	75.0%	.0%	100.0%
		% within Gender	11.5%	15.9%	.0%	14.5%
		% of Total	3.6%	10.9%	.0%	14.5%
	ONE TIME	Count	17	40	0	57
		Expected Count	17.9	38.9	.2	57.0
		% within PAST 30 DAYS:HEARD SOMEONE BRAG ABOUT ALCOHOL USE	29.8%	70.2%	.0%	100.0%
		% within Gender	7.5%	8.2%	.0%	7.9%
		% of Total	2.4%	5.6%	.0%	7.9%
	TWO TIMES	Count	34	71	0	105
		Expected Count	33.1	71.7	.3	105.0
		% within PAST 30 DAYS:HEARD SOMEONE BRAG ABOUT ALCOHOL USE	32.4%	67.6%	.0%	100.0%
		% within Gender	15.0%	14.5%	.0%	14.6%
		% of Total	4.7%	9.9%	.0%	14.6%
	3-5 TIMES	Count	67	162	1	230
		Expected Count	72.4	157.0	.6	230.0
		% within PAST 30 DAYS:HEARD SOMEONE BRAG ABOUT ALCOHOL USE	29.1%	70.4%	.4%	100.0%
		% within Gender	29.6%	33.1%	50.0%	32.0%
		% of Total	9.3%	22.6%	.1%	32.0%
		Count	29	65	0	94
		Expected Count	29.6	64.2	.3	94.0

		% within PAST 30 DAYS:HEARD SOMEONE BRAG ABOUT ALCOHOL USE	30.9%	69.1%	.0%	100.0%
		% within Gender	12.8%	13.3%	.0%	13.1%
		% of Total	4.0%	9.1%	.0%	13.1%
Total	6-9 TIMES	Count	53	74	1	128
		Expected Count	40.3	87.4	.4	128.0
		% within PAST 30 DAYS:HEARD SOMEONE BRAG ABOUT ALCOHOL USE	41.4%	57.8%	.8%	100.0%
		% within Gender	23.5%	15.1%	50.0%	17.8%
		% of Total	7.4%	10.3%	.1%	17.8%
Total	10 OR MORE TIMES	Count	226	490	2	718
		Expected Count	226.0	490.0	2.0	718.0
		% within PAST 30 DAYS:HEARD SOMEONE BRAG ABOUT ALCOHOL USE	31.5%	68.2%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.5%	68.2%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	11.198(a)	10	.342
Likelihood Ratio	11.627	10	.311
Linear-by-Linear Association	4.684	1	.030
N of Valid Cases	718		

a 6 cells (33.3%) have expected count less than 5. The minimum expected count is .16.

## PAST 30 DAYS:CARRIED A WEAPON \*

### Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
Total	0 TIMES	Count	201	483	2	686
		Expected Count	214.6	469.5	1.9	686.0
		% within PAST 30 DAYS:CARRIED A WEAPON	29.3%	70.4%	.3%	100.0%
		% within Gender	89.7%	98.6%	100.0%	95.8%
		% of Total	28.1%	67.5%	.3%	95.8%
		Count	7	2	0	9

PAST 30 DAYS:CARRIED A WEAPON	ONE TIME	Expected Count	2.8	6.2	.0	9.0
		% within PAST 30 DAYS:CARRIED A WEAPON	77.8%	22.2%	.0%	100.0%
		% within Gender	3.1%	.4%	.0%	1.3%
		% of Total	1.0%	.3%	.0%	1.3%
	TWO TIMES	Count	5	2	0	7
		Expected Count	2.2	4.8	.0	7.0
		% within PAST 30 DAYS:CARRIED A WEAPON	71.4%	28.6%	.0%	100.0%
		% within Gender	2.2%	.4%	.0%	1.0%
		% of Total	.7%	.3%	.0%	1.0%
	3-5 TIMES	Count	2	1	0	3
		Expected Count	.9	2.1	.0	3.0
		% within PAST 30 DAYS:CARRIED A WEAPON	66.7%	33.3%	.0%	100.0%
		% within Gender	.9%	.2%	.0%	.4%
		% of Total	.3%	.1%	.0%	.4%
	6-9 TIMES	Count	0	1	0	1
		Expected Count	.3	.7	.0	1.0
		% within PAST 30 DAYS:CARRIED A WEAPON	.0%	100.0%	.0%	100.0%
		% within Gender	.0%	.2%	.0%	.1%
		% of Total	.0%	.1%	.0%	.1%
	10 OR MORE TIMES	Count	9	1	0	10
		Expected Count	3.1	6.8	.0	10.0
		% within PAST 30 DAYS:CARRIED A WEAPON	90.0%	10.0%	.0%	100.0%
		% within Gender	4.0%	.2%	.0%	1.4%
		% of Total	1.3%	.1%	.0%	1.4%
	Total	Count	224	490	2	716
		Expected Count	224.0	490.0	2.0	716.0
		% within PAST 30 DAYS:CARRIED A WEAPON	31.3%	68.4%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.3%	68.4%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	33.808(a)	10	.000
Likelihood Ratio	31.805	10	.000
Linear-by-Linear Association	24.315	1	.000
N of Valid Cases	716		

a 14 cells (77.8%) have expected count less than 5. The minimum expected count is .00.

# PAST 30 DAYS:EXPERIENCED PEER PRESSURE,ALCOHOL \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
PAST 30 DAYS:EXPERIENCED PEER PRESSURE,ALCOHOL	0 TIMES	Count	123	285	1	409
		Expected Count	129.3	278.6	1.1	409.0
		% within PAST 30 DAYS:EXPERIENCED PEER PRESSURE,ALCOHOL	30.1%	69.7%	.2%	100.0%
		% within Gender	54.2%	58.3%	50.0%	57.0%
		% of Total	17.1%	39.7%	.1%	57.0%
	ONE TIME	Count	17	64	0	81
		Expected Count	25.6	55.2	.2	81.0
		% within PAST 30 DAYS:EXPERIENCED PEER PRESSURE,ALCOHOL	21.0%	79.0%	.0%	100.0%
		% within Gender	7.5%	13.1%	.0%	11.3%
		% of Total	2.4%	8.9%	.0%	11.3%
	TWO TIMES	Count	25	60	1	86
		Expected Count	27.2	58.6	.2	86.0
		% within PAST 30 DAYS:EXPERIENCED PEER PRESSURE,ALCOHOL	29.1%	69.8%	1.2%	100.0%
		% within Gender	11.0%	12.3%	50.0%	12.0%
		% of Total	3.5%	8.4%	.1%	12.0%
	3-5 TIMES	Count	39	61	0	100
		Expected Count	31.6	68.1	.3	100.0
		% within PAST 30 DAYS:EXPERIENCED PEER PRESSURE,ALCOHOL	39.0%	61.0%	.0%	100.0%
		% within Gender	17.2%	12.5%	.0%	13.9%
		% of Total	5.4%	8.5%	.0%	13.9%
	6-9 TIMES	Count	14	9	0	23
		Expected Count	7.3	15.7	.1	23.0
		% within PAST 30 DAYS:EXPERIENCED PEER PRESSURE,ALCOHOL	60.9%	39.1%	.0%	100.0%
		% within Gender	6.2%	1.8%	.0%	3.2%
		% of Total	1.9%	1.3%	.0%	3.2%

		Count	9	10	0	19
		Expected Count	6.0	12.9	.1	19.0
	10 OR MORE TIMES	% within PAST 30 DAYS:EXPERIENCED PEER PRESSURE,ALCOHOL	47.4%	52.6%	.0%	100.0%
		% within Gender	4.0%	2.0%	.0%	2.6%
		% of Total	1.3%	1.4%	.0%	2.6%
Total		Count	227	489	2	718
		Expected Count	227.0	489.0	2.0	718.0
		% within PAST 30 DAYS:EXPERIENCED PEER PRESSURE,ALCOHOL	31.6%	68.1%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.6%	68.1%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	21.715(a)	10	.017
Likelihood Ratio	20.534	10	.025
Linear-by-Linear Association	8.408	1	.004
N of Valid Cases	718		

a 6 cells (33.3%) have expected count less than 5. The minimum expected count is .05.

## PAST 30 DAYS:HELD A DRINK-DECOY \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
0 TIMES		Count	201	423	2	626
		Expected Count	197.6	426.6	1.7	626.0
		% within PAST 30 DAYS:HELD A DRINK-DECOY	32.1%	67.6%	.3%	100.0%
		% within Gender	88.5%	86.3%	100.0%	87.1%
		% of Total	28.0%	58.8%	.3%	87.1%
ONE TIME		Count	14	38	0	52
		Expected Count	16.4	35.4	.1	52.0
		% within PAST 30 DAYS:HELD A DRINK-DECOY	26.9%	73.1%	.0%	100.0%
		% within Gender	6.2%	7.8%	.0%	7.2%
		% of Total	1.9%	5.3%	.0%	7.2%

PAST 30 DAYS:HELD A DRINK-DECOY	TWO TIMES	Count	5	17	0	22
		Expected Count	6.9	15.0	.1	22.0
		% within PAST 30 DAYS:HELD A DRINK-DECOY	22.7%	77.3%	.0%	100.0%
		% within Gender	2.2%	3.5%	.0%	3.1%
		% of Total	.7%	2.4%	.0%	3.1%
	3-5 TIMES	Count	5	11	0	16
		Expected Count	5.1	10.9	.0	16.0
		% within PAST 30 DAYS:HELD A DRINK-DECOY	31.3%	68.8%	.0%	100.0%
		% within Gender	2.2%	2.2%	.0%	2.2%
		% of Total	.7%	1.5%	.0%	2.2%
	6-9 TIMES	Count	0	1	0	1
		Expected Count	.3	.7	.0	1.0
		% within PAST 30 DAYS:HELD A DRINK-DECOY	.0%	100.0%	.0%	100.0%
		% within Gender	.0%	.2%	.0%	.1%
		% of Total	.0%	.1%	.0%	.1%
	10 OR MORE TIMES	Count	2	0	0	2
		Expected Count	.6	1.4	.0	2.0
		% within PAST 30 DAYS:HELD A DRINK-DECOY	100.0%	.0%	.0%	100.0%
		% within Gender	.9%	.0%	.0%	.3%
		% of Total	.3%	.0%	.0%	.3%
	Total	Count	227	490	2	719
		Expected Count	227.0	490.0	2.0	719.0
		% within PAST 30 DAYS:HELD A DRINK-DECOY	31.6%	68.2%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.6%	68.2%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	6.535(a)	10	.768
Likelihood Ratio	7.423	10	.685
Linear-by-Linear Association	.003	1	.958
N of Valid Cases	719		

a 10 cells (55.6%) have expected count less than 5. The minimum expected count is .00.

# PAST 30 DAYS:THOUGHT A PARTNER NOT ATTRACTIVE,DRUNK \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
PAST 30 DAYS:THOUGHT A PARTNER NOT ATTRACTIVE,DRUNK	0 TIMES	Count	166	353	1	520
		Expected Count	163.7	354.9	1.4	520.0
		% within PAST 30 DAYS:THOUGHT A PARTNER NOT ATTRACTIVE,DRUNK	31.9%	67.9%	.2%	100.0%
		% within Gender	73.5%	72.0%	50.0%	72.4%
		% of Total	23.1%	49.2%	.1%	72.4%
	ONE TIME	Count	18	49	0	67
		Expected Count	21.1	45.7	.2	67.0
		% within PAST 30 DAYS:THOUGHT A PARTNER NOT ATTRACTIVE,DRUNK	26.9%	73.1%	.0%	100.0%
		% within Gender	8.0%	10.0%	.0%	9.3%
		% of Total	2.5%	6.8%	.0%	9.3%
	TWO TIMES	Count	19	41	0	60
		Expected Count	18.9	40.9	.2	60.0
		% within PAST 30 DAYS:THOUGHT A PARTNER NOT ATTRACTIVE,DRUNK	31.7%	68.3%	.0%	100.0%
		% within Gender	8.4%	8.4%	.0%	8.4%
		% of Total	2.6%	5.7%	.0%	8.4%
	3-5 TIMES	Count	16	35	1	52
		Expected Count	16.4	35.5	.1	52.0
		% within PAST 30 DAYS:THOUGHT A PARTNER NOT ATTRACTIVE,DRUNK	30.8%	67.3%	1.9%	100.0%
		% within Gender	7.1%	7.1%	50.0%	7.2%
		% of Total	2.2%	4.9%	.1%	7.2%
	6-9 TIMES	Count	3	9	0	12
		Expected Count	3.8	8.2	.0	12.0
		% within PAST 30 DAYS:THOUGHT A PARTNER NOT ATTRACTIVE,DRUNK	25.0%	75.0%	.0%	100.0%
		% within Gender	1.3%	1.8%	.0%	1.7%
		% of Total	.4%	1.3%	.0%	1.7%
	10 OR MORE TIMES	Count	4	3	0	7
		Expected Count	2.2	4.8	.0	7.0
		% within PAST 30 DAYS:THOUGHT A PARTNER NOT ATTRACTIVE,DRUNK	57.1%	42.9%	.0%	100.0%
		% within Gender	1.8%	.6%	.0%	1.0%

	% of Total	.6%	.4%	.0%	1.0%
Total	Count	226	490	2	718
	Expected Count	226.0	490.0	2.0	718.0
	% within PAST 30 DAYS:THOUGHT A PARTNER NOT ATTRACTIVE,DRUNK	31.5%	68.2%	.3%	100.0%
	% within Gender	100.0%	100.0%	100.0%	100.0%
	% of Total	31.5%	68.2%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	8.706(a)	10	.560
Likelihood Ratio	6.077	10	.809
Linear-by-Linear Association	.000	1	.994
N of Valid Cases	718		

a 9 cells (50.0%) have expected count less than 5. The minimum expected count is .02.

## PAST 30 DAYS:TOLD A PARTNER NOT ATTRACTIVE,DRUNK \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
0 TIMES	ONE TIME	Count	192	399	2	593
		Expected Count	186.9	404.4	1.7	593.0
		% within PAST 30 DAYS:TOLD A PARTNER NOT ATTRACTIVE,DRUNK	32.4%	67.3%	.3%	100.0%
		% within Gender	85.0%	81.6%	100.0%	82.7%
		% of Total	26.8%	55.6%	.3%	82.7%
TWO TIMES	ONE TIME	Count	16	43	0	59
		Expected Count	18.6	40.2	.2	59.0
		% within PAST 30 DAYS:TOLD A PARTNER NOT ATTRACTIVE,DRUNK	27.1%	72.9%	.0%	100.0%
		% within Gender	7.1%	8.8%	.0%	8.2%
		% of Total	2.2%	6.0%	.0%	8.2%
TWO TIMES	TWO TIMES	Count	11	25	0	36
		Expected Count	11.3	24.6	.1	36.0
		% within PAST 30 DAYS:TOLD A PARTNER NOT ATTRACTIVE,DRUNK	30.6%	69.4%	.0%	100.0%

PAST 30 DAYS:TOLD A PARTNER NOT ATTRACTIVE,DRUNK	3-5 TIMES	% within Gender	4.9%	5.1%	.0%	5.0%
		% of Total	1.5%	3.5%	.0%	5.0%
		Count	4	19	0	23
		Expected Count	7.2	15.7	.1	23.0
		% within PAST 30 DAYS:TOLD A PARTNER NOT ATTRACTIVE,DRUNK	17.4%	82.6%	.0%	100.0%
	6-9 TIMES	% within Gender	1.8%	3.9%	.0%	3.2%
		% of Total	.6%	2.6%	.0%	3.2%
		Count	2	3	0	5
		Expected Count	1.6	3.4	.0	5.0
		% within PAST 30 DAYS:TOLD A PARTNER NOT ATTRACTIVE,DRUNK	40.0%	60.0%	.0%	100.0%
	10 OR MORE TIMES	% within Gender	.9%	.6%	.0%	.7%
		% of Total	.3%	.4%	.0%	.7%
		Count	1	0	0	1
		Expected Count	.3	.7	.0	1.0
		% within PAST 30 DAYS:TOLD A PARTNER NOT ATTRACTIVE,DRUNK	100.0%	.0%	.0%	100.0%
	Total	% within Gender	.4%	.0%	.0%	.1%
		% of Total	.1%	.0%	.0%	.1%
		Count	226	489	2	717
		Expected Count	226.0	489.0	2.0	717.0
		% within PAST 30 DAYS:TOLD A PARTNER NOT ATTRACTIVE,DRUNK	31.5%	68.2%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.5%	68.2%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	5.690(a)	10	.841
Likelihood Ratio	6.412	10	.780
Linear-by-Linear Association	.542	1	.461
N of Valid Cases	717		

a 10 cells (55.6%) have expected count less than 5. The minimum expected count is .00.

## I FEEL VALUED AS A PERSON \* Gender

## Crosstab

		Gender			Total	
		Male	Female	Unknown		
I FEEL VALUED AS A PERSON	STRONGLY AGREE	Count	31	43	0	74
		Expected Count	23.4	50.4	.2	74.0
		% within I FEEL VALUED AS A PERSON	41.9%	58.1%	.0%	100.0%
		% within Gender	13.7%	8.8%	.0%	10.3%
		% of Total	4.3%	6.0%	.0%	10.3%
	AGREE	Count	108	190	1	299
		Expected Count	94.7	203.5	.8	299.0
		% within I FEEL VALUED AS A PERSON	36.1%	63.5%	.3%	100.0%
		% within Gender	47.6%	38.9%	50.0%	41.7%
		% of Total	15.1%	26.5%	.1%	41.7%
	NEUTRAL	Count	57	186	1	244
		Expected Count	77.2	166.1	.7	244.0
		% within I FEEL VALUED AS A PERSON	23.4%	76.2%	.4%	100.0%
		% within Gender	25.1%	38.1%	50.0%	34.0%
		% of Total	7.9%	25.9%	.1%	34.0%
	DISAGREE	Count	18	44	0	62
		Expected Count	19.6	42.2	.2	62.0
		% within I FEEL VALUED AS A PERSON	29.0%	71.0%	.0%	100.0%
		% within Gender	7.9%	9.0%	.0%	8.6%
		% of Total	2.5%	6.1%	.0%	8.6%
	STRONGLY DISAGREE	Count	8	15	0	23
		Expected Count	7.3	15.7	.1	23.0
		% within I FEEL VALUED AS A PERSON	34.8%	65.2%	.0%	100.0%
		% within Gender	3.5%	3.1%	.0%	3.2%
		% of Total	1.1%	2.1%	.0%	3.2%
	DONT KNOW	Count	5	10	0	15
		Expected Count	4.7	10.2	.0	15.0
		% within I FEEL VALUED AS A PERSON	33.3%	66.7%	.0%	100.0%
		% within Gender	2.2%	2.0%	.0%	2.1%
		% of Total	.7%	1.4%	.0%	2.1%
Total		Count	227	488	2	717
		Expected Count	227.0	488.0	2.0	717.0
		% within I FEEL VALUED AS A PERSON	31.7%	68.1%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%

	% of Total	31.7%	68.1%	.3%	100.0%
--	------------	-------	-------	-----	--------

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	14.996(a)	10	.132
Likelihood Ratio	15.673	10	.109
Linear-by-Linear Association	4.556	1	.033
N of Valid Cases	717		

a 7 cells (38.9%) have expected count less than 5. The minimum expected count is .04.

# I FEEL THAT FACULTY CARE ABOUT ME \*

## Gender

## Crosstab

		Gender			Total
		Male	Female	Unknown	
I FEEL THAT FACULTY CARE ABOUT ME	STRONGLY AGREE	Count	38	54	0 92
		Expected Count	29.1	62.6	.3 92.0
		% within I FEEL THAT FACULTY CARE ABOUT ME	41.3%	58.7%	.0% 100.0%
		% within Gender	16.7%	11.1%	.0% 12.8%
		% of Total	5.3%	7.5%	.0% 12.8%
	AGREE	Count	103	205	0 308
		Expected Count	97.5	209.6	.9 308.0
		% within I FEEL THAT FACULTY CARE ABOUT ME	33.4%	66.6%	.0% 100.0%
		% within Gender	45.4%	42.0%	.0% 43.0%
		% of Total	14.4%	28.6%	.0% 43.0%
	NEUTRAL	Count	62	143	1 206
		Expected Count	65.2	140.2	.6 206.0
		% within I FEEL THAT FACULTY CARE ABOUT ME	30.1%	69.4%	.5% 100.0%
		% within Gender	27.3%	29.3%	50.0% 28.7%
		% of Total	8.6%	19.9%	.1% 28.7%
	DISAGREE	Count	17	60	1 78
		Expected Count	24.7	53.1	.2 78.0
		% within I FEEL THAT FACULTY CARE ABOUT ME	21.8%	76.9%	1.3% 100.0%
		% within Gender	7.5%	12.3%	50.0% 10.9%
		% of Total	2.4%	8.4%	.1% 10.9%

		Count	6	17	0	23
		Expected Count	7.3	15.7	.1	23.0
	STRONGLY DISAGREE	% within I FEEL THAT FACULTY CARE ABOUT ME	26.1%	73.9%	.0%	100.0%
		% within Gender	2.6%	3.5%	.0%	3.2%
		% of Total	.8%	2.4%	.0%	3.2%
		Count	1	9	0	10
		Expected Count	3.2	6.8	.0	10.0
	DONT KNOW	% within I FEEL THAT FACULTY CARE ABOUT ME	10.0%	90.0%	.0%	100.0%
		% within Gender	.4%	1.8%	.0%	1.4%
		% of Total	.1%	1.3%	.0%	1.4%
Total		Count	227	488	2	717
		Expected Count	227.0	488.0	2.0	717.0
		% within I FEEL THAT FACULTY CARE ABOUT ME	31.7%	68.1%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.7%	68.1%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	14.679(a)	10	.144
Likelihood Ratio	15.069	10	.130
Linear-by-Linear Association	10.234	1	.001
N of Valid Cases	717		

a 7 cells (38.9%) have expected count less than 5. The minimum expected count is .03.

# I HAVE A RESPONSIBILITY TO CONTRIBUTE \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
	STRONGLY AGREE	Count	32	51	0	83
		Expected Count	26.2	56.6	.2	83.0
		% within I HAVE A RESPONSIBILITY TO CONTRIBUTE	38.6%	61.4%	.0%	100.0%
		% within Gender	14.2%	10.5%	.0%	11.6%
		% of Total	4.5%	7.1%	.0%	11.6%

I HAVE A RESPONSIBILITY TO CONTRIBUTE	AGREE	Count	99	228	0	327
		Expected Count	103.2	222.9	.9	327.0
		% within I HAVE A RESPONSIBILITY TO CONTRIBUTE	30.3%	69.7%	.0%	100.0%
		% within Gender	43.8%	46.7%	.0%	45.7%
		% of Total	13.8%	31.8%	.0%	45.7%
	NEUTRAL	Count	64	165	2	231
		Expected Count	72.9	157.4	.6	231.0
		% within I HAVE A RESPONSIBILITY TO CONTRIBUTE	27.7%	71.4%	.9%	100.0%
		% within Gender	28.3%	33.8%	100.0%	32.3%
		% of Total	8.9%	23.0%	.3%	32.3%
	DISAGREE	Count	18	33	0	51
		Expected Count	16.1	34.8	.1	51.0
		% within I HAVE A RESPONSIBILITY TO CONTRIBUTE	35.3%	64.7%	.0%	100.0%
		% within Gender	8.0%	6.8%	.0%	7.1%
		% of Total	2.5%	4.6%	.0%	7.1%
	STRONGLY DISAGREE	Count	6	5	0	11
		Expected Count	3.5	7.5	.0	11.0
		% within I HAVE A RESPONSIBILITY TO CONTRIBUTE	54.5%	45.5%	.0%	100.0%
		% within Gender	2.7%	1.0%	.0%	1.5%
		% of Total	.8%	.7%	.0%	1.5%
	DONT KNOW	Count	7	6	0	13
		Expected Count	4.1	8.9	.0	13.0
		% within I HAVE A RESPONSIBILITY TO CONTRIBUTE	53.8%	46.2%	.0%	100.0%
		% within Gender	3.1%	1.2%	.0%	1.8%
		% of Total	1.0%	.8%	.0%	1.8%
Total		Count	226	488	2	716
		Expected Count	226.0	488.0	2.0	716.0
		% within I HAVE A RESPONSIBILITY TO CONTRIBUTE	31.6%	68.2%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.6%	68.2%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)

Pearson Chi-Square	13.729(a)	10	.186
Likelihood Ratio	13.563	10	.194
Linear-by-Linear Association	.351	1	.554
N of Valid Cases	716		

a 8 cells (44.4%) have expected count less than 5. The minimum expected count is .03.

## MY CAMPUS ENCOURAGES ME TO HELP OTHERS \* Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
MY CAMPUS ENCOURAGES ME TO HELP OTHERS	STRONGLY AGREE	Count	16	32	0	48
		Expected Count	15.2	32.7	.1	48.0
		% within MY CAMPUS ENCOURAGES ME TO HELP OTHERS	33.3%	66.7%	.0%	100.0%
		% within Gender	7.0%	6.6%	.0%	6.7%
		% of Total	2.2%	4.5%	.0%	6.7%
	AGREE	Count	83	160	0	243
		Expected Count	76.9	165.4	.7	243.0
		% within MY CAMPUS ENCOURAGES ME TO HELP OTHERS	34.2%	65.8%	.0%	100.0%
		% within Gender	36.6%	32.8%	.0%	33.9%
		% of Total	11.6%	22.3%	.0%	33.9%
	NEUTRAL	Count	80	195	1	276
		Expected Count	87.4	187.8	.8	276.0
		% within MY CAMPUS ENCOURAGES ME TO HELP OTHERS	29.0%	70.7%	.4%	100.0%
		% within Gender	35.2%	40.0%	50.0%	38.5%
		% of Total	11.2%	27.2%	.1%	38.5%
	DISAGREE	Count	33	70	1	104
		Expected Count	32.9	70.8	.3	104.0
		% within MY CAMPUS ENCOURAGES ME TO HELP OTHERS	31.7%	67.3%	1.0%	100.0%
		% within Gender	14.5%	14.3%	50.0%	14.5%
		% of Total	4.6%	9.8%	.1%	14.5%
		Count	11	15	0	26

		Expected Count	8.2	17.7	.1	26.0
	STRONGLY DISAGREE	% within MY CAMPUS ENCOURAGES ME TO HELP OTHERS	42.3%	57.7%	.0%	100.0%
		% within Gender	4.8%	3.1%	.0%	3.6%
		% of Total	1.5%	2.1%	.0%	3.6%
		Count	4	16	0	20
	DONT KNOW	Expected Count	6.3	13.6	.1	20.0
		% within MY CAMPUS ENCOURAGES ME TO HELP OTHERS	20.0%	80.0%	.0%	100.0%
		% within Gender	1.8%	3.3%	.0%	2.8%
		% of Total	.6%	2.2%	.0%	2.8%
		Count	227	488	2	717
Total		Expected Count	227.0	488.0	2.0	717.0
		% within MY CAMPUS ENCOURAGES ME TO HELP OTHERS	31.7%	68.1%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.7%	68.1%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	6.980(a)	10	.727
Likelihood Ratio	7.277	10	.699
Linear-by-Linear Association	.610	1	.435
N of Valid Cases	717		

a 6 cells (33.3%) have expected count less than 5. The minimum expected count is .06.

## I ABIDE BY THE UNIVERSITY POLICY CONCERNING ALCOHOL \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
		Count	45	102	1	148
		Expected Count	46.7	100.9	.4	148.0
	STRONGLY AGREE	% within I ABIDE BY THE UNIVERSITY POLICY CONCERNING ALCOHOL	30.4%	68.9%	.7%	100.0%
		% within Gender	19.9%	20.9%	50.0%	20.7%
		% of Total	6.3%	14.2%	.1%	20.7%

I ABIDE BY THE UNIVERSITY POLICY CONCERNING ALCOHOL	AGREE	Count	59	150	0	209
		Expected Count	66.0	142.4	.6	209.0
		% within I ABIDE BY THE UNIVERSITY POLICY CONCERNING ALCOHOL	28.2%	71.8%	.0%	100.0%
		% within Gender	26.1%	30.7%	.0%	29.2%
		% of Total	8.2%	20.9%	.0%	29.2%
	NEUTRAL	Count	46	100	1	147
		Expected Count	46.4	100.2	.4	147.0
		% within I ABIDE BY THE UNIVERSITY POLICY CONCERNING ALCOHOL	31.3%	68.0%	.7%	100.0%
		% within Gender	20.4%	20.5%	50.0%	20.5%
		% of Total	6.4%	14.0%	.1%	20.5%
	DISAGREE	Count	33	79	0	112
		Expected Count	35.4	76.3	.3	112.0
		% within I ABIDE BY THE UNIVERSITY POLICY CONCERNING ALCOHOL	29.5%	70.5%	.0%	100.0%
		% within Gender	14.6%	16.2%	.0%	15.6%
		% of Total	4.6%	11.0%	.0%	15.6%
	STRONGLY DISAGREE	Count	30	18	0	48
		Expected Count	15.2	32.7	.1	48.0
		% within I ABIDE BY THE UNIVERSITY POLICY CONCERNING ALCOHOL	62.5%	37.5%	.0%	100.0%
		% within Gender	13.3%	3.7%	.0%	6.7%
		% of Total	4.2%	2.5%	.0%	6.7%
	DONT KNOW	Count	13	39	0	52
		Expected Count	16.4	35.4	.1	52.0
		% within I ABIDE BY THE UNIVERSITY POLICY CONCERNING ALCOHOL	25.0%	75.0%	.0%	100.0%
		% within Gender	5.8%	8.0%	.0%	7.3%
		% of Total	1.8%	5.4%	.0%	7.3%
Total		Count	226	488	2	716
		Expected Count	226.0	488.0	2.0	716.0
		% within I ABIDE BY THE UNIVERSITY POLICY CONCERNING ALCOHOL	31.6%	68.2%	.3%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.6%	68.2%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)

<b>Pearson Chi-Square</b>	26.559(a)	10		.003
<b>Likelihood Ratio</b>	25.359	10		.005
<b>Linear-by-Linear Association</b>	2.483	1		.115
<b>N of Valid Cases</b>	716			

a 6 cells (33.3%) have expected count less than 5. The minimum expected count is .13.

## OTHERS : INTERRPUTS STUDYING \*

### Gender

Crosstab

			Gender			<b>Total</b>
			Male	Female	Unknown	
<b>OTHERS : INTERRPUTS STUDYING</b>	YES	Count	76	203	1	280
		Expected Count	89.0	189.8	1.2	280.0
		% within OTHERS : INTERRPUTS STUDYING	27.1%	72.5%	.4%	100.0%
		% within Gender	33.3%	41.8%	33.3%	39.1%
		% of Total	10.6%	28.3%	.1%	39.1%
	2	Count	152	283	2	437
		Expected Count	139.0	296.2	1.8	437.0
		% within OTHERS : INTERRPUTS STUDYING	34.8%	64.8%	.5%	100.0%
		% within Gender	66.7%	58.2%	66.7%	60.9%
		% of Total	21.2%	39.5%	.3%	60.9%
<b>Total</b>		Count	228	486	3	717
		Expected Count	228.0	486.0	3.0	717.0
		% within OTHERS : INTERRPUTS STUDYING	31.8%	67.8%	.4%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.8%	.4%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	4.682(a)	2	.096
<b>Likelihood Ratio</b>	4.735	2	.094
<b>Linear-by-Linear Association</b>	4.330	1	.037
<b>N of Valid Cases</b>	717		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is 1.17.

# OTHERS : MAKES YOU FEEL UNSAFE \*

## Gender

Crosstab

			Gender			Total
			Male	Female	Unknown	
OTHERS : MAKES YOU FEEL UNSAFE	YES	Count	24	102	0	126
		Expected Count	40.1	85.4	.5	126.0
		% within OTHERS : MAKES YOU FEEL UNSAFE	19.0%	81.0%	.0%	100.0%
		% within Gender	10.5%	21.0%	.0%	17.6%
		% of Total	3.3%	14.2%	.0%	17.6%
	2	Count	204	384	3	591
		Expected Count	187.9	400.6	2.5	591.0
		% within OTHERS : MAKES YOU FEEL UNSAFE	34.5%	65.0%	.5%	100.0%
		% within Gender	89.5%	79.0%	100.0%	82.4%
		% of Total	28.5%	53.6%	.4%	82.4%
Total		Count	228	486	3	717
		Expected Count	228.0	486.0	3.0	717.0
		% within OTHERS : MAKES YOU FEEL UNSAFE	31.8%	67.8%	.4%	100.0%
		% within Gender	100.0%	100.0%	100.0%	100.0%
		% of Total	31.8%	67.8%	.4%	100.0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	12.368(a)	2	.002
Likelihood Ratio	13.760	2	.001
Linear-by-Linear Association	10.379	1	.001
N of Valid Cases	717		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .53.

# OTHERS : MESSES UP YOUR SPACE \*

## Gender

Crosstab

		Gender

			Male	Female	Unknown	Total
OTHERS : MESSES UP YOUR SPACE	YES	Count	79	152	0	231
		Expected Count	73.3	156.7	1.0	231.0
		% within OTHERS : MESSES UP YOUR SPACE	34.2%	65.8%	.0%	100.0%
		% within Gender	34.8%	31.3%	.0%	32.3%
		% of Total	11.0%	21.3%	.0%	32.3%
	2	Count	148	333	3	484
		Expected Count	153.7	328.3	2.0	484.0
		% within OTHERS : MESSES UP YOUR SPACE	30.6%	68.8%	.6%	100.0%
		% within Gender	65.2%	68.7%	100.0%	67.7%
		% of Total	20.7%	46.6%	.4%	67.7%
Total	Count	227	485	3	715	
	Expected Count	227.0	485.0	3.0	715.0	
	% within OTHERS : MESSES UP YOUR SPACE	31.7%	67.8%	.4%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	31.7%	67.8%	.4%	100.0%	

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	2.285(a)	2	.319
Likelihood Ratio	3.187	2	.203
Linear-by-Linear Association	1.256	1	.262
N of Valid Cases	715		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .97.

## OTHERS : AFFECTS YOUR INVOLVEMENTS IN ORGANIZED GROUPS \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
OTHERS : AFFECTS YOUR	YES	Count	13	22	0	35
		Expected Count	11.2	23.7	.1	35.0
		% within OTHERS : AFFECTS YOUR INVOLVEMENTS IN ORGANIZED GROUPS	37.1%	62.9%	.0%	100.0%
		% within Gender	5.7%	4.6%	.0%	4.9%

INVOLVEMENTS IN ORGANIZED GROUPS	% of Total		1.8%	3.1%	.0%	4.9%
	Count		214	459	2	675
	Expected Count		215.8	457.3	1.9	675.0
	% within OTHERS : AFFECTS YOUR INVOLVEMENTS IN ORGANIZED GROUPS		31.7%	68.0%	.3%	100.0%
	% within Gender		94.3%	95.4%	100.0%	95.1%
	% of Total		30.1%	64.6%	.3%	95.1%
Total	Count		227	481	2	710
	Expected Count		227.0	481.0	2.0	710.0
	% within OTHERS : AFFECTS YOUR INVOLVEMENTS IN ORGANIZED GROUPS		32.0%	67.7%	.3%	100.0%
	% within Gender		100.0%	100.0%	100.0%	100.0%
	% of Total		32.0%	67.7%	.3%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	.542(a)	2	.763
Likelihood Ratio	.629	2	.730
Linear-by-Linear Association	.492	1	.483
N of Valid Cases	710		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .10.

## OTHERS : PREVENTS YOU FROM ENJOYING EVENTS \* Gender

## Crosstab

		Gender			Total
		Male	Female	Unknown	
OTHERS : PREVENTS YOU FROM ENJOYING EVENTS	YES	Count	22	59	0
		Expected Count	25.9	54.8	.3
		% within OTHERS : PREVENTS YOU FROM ENJOYING EVENTS	27.2%	72.8%	.0%
		% within Gender	9.7%	12.3%	.0%
		% of Total	3.1%	8.3%	.0%
	2	Count	205	421	3
		Expected Count	201.1	425.2	2.7
		% within OTHERS : PREVENTS YOU FROM ENJOYING EVENTS	32.6%	66.9%	.5%
		% within Gender	90.3%	87.7%	100.0%
		% of Total			88.6%

	% of Total	28.9%	59.3%	.4%	88.6%
Total	Count	227	480	3	710
	Expected Count	227.0	480.0	3.0	710.0
	% within OTHERS : PREVENTS YOU FROM ENJOYING EVENTS	32.0%	67.6%	.4%	100.0%
	% within Gender	100.0%	100.0%	100.0%	100.0%
	% of Total	32.0%	67.6%	.4%	100.0%

## Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)
Pearson Chi-Square	1.419(a)	2	.492
Likelihood Ratio	1.783	2	.410
Linear-by-Linear Association	.784	1	.376
N of Valid Cases	710		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is .34.

## OTHERS : INTERFERES IN OTHER WAYS \* Gender

## Crosstab

			Gender			Total
			Male	Female	Unknown	
OTHERS : INTERFERES IN OTHER WAYS	YES	Count	66	173	0	239
		Expected Count	75.8	162.2	1.0	239.0
		% within OTHERS : INTERFERES IN OTHER WAYS	27.6%	72.4%	.0%	100.0%
		% within Gender	29.1%	35.6%	.0%	33.4%
		% of Total	9.2%	24.2%	.0%	33.4%
	2	Count	161	313	3	477
		Expected Count	151.2	323.8	2.0	477.0
		% within OTHERS : INTERFERES IN OTHER WAYS	33.8%	65.6%	.6%	100.0%
		% within Gender	70.9%	64.4%	100.0%	66.6%
		% of Total	22.5%	43.7%	.4%	66.6%
Total	Count	227	486	3	716	
	Expected Count	227.0	486.0	3.0	716.0	
	% within OTHERS : INTERFERES IN OTHER WAYS	31.7%	67.9%	.4%	100.0%	
	% within Gender	100.0%	100.0%	100.0%	100.0%	
	% of Total	31.7%	67.9%	.4%	100.0%	

**Chi-Square Tests**

	Value	df	Asymp. Sig. (2-sided)
<b>Pearson Chi-Square</b>	4.469(a)	2	.107
<b>Likelihood Ratio</b>	5.436	2	.066
<b>Linear-by-Linear Association</b>	2.160	1	.142
<b>N of Valid Cases</b>	716		

a 2 cells (33.3%) have expected count less than 5. The minimum expected count is 1.00.

# **OTHERS : DOESNT INTERFERE WITH MY LIFE \* Gender**

**Crosstab**

		Count	Gender			Total
			Male	Female	Unknown	
OTHERS : DOESNT INTERFERE WITH MY LIFE	YES	Count	107	209	2	318
		Expected Count	102.1	214.5	1.3	318.0
		% within OTHERS : DOESNT INTERFERE WITH MY LIFE	33.6%	65.7%	.6%	100.0%
		% within Gender	46.9%	43.6%	66.7%	44.8%
		% of Total	15.1%	29.4%	.3%	44.8%
	2	Count	121	270	1	392
		Expected Count	125.9	264.5	1.7	392.0
		% within OTHERS : DOESNT INTERFERE WITH MY LIFE	30.9%	68.9%		